

Table 1: Participant Clinical and Demographic Characteristics.

Characteristic	FD (n=10)	CNVS or Gastroparesis (n = 42)	Scintigraphy (n = 27)	Total Sample (N=79)
Age (years; <i>median</i>) (IQR)	51 (32-64)	34 (26-44)	45 (31-56)	41 (27-52)
Female, n (%)	9 (90)	37 (88)	19 (70)	65 (82)
Ethnicity, n (%)				
Caucasian	9 (90)	31 (74)	16 (59)	56 (71)
Māori	0 (0)	3 (7)	4 (15)	7 (9)
Chinese	1 (10)	1 (2)	1 (4)	3 (4)
Pasifika	0 (0)	1 (2)	1 (4)	2 (3)
Indian	0 (0)	1 (2)	1 (4)	2 (3)
Other	0 (0)	5 (12)	4 (15)	9 (11)
BMI (kgm ²) <i>mean</i> (Std) ^a	22.6 (4.8)	24.1 (4.4)	24.0 (4.5)	23.9 (4.5)
Comorbidities, n (%)				
Diabetes	1(10)	5 (12)	1 (4)	7 (9)
Hypo/Hyper- thyroidism	1(10)	0 (0)	1 (4)	2 (3)
Cardio- Respiratory	1(10)	8 (19)	6 (22)	15 (19)
IBS	1(10)	7 (17)	5 (19)	13 (16)
Anxiety / Depression / PTSD	2 (20)	16 (38)	3 (11)	21 (27)

Table 1 continued

Table 1 Continued

Characteristic	FD (only) (n=10)	CNVS or Gastroparesis (n = 42)	Scintigraphy (n = 27)	Total Sample (n=79)
PAGI-SYM Score <i>Mdn</i> (IQR)	1.23 (1.03 - 1.73)	2.45 (1.83 – 3.30)	2.60 (1.40 - 3.45)	2.40 (1.50 - 3.30)
PAGI-QOL Score <i>Mdn</i> (IQR)	3.34 (2.74 - 4.39)	2.68 (1.82- 3.44)	3.50 (2.40 - 3.80)	2.86 (2.03 - 3.71)
GCSI Score <i>Mdn</i> (IQR)	1.71 (1.31 - 2.47)	3.17 (2.52 – 3.67)	2.83 (2.08 - 3.64)	2.83 (2.06-3.64)

Note: BMI = Body Mass Index; IBS = Irritable Bowel Syndrome; PTSD = post-traumatic stress disorder; GCSI = Gastroparesis Cardinal Symptom Index.