

Health-Related Quality of Life and psychological distress of adolescents during the COVID-19 pandemic in Geneva

Authors: Roxane Dumont^{1*} M.Sc., Viviane Richard^{1*} M.Sc., H el ene Baysson^{1,2} Ph.D., Elsa Lorthe¹ RM, Ph.D., Giovanni Piumatti^{3,4} Ph.D., Stephanie Schrempft¹ Ph.D., Ania Wisniak^{1,5} M.D., R emy P. Barbe⁶, M.D., Klara M. Posfay-Barbe⁷, M.D., M.Sc., Idris Guessous^{2,8} M.D., Ph.D., Silvia Stringhini^{1,2,10} Ph.D., on behalf of the Specchio-COVID19 study group.

- 1 Unit of Population Epidemiology, Division of Primary Care Medicine, Geneva University Hospitals, Geneva, Switzerland
 - 2 Department of Health and Community Medicine, Faculty of Medicine, University of Geneva, Geneva, Switzerland
 - 3 Institute of Public Health, Faculty of BioMedicine, Universit  della Svizzera Italiana, Lugano, Switzerland
 - 4 Fondazione Agnelli, Turin, Italy
 - 5 Institute of Global Health, Faculty of Medicine, University of Geneva, Geneva, Switzerland
 - 6 Division of Child and Adolescent Psychiatry, Department of Woman, Child, and Adolescent Medicine, Geneva University Hospitals, Switzerland
 - 7 Division of General Pediatrics, Department of Woman, Child, and Adolescent Medicine, Geneva University Hospitals, Switzerland
 - 8 Division and Department of Primary Care Medicine, Geneva University Hospitals, Geneva, Switzerland
 - 10 University Center for General Medicine and Public Health, University of Lausanne, Switzerland
- * Both authors contributed equally to this work.

Specchio-COVID19 study group

Isabelle Arm-Vernez, Andrew S Azman, Fatim Ba, Oumar Ba, Delphine Bachmann, Jean-Fran ois Balavoine, Michael Balavoine, H el ene Baysson, Lison Beigbeder, Julie Berthelot, Patrick Bleich, Ga elle Bryand, Fran ois Chappuis, Prune Collombet, Delphine Courvoisier, Alain Cudet, Carlos de Mestral Vargas, Paola D'ippolito, Richard Dubos, Roxane Dumont, Isabella Eckerle, Nacira El Merjani, Antoine Flahault, Natalie Francioli, Marion Frangville, Idris Guessous, S everine Harnal, Samia Hurst, Laurent Kaiser, Omar Kherad, Julien Lamour, Pierre Lescuyer, Fran ois L'Huissier, Fanny-Blanche Lombard, Andrea Jutta Loizeau, Elsa Lorthe, Chantal Martinez, Lucie M enard, Lakshmi Menon, Ludovic Metral-Boffod, Benjamin Meyer, Alexandre Moulin, Mayssam Nehme, Natacha No el, Francesco Pennacchio, Javier Perez-Saez, Giovanni Piumatti, Didier Pittet, Jane Portier, Klara M Posfay-Barbe, G eraldine Poulain, Caroline Pugin, Nick Pullen, Zo Francia Randrianandrasana, Aude Richard, Viviane Richard, Frederic Rinaldi, Jessica Rizzo, Khadija Samir, Claire Semaani, Silvia Stringhini, St ephanie Testini, Guillemette Violot, Nicolas Vuilleumier, Ania Wisniak, Sabine Yerly, Mar ia-Eugenia Zaballa

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Corresponding author

Silvia Stringhini

silvia.stringhini@hcuge.ch

Unité d'épidémiologie populationnelle

Rue Jean-Violette 29

1205 Genève

Switzerland

+41 22 305 58 61

Acknowledgment

We are grateful to the staff of the Unit of Population Epidemiology of the HUG Division of Primary Care Medicine as well as to all the participants whose contributions were invaluable to the study.

Funding

The Specchio-COVID19 study was funded by the Swiss Federal Office of Public Health, the General Directorate of Health of the Department of Safety, Employment and Health of the canton of Geneva, the Private Foundation of the Geneva University Hospitals, the Swiss School of Public Health (Corona Immunitas Research Program) and the Fondation des Grangettes.

Conflict of interest

None of the funding institutions participated in the conduct of the research or in the preparation of the article.

Author Contributions

The idea for the study originally came from RD, VR, HB and SS. RD, VR and HB carried out the literature search. SS, HB and EL conceptualized and designed the study. RD performed data analysis. RD and VR wrote the first draft of the manuscript. All authors have read, critically revised and approved the final version of this manuscript.

Ethics and dissemination

The Specchio-COVID19 study was approved by the Cantonal Research Ethics Commission of Geneva, Switzerland (CCER Project ID 2020-00881).

ABSTRACT

Purpose

Our objective was to assess adolescent's Health-Related Quality of Life (HRQoL) and psychological distress, from their own and their parents' perspective, and to examine associated risk factors during the COVID-19 pandemic in Geneva, Switzerland.

Methods

A random sample of adolescents, aged 14-17 years, and their families was invited to a serosurvey in November and December 2020. Adolescents' HRQoL was evaluated using the validated adolescent-reported KIDSCREEN-10 and parent-reported KINDL[®] scales. Psychological distress was assessed with self-reported sadness and loneliness, and using the KINDL[®] emotional well-being scale. Risk factors for adolescents' low HRQoL and psychological distress were identified using generalized estimating equations and both adolescents' and their parents' perceptions were compared.

Results

Among 240 adolescents, 11% had a low HRQoL, 35% reported sadness and 23% reported loneliness. Based on parents' perception, 12% of the adolescents had a low HRQoL and 16% a low emotional well-being. Being a girl (aOR=3.29; 95%CI: 1.64-6.57), increased time on social media (aOR=2.05; 95%CI: 1.08-3.88), parents' *average to poor* mood (aOR=2.81; 95%CI: 1.21-6.56) and *average to poor* household financial situation (aOR=2.30; 95%CI: 1.00-5.29) were associated with an increased risk of sadness. Mismatches between adolescents' and their parents' perception of HRQoL were more likely for girls (aOR=2.88; 95%CI: 1.54-5.41) and in households with lower family well-being (aOR=0.91; 95%CI: 0.86-0.96).

Conclusion

A meaningful proportion of adolescents experienced low well-being during the second wave of COVID-19. Adolescents living in underprivileged or distressed families seemed particularly affected. Monitoring is necessary to evaluate the long-term effects of the pandemic on adolescents.

Implications and Contribution

This study describes the psychological well-being of a population-based sample of adolescents in Geneva, Switzerland amid the COVID-19 pandemic, and identifies adolescents at risk of distress. This study provides further insight by comparing adolescents' well-being as reported by themselves and their parents.

Keywords: COVID-19, adolescents, health-related quality of life, well-being, psychological distress

Word count

Abstract: 250

Main text: 3486

1. INTRODUCTION

The COVID-19 pandemic and the measures put in place by public health authorities to contain its spread have caused significant disruptions in daily life and raised concerns for mental health of the entire population, and in particular adolescents [1]. Adolescence, defined as the developmental phase from childhood to adulthood [1,2], is characterized by important psychological and physical changes, greater vulnerability to external events, and less resources to adapt to hard circumstances [3]. During this sensitive stage of life, the importance of peer-interactions increases in parallel with a desire for greater autonomy from parents [4].

Because of increased difficulties to meet these developmental needs, adolescents may have been particularly affected by the pandemic [5], despite being at low risk of severe SARS-CoV-2 infection [6]. Educational disruptions due to school closures or quarantine, the widespread use of distance learning, the introduction of social distancing, or cancellation of extra-curricular activities combined with the general stay-at-home message, led to significant changes in daily life, less time spent with peers, and more with the family. This situation could have exacerbated tensions within the family or, represented an opportunity to strengthen family relationships [7]. These unprecedented circumstances also resulted in changes in health behaviours among adolescents such as an increase in screen time [8] and “junk food” consumption [9], together with a decrease in physical activity [10]. Adolescents may have been overly burdened by these life changes along with the worrying pandemic environment, as confirmed by a narrative review showing that adolescents’ mental health was negatively impacted by the COVID-19 pandemic situation [11]. Adolescents living in underprivileged families or whose parents present a decreased mental or health state, including a SARS-CoV-2 infection, could be particularly vulnerable as they may have less support and resources to adapt to the changes induced by the pandemic [12,13].

Sanitary restrictions and mortality due to the COVID-19 pandemic were relatively light in Switzerland compared with other countries [14], which may have resulted in a reduced impact on adolescents. After a first 10 week lockdown in spring 2020, schools remained open, while restrictions such as limitations on size of gatherings, remote work and mask wearing in public spaces from 12 years old were maintained. Inland travel was never limited; non-essential shops, sport facilities and cultural places intermittently closed according to the epidemiological situation. Yet, it is important to assess adolescents' well-being and psychological distress in this specific context to be able to internationally compare the impact of different measures. A better understanding of adolescents' well-being during this period is also useful for the design of new measures as the COVID-19 pandemic continues.

In this study, we assess adolescents' health related quality of life (HRQoL) and psychological distress, as well as their risk factors. We also compare their own and their parents' perceptions. Data comes from a population-based sample of adolescents in the canton of Geneva, Switzerland, primarily recruited for a SARS-CoV-2 serosurvey during the second COVID-19 wave.

2. METHODS

Survey design

A serological study was conducted between November 23rd and December 23rd 2020, in the general population of the canton of Geneva, Switzerland [15]. Children and adolescents aged 6 months to 17 years were randomly selected from the Swiss Federal Statistical Office's registers and were invited with their family members for a free anti-SARS-CoV-2 serology. 597 families participated (participation rate: 17%). Of these, 194 families had adolescents aged between 14

and 17 years, and were included in the current study. This age range was chosen for adolescents to be mature enough to autonomously answer the study questionnaire.

Each family designated a “referent parent” who completed comprehensive health and socio-demographic online questionnaires about themselves and their child(ren). Additionally, adolescents were asked to complete a paper questionnaire about their well-being and life habits since the start of the COVID-19 pandemic, mid-March 2020. This questionnaire was completed during the serology appointment in a separate area ensuring confidentiality, especially from parents. Adolescents were included in this analysis if the three above-mentioned questionnaires were completed (Figure 1).

Measures

HRQoL and psychological distress - adolescent's perception

HRQoL as perceived by the adolescents was evaluated over the previous week using the validated French version of the standardized KIDSCREEN-10 scale [16]. KIDSCREEN-10 evaluates HRQoL in children and adolescents aged 8 to 18 years based on 10 items, enabling a normalized score calculation from 0 to 100. The internal consistency of the overall self-reported KIDSCREEN-10 index was good ($\alpha=0.81$). Adolescents with a HRQoL score lower than one standard deviation below the study population mean were considered as having a low HRQoL [17]. The 2 items of the KIDSCREEN-10 targeting mood and feelings (sadness and loneliness) were separately used as binary proxies for psychological distress. Adolescents were considered to feel sad or lonely if they reported feeling so “quite often”, “very often” or “always” over the last week.

HRQoL and psychological distress - parent's perception

The parent's perception of their children's well-being over the previous week was based on the French version of the KINDL[®] for parents [18]. This scale measures HRQoL of children and adolescents aged from 7 to 17 years based on their parent's answers, by combining 24 items covering 6 dimensions: physical well-being, emotional well-being, self-esteem, family, friends and school. Adolescents with an overall HRQoL score lower than one standard deviation below the study population mean were considered as having a low HRQoL [17]. Overall internal consistency of the KINDL[®] was high ($\alpha=0.85$). Focusing on the distinct KINDL[®] dimensions, psychological distress was defined as an emotional well-being score lower than one standard deviation below the study population mean. This dimension showed acceptable consistency ($\alpha=0.69$).

Outcomes

The 5 outcomes of interest were HRQoL from both the adolescents' and the parents' perception, as well as self-reported sadness and loneliness, and emotional well-being reported by the referent parent.

Potential risk factors

Potential risk factors were identified based on a literature review and descriptive analyses. Age, sex, screen time (hours per day), change in time spent on social networks since the start of the pandemic (increase, decrease, no change) and anti-SARS-CoV-2 serological status of the adolescent were selected, as well as age, sex, education, self-perceived mood and anti-SARS-CoV-2 serological status of the referent parent. Values of the anti-SARS-CoV-2 serology ≥ 0.8 μmL were considered positive (Elecsys anti-SARS-CoV-2 S; Roche Diagnostics, Rotkreuz, Switzerland [19]); the serological assessment was conducted before the start of the vaccination

campaign in Switzerland. Parent education was measured with a three-level scale: lower (compulsory or no formal education), middle (secondary education), and higher (tertiary education). Self-perceived mood of the parent was defined as good if the answer to the question “In general, outside the pandemic context, how would you assess your mood ?” was “good” or “very good”, and average to poor for answers such as “average”, “poor” or “very poor”. Household level variables such as household size, density, financial situation and family well-being were also included. Household size was expressed as the number of people living in the household. Household density was defined using the household crowded index [20], calculated as the household size divided by the number of bedrooms. Households with a ratio higher than 1.5 were considered crowded. Household financial situation was considered as good if the referent parent answered that they could save money or face minor unexpected expenses, and average to poor if they selected one of the following statements: “I have to be careful with my expenses and an unexpected event could put me into financial difficulty” or “I cannot cover my needs with my income and I need external support”. Family well-being was measured using the family dimension of the KINDL[®] scale for parents ($\alpha=0.71$), which focuses on the relationship between the children and their parents [18].

Statistical analysis

After excluding participants with missing data, multivariable models were performed for each outcome with adolescents’ age, sex, anti-SARS-CoV-2 serological status, screen time and change in time spent on social network, referent parents’ age, sex, anti-SARS-CoV-2 serological status and mood, as well as household size, density and financial situation as predictors. As some of the adolescents were siblings, a generalized estimating equation (GEE) function [21] was used to correct for the familial dependency in the observations with a covariance matrix defined as “exchangeable” and tests based on sandwich-corrected robust standard errors. Multivariable

model results were reported as adjusted odds ratios (aOR) with 95% confidence intervals (95%CI).

Discrepancies between self-reported and parent-reported low HRQoL were coded as a binary variable: 1 if perceptions were different and 0 if they were similar. Risk factors of discrepancy were assessed with GEE adjusting for the above-mentioned covariates, as well as the family well-being score from the KINDL[®] scale.

Statistical significance was defined at a level of confidence of 95% and all analyses were performed with R (version 4.0.3).

3. RESULTS

The sample consisted of 240 adolescents from 194 households. Mean age was 15 years (SD 2.1 years) and 47% were females. Referent parents' mean age was 47 years (SD 9.0 years), 75% being mothers. Table 1 presents a descriptive overview of the adolescents' and their parents' socio-demographic characteristics.

Adolescent-reported health-related quality of life, sadness, and loneliness

Overall, 26 (11%) adolescents reported a low HRQoL (Table 1). Multivariable analysis showed that younger adolescents (aOR=0.84; 95%CI: 0.73-0.97) were more likely to present a low HRQoL (Table 2). Referent parents' positive anti-SARS-CoV2 serology (aOR=3.44; 95%CI: 1.3-9.09) was associated with a lower HRQoL of the adolescent, whereas there was no association with the adolescents' own serological status ($P > 0.1$). Moreover, engaging in less screen time was associated with a low HRQoL (aOR=0.91; 95%CI: 0.84-0.98), whereas an

increase in time spent on social networks since the beginning of the pandemic seemed negatively associated with HRQoL, with borderline significance (aOR=1.82; 95%CI: 0.92-3.66).

Focusing on psychological distress, 35% of the adolescents reported feeling sad during the previous week, and 23% felt alone (Table 3). Sadness was more likely among girls, compared to boys (aOR=3.29; 95%CI: 1.64-6.57), and was associated with an increase in the use of social networks (aOR=2.05; 95%CI: 1.08-3.88) and an average to poor household financial situation (aOR=2.30; IC95%: 1.00-5.29) with a borderline significance level. Sadness was also associated with the referent parent's positive anti-SARS-CoV2 serology (aOR=2.38; 95%CI: 1.10-5.14), older age (aOR=1.07; 95%CI: 1.00-1.15) or average to poor mood (aOR=2.81; 95%CI: 1.21-6.56). Female adolescents (aOR=4.15; 95%CI: 2.02-8.51) and those who increased their time spent on social networks (aOR=1.93; 95%CI: 0.96-0.3.90) were more likely to report feeling lonely, while a positive anti-SARS-CoV-2 serology was associated with less loneliness (aOR=0.20; 95%CI: 0.07-0.63) (Table 2).

Parent-reported Health-related quality of life and emotional well-being

Based on parents' perception, 28 (12%) adolescents presented a low HRQoL (Table 1). The latter was only associated with adolescents' sex, girls being more likely to be perceived as having a low HRQoL compared to boys (aOR=4.21; 95%CI: 1.76-10.10; Table 2). When focusing on psychological distress, 38 (16%) adolescents presented a low emotional well-being according to their parents (Table 1), which was associated with being female (aOR=3.03; 95%CI: 1.32-6.90), living in a crowded household (aOR=3.20; 95%CI: 1.26-8.10), and having an average to poor financial situation (aOR=2.08; 95%CI: 1.00-7.79; Table 3).

Comparison of adolescents' and parents' perceptions

Based on the dichotomous classification of the KIDSCREEN-10 and the KINDL[®], adolescents' and parents' perceptions matched in 184 (79%) of cases. However, 22 (9.5%) of the adolescents

presented a low HRQoL that did not seem identified by their parents. Misperception was more likely among girls compared to boys (aOR=2.88; 95%CI: 1.54-5.41) and in households with lower family well-being (aOR=0.91; 95%CI: 0.86-0.96; Table 4).

4. DISCUSSION

In this population-based study, we observed that a meaningful proportion of adolescents reported a low well-being HRQoL and some psychological distress during the second COVID-19 wave in Switzerland, 8 to 9 months after the start of the pandemic. Adolescents' self-reported well-being was generally corroborated by their parent's observation. Risk factors for a low well-being from the adolescents' perspective included a younger age and a positive anti-SARS-CoV-2 serology of the referent parent, while being a girl was the only risk factor for low HRQoL as perceived by parents. Girls compared to boys, and adolescents living in households with lower family well-being, were at higher risk of misperception by their parents. When looking at psychological distress, risk factors for adolescent-reported sadness or loneliness included being a girl, spending more time on social networks or living in a household with a disadvantaged financial situation, as well as characteristics of the referent parent such as older age, positive anti-SARS-CoV-2 serology and lower mood.

Compared to the Swiss pre-pandemic reference level where the KIDSCREEN-10 median corresponded to a score of 80 [16,22], adolescents presented lower HRQoL in this study with a median of 72.5 (MAD = 14.8), which may be partly explained by the impact of the pandemic. This is in line with results from other studies in China [23], Brazil [24], Germany [17], Australia [25], India [26], Spain [27] and Italy [27] which reported high levels of adolescent stress, worry and anxiety during the pandemic, likely due to restrictive measures to contain COVID-19 spread and the worrying pandemic environment. Estimates of sadness and loneliness among adolescents

were lower in the present study than what reported in other studies in China [23], India [26], Spain [27] and Italy [27]. However, most previous studies were based on convenience samples and were not population-based. Further, in Switzerland measures to contain virus spread were comparatively less strict than in other countries. For example, inland travel, small gatherings and outdoor sports were never forbidden [14,28]. Finally, during the study period sanitary measures were regularly changing in Geneva, including restrictions on extra-curricular activities and mask wearing at schools, but were never as strict as during the first pandemic wave when schools were closed and activities suspended [14]. The psychological distress estimates are thus likely to be lower than what would have been observed during the first wave, when most other studies have been conducted [23,26,27].

Lower well-being was associated with a younger age, as also found by Ravens-Sieberer et al [17] and with being a girl, as generally observed [25,29]. Parental lower mood was a risk factor for adolescent psychological distress, which is consistent with studies showing that parent's mental health directly impacts children's functioning [12]. Living in a crowded household was associated with adolescents' psychological distress, which could be explained by difficulties to maintain privacy and a personal space at home [30]. This aspect has possibly worsened since the start of the pandemic as household members were likely to spend more time at home than usual. On the opposite, household size was not associated with adolescents' well-being, consistent with other findings [31].

In accordance with warnings issued by psychologists, an increase in time spent on social networks was associated with sadness and loneliness [32]. Causality could be bidirectional. As a consequence of other activities being restricted, such as sport or music lessons, adolescents may have felt lonely, sad and bored, and thus have spent more time on social media. Conversely, it

may have increased their exposure to alarming and contradictory information and affected their well-being [33]. Interestingly, overall screen time was not related to psychological distress, suggesting that examining the type of online activities may be of more significance than the overall screen time [34].

Another unexpected finding was that adolescents' low HRQoL was associated with a positive anti-SARS-CoV-2 serology of the parent but not with their own serological result. In line with other results [35], this may reflect that adolescents were more worried or impacted by their parents health rather than by their own during the COVID-19 pandemic. It may also mirror the impact on adolescents of potential difficult circumstances linked with a parent infected by SARS-CoV-2, such as a severe disease or long-term effects due to long COVID [36]. Adolescents' negative anti-SARS-CoV-2 serology was associated with loneliness. A possible explanation could be that adolescents respecting social distancing measures more carefully were less infected but also felt lonelier, although this remains speculative at this stage.

An association between poor socioeconomic conditions and adolescents' psychological distress was probably already present before the COVID-19 pandemic [37]. However, its significance was likely to increase as restrictions may have resulted in fewer quiet spaces, less resources to help with school work or increased likelihood of financial instability for adolescents in general, and more particularly vulnerable ones [13]. Longitudinal studies are necessary to assess the effect of the pandemic on adolescents' mental health over time. More specifically, the impact of socioeconomic conditions should be examined, as deprivation could result in lower resilience and constitute a risk factor for persistent low well-being.

Adolescents' and parents' perceptions of adolescents' HRQoL matched in most cases. This finding is important as parents play a role in the early detection of psychological distress among their children and in care seeking [38]. Especially during the COVID-19 pandemic, parents play a key role in reducing the negative psychological impact on their children, for example through attention and supportive behaviours, such as keeping regular routines as much as possible or providing clear explanations about the situation [39]. However, 9.5% of the adolescents presented a low HRQoL that did not seem to be identified by their parents. Misperception was more likely among girls and adolescents living in families with lower well-being; these adolescents may represent a vulnerable group whose mental health issues are under-recognized.

The study presents several limitations. First, it is a cross-sectional study relying on self-reported data, without any pre-pandemic hindsight coming from the same sample. It does not allow us to firmly conclude whether the reported adolescents' low quality of life is caused by the pandemic. The sample size is quite small, which limited statistical power. Finally, though randomly invited, study participants come from a higher socioeconomic background than the Geneva population. As a consequence, estimates of low well-being and psychological distress may have been underestimated. This study also presents several strengths. First, it relies on a randomly selected population-based sample. Thanks to the family-based design, data comes from both adolescents and their parents, which enables comparisons of both perceptions. Furthermore, previous SARS-CoV-2 infection is assessed with an objective measure. Finally, the demographic and socioeconomic characteristics of the sample are assessed through a wide range of indicators, at the individual and household levels.

5. CONCLUSION

Amid the COVID-19 second wave in Geneva, a meaningful share of adolescents reported low psychological well-being, especially those living in unfavourable family environments, including crowded households, in poor financial situation or with distressed parents. Tailored measures, especially targeting these vulnerable adolescents, are necessary and could include free psychological consultations or resources for parents to react adequately to this special situation. Finally, monitoring is necessary to evaluate the long-term effects of the pandemic on adolescents' mental health and inform decision-makers as the pandemic continues. Indeed some of the impacts may be temporary thanks to the lifting of restrictions, while others may persist over time, especially among vulnerable individuals.

REFERENCES

- [1] Sawyer SM, Afifi RA, Bearinger LH, et al. Adolescence: a foundation for future health. *The Lancet* 2012;379:1630–40.
- [2] Dick B, Ferguson BJ. Health for the World's Adolescents: A Second Chance in the Second Decade. *Journal of Adolescent Health* 2015;56:3–6.
- [3] Arnett JJ. *The Oxford Handbook of Emerging Adulthood*. Oxford University Press; 2015.
- [4] American Psychological Association. *Developing adolescents: A reference for professionals*.
- [5] Patrick SW, Henkhaus LE, Zickafoose JS, et al. Well-being of Parents and Children During the COVID-19 Pandemic: A National Survey. *Pediatrics* 2020;146.
- [6] Souza TH de, Nadal JA, Nogueira RJN, et al. Clinical manifestations of children with COVID-19: A systematic review. *Pediatric Pulmonology* 2020;55:1892–9.
- [7] Donker MH, Mastrotheodoros S, Branje S. *Development of Parent-Adolescent Relationships During the COVID-19 Pandemic: The Role of Stress and Coping* 2020.
- [8] Xiang M, Zhang Z, Kuwahara K. Impact of COVID-19 pandemic on children and adolescents' lifestyle behavior larger than expected. *Prog Cardiovasc Dis* 2020;63:531–2.
- [9] Segre G, Campi R, Scarpellini F, et al. Interviewing children: the impact of the COVID-19 quarantine on children's perceived psychological distress and changes in routine. *BMC Pediatrics* 2021;21:231.
- [10] Chaffee BW, Cheng J, Couch ET, et al. Adolescents' Substance Use and Physical Activity Before and During the COVID-19 Pandemic. *JAMA Pediatrics* 2021;175:715.
- [11] Singh S, Roy D, Sinha K, et al. Impact of COVID-19 and lockdown on mental health of children and adolescents: A narrative review with recommendations. *Psychiatry Research* 2020;293:113429.
- [12] Spinelli M, Lionetti F, Pastore M, et al. Parents' Stress and Children's Psychological Problems in Families Facing the COVID-19 Outbreak in Italy. *Front Psychol* 2020;11.
- [13] Fisher J, Languilaire J-C, Lawthom R, et al. Community, work, and family in times of COVID-19. *Community, Work & Family* 2020;23:247–52.
- [14] COVID-19: Décisions des autorités. Available at: <https://www.geneve.ch/fr/actualites/dossiers-information/covid-recommandations-sanitaires-mesures-prises-ville-geneve/informations-pratiques/decisions-autorites>. Accessed July 18, 2021.
- [15] Stringhini S, Zaballa M-E, Perez-Saez J, et al. Seroprevalence of anti-SARS-CoV-2 antibodies after the second pandemic peak. *The Lancet Infectious Diseases* 2021;21:600–1.
- [16] Ravens-Sieberer U, and KIDSCREEN Group Europe. *The KIDSCREEN questionnaires: Quality of life questionnaires for children and adolescents - Handbook*. Pabst Science Publishers 2006.
- [17] Ravens-Sieberer U, Kaman A, Erhart M, et al. Impact of the COVID-19 pandemic on quality of life and mental health in children and adolescents in Germany. *Eur Child Adolesc Psychiatry* 2021.
- [18] Ravens-Sieberer U, Bullinger M. News from the KINDL-Questionnaire – A new version for adolescents. *Quality of Life Research* 1998;7, 653.
- [19] Elecsys® Anti-SARS-CoV-2 S - Immunoassay for the quantitative determination of antibodies to the SARS-CoV-2 spike protein. 2020.
- [20] WHO Housing and health guidelines. Available at: <https://www.who.int/publications-detail-redirect/9789241550376>. Accessed July 18, 2021.
- [21] Carey VJ, Ripley B. *GEE: Generalized Estimation Equation Solver*. 2019.

- [22] Rajmil L, Herdman M, Ravens-Sieberer U, et al. Socioeconomic inequalities in mental health and health-related quality of life (HRQOL) in children and adolescents from 11 European countries. *Int J Public Health* 2014;59:95–105.
- [23] Zhou S-J, Zhang L-G, Wang L-L, et al. Prevalence and socio-demographic correlates of psychological health problems in Chinese adolescents during the outbreak of COVID-19. *Eur Child Adolesc Psychiatry* 2020;29:749–58.
- [24] Garcia de Avila M, Hamamoto Filho P, Jacob F, et al. Children’s Anxiety and Factors Related to the COVID-19 Pandemic: An Exploratory Study Using the Children’s Anxiety Questionnaire and the Numerical Rating Scale. *International Journal of Environmental Research and Public Health* 2020;17.
- [25] Magson NR, Freeman JYA, Rapee RM, et al. Risk and Protective Factors for Prospective Changes in Adolescent Mental Health during the COVID-19 Pandemic. *J Youth Adolescence* 2021;50:44–57.
- [26] Saurabh K, Ranjan S. Compliance and Psychological Impact of Quarantine in Children and Adolescents due to Covid-19 Pandemic. *Indian J Pediatr* 2020:1–5.
- [27] Orgilés M, Morales A, Delvecchio E, et al. Immediate Psychological Effects of the COVID-19 Quarantine in Youth From Italy and Spain. *Front Psychol* 2020;0.
- [28] Hale T, Angrist N, Goldszmidt R, et al. A global panel database of pandemic policies (Oxford COVID-19 Government Response Tracker). *Nat Hum Behav* 2021;5:529–38.
- [29] Pigaiani Y, Zoccante L, Zocca A, et al. Adolescent Lifestyle Behaviors, Coping Strategies and Subjective Wellbeing during the COVID-19 Pandemic: An Online Student Survey. *Healthcare* 2020;8:472.
- [30] Ruiz-Tagle J, Urria I. Household overcrowding trajectories and mental well-being. *Social Science & Medicine* 2021:114051.
- [31] Okabe-Miyamoto K, Folk D, Lyubomirsky S, et al. Changes in social connection during COVID-19 social distancing: It’s not (household) size that matters, it’s who you’re with. *PLOS ONE* 2021;16:e0245009.
- [32] Davies T. Mind change: How digital technologies are leaving their mark on our brains. *New Media & Society* 2016;18:2139–41.
- [33] Keles B, McCrae N, Grealish A. A systematic review: the influence of social media on depression, anxiety and psychological distress in adolescents. *International Journal of Adolescence and Youth* 2020;25:79–93.
- [34] Boers E, Afzali M, Newton N, et al. Association of Screen Time and Depression in Adolescence. *JAMA Pediatrics* 2019;173.
- [35] Vogel M, Meigen C, Sobek C, et al. Well-being and COVID-19-related worries of German children and adolescents: A longitudinal study from pre-COVID to the end of lockdown in Spring 2020. *JCPP Advances* 2021;1:e12004.
- [36] Shah R, Ali FM, Nixon SJ, et al. Measuring the impact of COVID-19 on the quality of life of the survivors, partners and family members: a cross-sectional international online survey. *BMJ Open* 2021;11:e047680.
- [37] Ravens-Sieberer U, Erhart M, Gosch A, et al. Mental health of children and adolescents in 12 European countries-results from the European KIDSCREEN study. *Clinical Psychology & Psychotherapy* 2008;15.
- [38] Perrin E, Leslie L, Boat T. Parenting as Primary Prevention. *JAMA Pediatrics* 2016;170.
- [39] Helping Children Cope With Stress During the 2019-nCoV Outbreak. Available at: <https://resourcecentre.savethechildren.net/library/helping-children-cope-stress-during-2019-ncov-outbreak>. Accessed July 18, 2021.

Figure 1: Study flow diagram




Table 1: Characteristics of the study population

	Adolescents (N=240)	Referent parents (N=194)
	N (%)	N (%)
Age in years, mean (SD) (N=240/N=194)	15.2 (2.1)	46.5 (9.0)
Sex (N=240/N=194)		
Male	128 (53.3)	48 (24.8)
Female	112 (46.7)	146 (75.2)
Nationality (N=240/N=194)		
Swiss	188 (78.3)	144 (74.2)
Portuguese	10 (4.2)	8 (4.1)
Italian	14 (5.8)	14 (7.2)
French	13 (5.4)	9 (4.6)
Others	15 (6.3)	19 (9.9)
Ethnicity (N=240/N=194)		
European-Caucasian	194 (80.1)	165 (85.1)
Other	46 (19.9)	29 (14.9)
Marital status (N=194)		
Married or in couple	-	152 (78.4)
Separated, single, widowed	-	42 (21.6)
Household size in individuals, mean (SD) (N=194)	-	3.2 (1.1)
Education (N=194)		
Lower	-	12 (6.2)
Medium	-	62 (32.0)
Higher	-	120 (61.8)
Household financial situation (N=194)		
High	-	139 (71.6)
Average to poor	-	40 (20.6)
Does not want to answer	-	15 (7.8)
Crowded household (N=193)	-	20 (10.3)
Positive anti-SARS-CoV-2 serology (N=240/N=194)	53 (22.1)	45 (23.2)
Low self-reported HRQoL¹ (N=235)	26 (11.0)	-
Self-reported sadness (N=236)	84 (35.5)	-
Self-reported loneliness (N=237)	55 (23.2)	-
Low parent-reported HRQoL² (N=234)	28 (11.9)	-
Low parent-reported emotional well-being² (N=234)	38 (16.2)	-
Average to poor mood (N=194)	-	30 (15.5)

Results are N (%), unless stated otherwise. All variables are self-reported, except serology (see Methods). Other ethnicities include Arab, Asian, African, Indian and South-American. HRQoL stands for health-related quality of life.

¹ Based on the KIDSCREEN-10 scale

² Based on the KINDL[®] scale

Table 2: Generalized estimating equations on adolescents' self-reported Health-Related Quality of Life, HRQoL (KIDSCREEN-10) as well as reported by their parents (KINDL®).

	Adolescent-reported HRQoL (KIDSCREEN-10)				Parent-reported HRQoL (KINDL®)			
	N	High N (%)	Low N (%)	Low HRQoL OR (95%CI) ^a	N	High N (%)	Low N (%)	Low HRQoL OR (95%CI) ^b
Age of the adolescent (years)¹	235	15.4 (1.6)	13.8 (4.2)	0.84 (0.73-0.97)*		15.4 (1.2)	15.4 (1.1)	0.92 (0.63-1.35)
Age of the parent (years)¹	235	48.4 (5.1)	48.5 (7.3)	1.02 (0.92-1.13)		48.3 (5.5)	48.7 (4.4)	1.02 (0.94-1.10)
Sex of the adolescent	235				234			
Boy		113 (90.4)	12 (9.6)	1		118 (94.4)	7 (5.6)	1
Girl		96 (87.3)	14 (12.7)	1.08 (0.57-2.02)		88 (80.7)	21 (19.3)	4.21 (1.76-10.10)**
Self-perceived mood of the parent	235				234			
Good		178 (89.9)	20 (10.1)	1		178 (89.4)	21 (10.6)	1
Average to poor		31 (83.8)	6 (16.2)	1.85 (0.57-5.93)		28 (80.0)	7 (20.0)	1.57 (0.62-3.98)
Financial situation of the household	235				234			
Good		156 (90.7)	16 (9.3)	1		152 (88.4)	20 (11.6)	1
Average to poor		35 (79.5)	9 (20.5)	2.05 (0.75-5.59)		36 (83.7)	7 (16.3)	1.06 (0.56-4.59)
No answer		18 (94.7)	1 (5.3)	0.63 (0.07-5.27)		18 (94.7)	1 (5.3)	0.67 (0.08-5.49)
Household size (individuals)¹	235	3.2 (1.0)	3.8 (2.0)	1.41 (0.81-2.44)		3.3 (1.2)	3.4 (1.1)	1.18 (0.80-1.74)
Household density	234				233			
Non-crowded		192 (89.7)	22 (10.3)	1		190 (89.2)	23 (10.8)	1
Crowded		16 (80.0)	4 (20.0)	2.10 (0.59-6.50)		15 (75.0)	5 (25.0)	2.09 (0.78-5.59)
Change in social networks habits	229				228			
Same or less		115 (92.0)	10 (8.0)	1		109 (87.9)	15 (12.1)	1
More		88 (84.6)	16 (15.4)	1.82 (0.92-3.66)		91 (87.5)	13 (12.5)	0.77 (0.30-2.00)
Screen time (hours)¹	229	1.8 (2.0)	1.6 (1.1)	0.91 (0.84-0.98)*	228	1.9 (2.0)	1.7 (1.3)	0.95 (0.85-1.06)
Parent anti-SARS-CoV-2 serology	235				234			
Negative		163 (92.1)	14 (7.9)	1		159 (90.3)	17 (9.7)	1
Positive		46 (79.3)	12 (20.7)	3.44 (1.30-9.09)*		47 (81.0)	11 (19.0)	1.94 (0.64-5.81)
Adolescent anti-SARS-CoV-2 serology	235				234			
Negative		167 (90.8)	17 (9.2)	1		162 (89.0)	20 (11.0)	1
Positive		42 (82.4)	9 (17.6)	1.12 (0.55-2.27)		44 (84.6)	8 (15.4)	1.57 (0.48-5.06)

Results are odds ratio (OR) and 95% confidence intervals (CI) from multivariable generalized estimating equations adjusted for all covariates in first column.

^a based on 224 observations, ^b based on 225 observations

* indicates P < 0.05; ** indicates P < 0.01

¹Descriptive analysis presented as mean (SD); OR applicable for each additional unit of continuous variables

Table 3: Generalized estimating equations on adolescents' self-reported loneliness and sadness and emotional well-being as reported by their parents

	Self-reported sadness				Self-reported Loneliness				Parent-reported low emotional well-being (KINDL®)			
	N	No N(%)	Yes N(%)	Sadness OR (95% CI) ^a	N	No N(%)	Yes N(%)	Loneliness OR (95% CI) ^a	N	High N(%)	Low N(%)	Low emotional well-being OR (95% CI) ^b
Age of the adolescent (years)¹	236	15.4 (1.1)	14.9 (3.1)	0.88 (0.76-1.03)	237	15.3 (1.6)	14.9 (3.2)	0.95 (0.81-1.11)	234	15.4 (1.2)	15.3 (1.1)	0.78 (0.57-1.07)
Age of the parent (years)¹	236	47.8 (4.9)	49.5 (6.0)	1.07 (1.00-1.15) *	237	48.2 (5.2)	49.3 (6.0)	1.05 (0.96-1.15)	234	48.0 (5.2)	50.2 (5.9)	1.08 (0.99-1.17)
Sex of the adolescent	236				237							
Boy		94 (75.2)	31 (24.8)	1		109 (86.5)	17 (13.5)	1	234	112 (89.6)	13 (10.4)	1
Girl		58 (52.3)	53 (47.7)	3.29 (1.64-6.57) **		73 (65.8)	38 (34.2)	4.15 (2.02-8.51) **		84 (77.1)	25 (22.9)	3.03 (1.32-6.90) *
Self-perceived mood of the parent	236											
Good		138 (69.3)	61 (30.7)	1	237	158 (79.0)	42 (21.0)	1	234	168 (84.4)	31 (15.6)	1
Average to poor		14 (37.8)	23 (62.2)	2.81 (1.21-6.56) *		24 (64.9)	13 (35.1)	1.67 (0.71-3.98)		28 (80.0)	7 (20.0)	0.86 (0.29-2.55)
Financial situation of the household	236											
Good		115 (66.9)	57 (33.1)	1	237	136 (78.6)	37 (21.4)	1	234	147 (85.5)	25 (14.5)	1
Average to poor		22 (48.9)	23 (51.1)	2.30 (1.00 -5.29) *		30 (66.7)	15 (33.3)	2.37 (0.95-5.85)		32 (74.4)	11 (25.6)	2.08 (1.00-7.79) *
No answer		15 (78.9)	4 (21.1)	0.77 (0.20-2.92)		16 (84.2)	3 (15.8)	1.41 (0.36-5.50)		17 (89.5)	2 (10.5)	1.12 (0.26-4.81)
Household size (individuals)¹	236	3.3 (1.1)	3.3 (1.3)	1.16 (0.92-1.45)	237	3.3 (1.1)	3.4 (1.5)	1.15 (0.82-1.61)		3.4 (1.2)	3.1 (1.4)	0.75 (0.44-1.27)
Household density	235				236				233			
Non-crowded		142 (66.0)	73 (34.0)	1		167 (77.3)	49 (22.7)	1		182 (85.4)	31 (14.6)	1
Crowded		10 (50.0)	10 (50.0)	1.48 (0.48-4.54)		15 (75.0)	5 (25.0)	1.13 (0.84-1.52)		13 (65.0)	7 (35.0)	3.20 (1.26-8.10) *
Change in social networks habits	230				231				228			
Same or less		93(73.8)	33 (26.2)	1		105 (83.3)	21 (16.7)	1		107 (86.3)	17 (13.7)	1
More		55 (52.9)	49 (47.1)	2.05 (1.08-3.88) *		72 (68.6)	33 (31.4)	1.93 (0.96-3.90)		84 (80.8)	20 (19.2)	1.33 (0.52-3.36)
Screen time (hours)¹	230	1.7 (1.8)	2.0 (2.0)	1.04 (0.97-1.11)	231	1.8 (1.8)	2.1 (2.1)	1.03 (0.96-1.12)	228	1.9 (2.0)	1.5 (1.5)	1.04 (0.96-1.12)
Parent anti-SARS-CoV-2 serology	236				237				234			
Negative		121 (68.4)	56 (31.6)	1		139 (78.1)	39 (21.9)	1		148 (84.1)	28 (15.9)	1
Positive		31 (52.5)	28 (47.5)	2.38 (1.10-5.14) *		43 (72.9)	16 (27.1)	2.04 (0.89-5.17)		48 (82.8)	10 (17.2)	0.67 (0.22-2.02)
Adolescent anti-SARS-CoV-2 serology	236				237				234			
Negative		116 (63.4)	67 (36.6)	1		135 (73.4)	49 (26.6)	1	234	153 (84.1)	29 (15.9)	1
Positive		36 (67.9)	17 (32.1)	0.55 (0.24-1.25)		47 (88.7)	6 (11.3)	0.20 (0.07-0.63) **		43 (82.7)	9 (17.3)	1.59 (0.60-4.22)

Results are odds ratio (OR) and 95% confidence intervals (CI) from multivariable generalized estimating equations adjusted for all covariates in first column

^a based on 229 observations; ^b based on 225 observations

* indicates P < 0.05; ** indicates P < 0.01

¹ Descriptive analysis presented as mean (SD); OR applicable for each additional unit of continuous variables

Table 4: Risk factors of discrepancies between adolescents' and parents' perception of adolescents' low HRQoL

Adolescent and parent perception of adolescent HRQoL				
	N	Match N(%)	Mismatch N(%)	Perception mismatch OR (95%CI)
Age of the adolescent (years)¹	232	15.4 (1.2)	15.2 (1.1)	0.86 (0.68-1.16)
Age of the parent (years)¹	232	48.4 (5.3)	48.4 (5.8)	1.02 (0.96-1.10)
Sex of the adolescent	232			
Boy		108 (87.1)	16 (12.9)	1
Girl		76 (70.4)	32 (29.6)	2.88 (1.54-5.41)**
Self-perceived mood of the parent	232			
Good		161 (81.7)	36 (18.3)	1
Average to poor		23 (65.7)	12 (34.3)	1.91 (0.67-5.39)
Financial situation of the household	232			
Good		135 (78.9)	36 (21.1)	1
Average to poor		32 (76.2)	10 (23.8)	1.03 (0.68-2.88)
No answer		17 (89.5)	2 (10.5)	0.37 (0.07-1.90)
Household size (individuals)¹	232	3.2 (1.0)	3.6 (1.7)	1.24 (0.85-1.82)
Household density	231			
Non-Crowded		172 (81.5)	39 (18.5)	1
Crowded		11 (55.0)	9 (45.0)	2.51 (0.76-8.09)
Change in social networks habits	226			
Same or less		99 (80.5)	24 (19.5)	1
More		79 (76.7)	24 (23.3)	0.88 (0.42-1.88)
Screen time (hours)¹	226	1.9 (2.0)	1.6 (1.2)	0.93 (0.87-1.01)
Serology result of the parent	232			
Negative		145 (82.9)	30 (17.1)	
Positive		39 (68.4)	18 (31.6)	1.72 (0.67-4.37)
Serology result of the adolescent	232			
Negative		148 (81.3)	34 (18.7)	1
Positive		36 (72.0)	14 (28.0)	2.54 (0.98-6.62)
Family well-being score¹	232	66.3 (7.8)	58.8 (10.0)	0.91 (0.86-0.96)**

Results are odds ratio (OR) and 95% confidence intervals (CI) from multivariable generalized estimating equations adjusted for all covariates in first column based on 225 observations. Perception mismatch is coded as 1 if low HRQoL from adolescents' and parents' perception do not match and 0 otherwise.

* indicates $P < 0.05$; ** indicates $P < 0.01$.

¹ Descriptive analysis presented as mean (SD); OR applicable for each additional unit of continuous variables