

1 Supplementary Appendix
2
3 Low Gut Ruminococcaceae Levels are Associated with Occurrence of Antibiotic-associated
4 Diarrhea
5
6 **Table S1. Demographic Characteristics of Healthy Volunteers. Subject 18, 28, and 32 did**
7 **not meet the requirements to participate in the study during screening and were not**
8 **dosed with antibiotics. Individuals with AAD are highlighted in yellow.2**
9 **Table S2. Demographic Characteristics of Individuals Between the AAD And Non-AAD**
10 **Groups. Two-tailed Mann-Whitney tests were performed to test for significant differences**
11 **between the AAD and non-AAD groups.3**
12 **Table S3. Frequency and Bristol Stool Scale type. Faeces samples collection log book from**
13 **20 Aug 2019 to 20 Jan 2020. Subject 18, 28 and 32 failed during screening and were not**
14 **dosed with antibiotics. Individuals with AAD are highlighted in yellow. Gray shaded boxes**
15 **indicate that donors could not produce the stool samples or the stool samples were not**
16 **collected. A total of 197 faeces samples were collected.4**
17 **Table S4. Primer-probe sets used in this study6**
18 **Table S5. gBlocks Gene Fragments sequences for Faecalibacterium prausnitzii, 16S rRNA, C.**
19 **difficile TcdA and TcdB gene.....7**
20 **Figure S1. Enterobacteriaceae bloom (EB) were more common in the AAD group with a**
21 **much higher magnitude.....9**
22 **Figure S2. PCoA of baseline samples based on the ASV level bray-curtis dissimilarity.10**
23
24

25 **Table S1. Demographic Characteristics of Healthy Volunteers.** Subject 18, 28, and 32 did not meet the
 26 requirements to participate in the study during screening and were not dosed with antibiotics. Individuals with
 27 AAD are highlighted in yellow.

ID	Age range	Sex	Ethnicity	Height (m)	Weight (kg)	BMI ^a	Total doses of Amoxicillin-Clavulanate
1	36-40	Female	Chinese	1.61	68	26.4	6
2	26-30	Male	Malay	1.68	71	25.0	4 ^c
3	31-35	Female	Chinese	1.61	58	22.2	6
4	36-40	Male	Chinese	1.79	68	21.1	6
5	26-30	Male	Chinese	1.77	96	30.5	6
6	36-40	Male	Chinese	1.77	82	26.1	6
7	36-40	Female	Chinese	1.71	73	24.9	6
8	26-30	Female	Chinese	1.63	69	25.8	6
9	21-25	Female	Chinese	1.62	52	20.0	6
10	21-25	Male	Chinese	1.67	67	24.1	6
11	21-25	Female	Chinese	1.65	52	19.1	6
12	26-30	Female	Chinese	1.56	58	23.8	6
13	31-35	Female	Chinese	1.65	64	23.4	6
14	36-40	Female	Chinese	1.58	61	24.4	6
15	21-25	Female	Chinese	1.63	48	18.1	1 ^c
16	26-30	Male	Chinese	1.6	74	28.9	5 ^c
17	36-40	Female	Chinese	1.59	74	29.4	6
19	21-25	Male	Chinese	1.8	88	27.2	6
20	36-40	Female	Chinese	1.58	57	22.7	6
21	31-35	Male	Malay	1.69	68	23.9	6
22	21-25	Female	Chinese	1.60	74	28.9	6
23	31-35	Male	Chinese	1.78	108	34.1	6
24	26-30	Male	Chinese	1.71	73	24.8	3 ^c
25	21-25	Male	Chinese	1.53	50	21.7	1 ^b
26	21-25	Male	Chinese	1.75	77	25.0	6
27	21-25	Male	Chinese	1.72	69	23.3	6
29	21-25	Male	Chinese	1.77	70	22.4	6
30	36-40	Male	Chinese	1.76	78	25.3	6
31	26-30	Female	Indian	1.64	73	27.1	6
33	26-30	Female	Vietnamese	1.51	52	22.9	6

28
 29 ^bSubject 25 developed severe vomiting after a single dose of amoxicillin-clavulanate, resulting in discontinuation
 30 of antibiotics on day 1.

31 ^cSubject 2, 15, 16 and 24 experienced 3 or more episodes of watery stool in a 24-hour period (Table S2), resulting
 32 in discontinuation of antibiotics.
 33

34 **Table S2. Demographic Characteristics of Individuals Between the AAD And Non-AAD Groups.** Two-tailed
 35 Mann-Whitney tests were performed to test for significant differences between the AAD and non-AAD groups.

	AAD individuals (n=13)	non-AAD individuals (n=17)	p value
Sex	--	--	0.46
Female	5(38%)	10(59%)	
Male	8(62%)	7(41%)	
Age	30(26-37)	28(24-34)	0.34
BMI	24.86(23.91-26.40)	23.75(22.19-25.84)	0.39
Height	1.68(1.63-1.71)	1.63(1.59-1.75)	0.31
Weight	70.65(67.10-74.00)	68.65(57.8-73.95)	0.46

Data are n/N (%) for proportional data and median (IQR) for continuous data, unless specified otherwise.

36

Table S3. Frequency and Bristol Stool Scale type. Faeces samples collection log book from 20 Aug 2019 to 20 Jan 2020. Subject 18, 28 and 32 failed during screening and were not dosed with antibiotics. Individuals with AAD are highlighted in yellow. Gray shaded boxes indicate that donors could not produce the stool samples or the stool samples were not collected. A total of 197 faeces samples were collected.

ID	Screening		Day 0		Day 1		Day 2		Day 3		Day 7		Day 14 +/- 2		Day 28 +/- 2	
	BS	Frequency	BS	Frequency	BS	Frequency	BS	Frequency	BS	Frequency	BS	Frequency	BS	Frequency	BS	Frequency
1	4	2	4	2	5	3	6	2	6	2	4	2	4	1	4	2
2	3	3	3	3	3	4	5	5 ^a	7	8	NA	NA	3	4	4	3
3	4	1	4	1	4	1	4	1	4	2	4	2	4	1	4	2
4	3	1	3	1	3	1	3	1	4	1	3	1	4	1	NA	0
5	5	1	5	3	7	2	5	1	6	2	4	1	4	1	4	3
6	4	2	5	2	6	2	6	1	6	1	4	1	4	1	1	1
7	4	1	4	1	6	2	6	2	5	1	4	1	4	2	2	1
8	4	2	4	1	5	1	5	1	5	1	4	1	3	2	4	2
9	4	1	4	1	4	1	5	1	5	2	4	1	3	1	4	2
10	4	2	4	1	4	1	4	1	7	4	NA	NA	NA	NA	4	4
11	3	1	4	1	4	1	4	2	4	1	2	1	3	1	4	1
12	4	1	4	1	NA	NA	NA	NA	4	1	4	1	4	1	4	1
13	5	1	6	1	6	1	5	1	4	2	6	2	4	1	6	2
14	3	1	4	1	4	1	4	2	4	1	4	2	4	2	3	1
15	3	1	3	1	7	3 ^b	6	3	4	1	5	1	4	2	3	1
16	4	1	6	1	6	1	6	2	7	5 ^c	6	1	6	2	6	1
17	4	1	4	2	4	1	4	1	4	1	4	1	4	1	4	1
19	6	1	3	2	7	2	2	1	3	1	2	1	4	1	3	1
20	3	1	4	1	6	1	6	1	3	1	3	1	3	1	4	2
21	2	1	4	1	4	2	6	1	3	1	4	1	3	2	6	1
22	5	1	6	1	4	1	4	1	4	1	4	1	4	1	4	2
23	4	1	3	2	4	1	4	1	4	1	4	2	4	1	4	2
24	5	2	5	1	7	3 ^d	4	2	1	2	4	2	4	1	4	2

25	3	1	3	1	4	1	5	1	4	1	5	1	4	1	4	1
26	4	1	3	2	3	1	4	2	4	2	4	2	4	2	4	2
27	4	1	4	2	5	1	5	3	5	1	4	1	4	2	4	1
29	4	1	4	2	4	2	4	3	4	2	5	2	4	2	4	3
30	4	1	4	2	2	1	2	2	2	1	5	4	4	1	4	1
31	4	3	4	4	4	2	4	2	4	2	4	2	5	1	4	2
33	2	1	1	2	2	1	3	1	2	1	1	1	NA	NA	NA	NA

^aSubject 2 experienced 5 episodes of watery stool in a 24-hour period, resulting in discontinuation of antibiotics on day 2, with the total doses of amoxicillin-clavulanate as 4.

^bSubject 15 experienced 3 episodes of watery stool in a 24-hour period, resulting in discontinuation of antibiotics on day 1, with the total doses of amoxicillin-clavulanate as 1.

^cSubject 16 experienced 5 episodes of watery stool in a 24-hour period, resulting in discontinuation of antibiotics on day 3, with the total doses of amoxicillin-clavulanate as 5.

^dSubject 24 experienced 3 episodes of watery stool in a 24-hour period, resulting in discontinuation of antibiotics on day 1, with the total doses of amoxicillin-clavulanate as 3.

Table S4. Primer-probe sets used in this study

Name	Sequences	References
16S_V4_515F primer	GTGCCAGCMGCCGCGGTAA	(1)
16S_V4_806R primer	GGACTACHVGGGTWTCTAAT	
F. prau_FP	GATGGCCTCGCGTCCGATTAG	(2)
F. prau_RP	CCGAAGACCTTCTCCTCC	
16S_FP	ACTCCTACGGGAGGCAGCAG	(3)
16S_RP	ATTACCGCGGCTGCTGG	
TcdA_FP	CAGTCGGATTGCAAGTAATTGACAAT	(4)
TcdA_RP	AGTAGTATCTACTACCATTAACAGTCTGC	
TcdA_Probe	/56-FAM/TTGAGATGATAGCAGTGTCAGGATTG/36-TAMSp/	
TcdB_FP	TACAAACAGGTGTATTTAGTACAGAAGATGGA	
TcdB_RP	CACCTATTTGATTTAGMCCTTTAAAAGC	
TcdB_Probe	/56-FAM/TTTKCCAGTAAAATCAATTGCTTC/36-TAMSp/	

Table S5. gBlocks Gene Fragments sequences for *Faecalibacterium prausnitzii*, 16S rRNA, *C. difficile* TcdA and TcdB gene.

gBlocks Gene Fragments	Sequences
gBlocks Gene Fragments for <i>Faecalibacterium prausnitzii</i> and 16S rRNA positive control	GGTAGAGGGAAAAGGAGCAATCCGCTTTGAGATGGCCTCGCGTCCGATTAGCTAGTTGGTGA GGTAATGGCCCACCAAGGCGACGATCGGTAGCCGGACTGAGAGGTTGAACGGCCACATTGG GACTGAGACACGGCCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAATGGGGG AAACCCTGATGCAGCGACGCCGCTGGAGGAAGAAGGTCTTCGGATTGTAAACTCCTGTTGT TGAGGAAGATAATGACGGTACTCAACAAGGAAGTGACGGCTAACTACGTGCCAGCAGCCGC GGTAATACGTAGGTCACAAGCGTTGTCCGGAATTAC
gBlocks Gene Fragments for <i>C. difficile</i> TcdA positive control	GGTAGAGGGAAAAGGAGCAATCCGCTTTGAGATGGCCTCGCGTCCGATTAGCTAGTTGGTGA GGTAATGGCCCACCAAGGCGACGATCGGTAGCCGGACTGAGAGGTTGAACGGCCACATTGG GACTGAGACACGGCCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAATGGGGG AAACCCTGATGCAGCGACGCCGCTGGAGGAAGAAGGTCTTCGGATTGTAAACTCCTGTTGT TGAGGAAGATAATGACGGTACTCAACAAGGAAGTGACGGCTAACTACGTGCCAGCAGCCGC GGTAATACGTAGGTCACAAGCGTTGTCCGGAATTAC
gBlocks Gene Fragments for <i>C. difficile</i> TcdB positive control	TACACCCCTGCGGCAACGTTGAAGCTCCTGGATTACACTGGCTGGATCTAAGCCGTGACACC CGTCATACTCCATAACCGTCTGTAACCTCACGGCTTGTCTGGACTGGATTGCCATTCTCTCAG AGTATTATGCAGGCCGGCGTACGGGTCCCATATAAACCTGTCATAGCTTACCTGACTCTACTT GGAAATGTGGCTTACAAACAGGTGTATTTAGTACAGAAGATGGATTTAAATATTTTGCCCA GCTAATAACACTTGATGAAAACCTAGAAGGAGAAGCAATTGATTTTACTGGAAAATTAATTAT TGACGAAAATATTTATTTTATTTGATGATAATTATAGAGGAGCTGTAGAATGGAAAAGAATTAG ATGGTGAAAATGCACTATTTTAGCCAGAAAACAGGTAAAGCTTTTAAAGTCTAAATCAAATA GGTG







Figure S1. Enterobacteriaceae bloom (EB) were more common in the AAD group with a much higher magnitude.

Related to Figure 2. (A) A panel list of the 10 most abundant families in individuals in the AAD (top panel) and non-AAD groups (bottom panel). (B) Enterobacteriaceae blooms (EB) were observed more frequently in the AAD group compared to the non-AAD group (76.9% vs 29.4%). As the increase in Enterobacteriaceae varies between days 1-3 across different individuals, we defined Enterobacteriaceae blooms as having the maximum of Enterobacteriaceae abundance across days 1-3 in each individual being larger than the median(max(days 1-3)-baseline). (C) For all the individuals who experienced EB, the relative abundance of Enterobacteriaceae is higher in the individuals with AAD compared to the individuals with non-AAD (mean 0.59 vs 0.21, FDR corrected $p < 0.0001$). (D) Pie charts of the most abundant genera in Enterobacteriaceae across days 1-3. Within the Enterobacteriaceae family, we observed a higher abundance of Escherichia-Shigella (49.9%) at the genus level.




Figure S2. PCoA of baseline samples based on the ASV level bray-curtis dissimilarity.

Related to Figure 3. Display is based on sample scores on the primary axis (PCoA1, 17.0% variance explained) and secondary axis (PCoA2, 15.5% variance explained). Vector loading plots at the ASV level indicate that ASV assigned to *Faecalibacterium* points to the non-AAD group while many unified features (e.g., ASVs assigned to *Blautia*, *Bifidobacterium*, *Collinsella*, *Bacteroides_vulgatus*) point to the AAD groups (multiple linear correlation with the threshold setting as 0.3).

Reference

1. Caporaso JG et al. Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *Proc. Natl. Acad. Sci. U. S. A.* 2011;108(SUPPL. 1). doi:10.1073/pnas.1000080107
2. Carroll IM, Ringel-Kulka T, Siddle JP, Ringel Y. Alterations in composition and diversity of the intestinal microbiota in patients with diarrhea-predominant irritable bowel syndrome. *Neurogastroenterol. Motil.* 2012;24(6). doi:10.1111/j.1365-2982.2012.01891.x
3. Van Hul M et al. From correlation to causality: the case of Subdoligranulum. *Gut Microbes* 2020;12(1):1–13.
4. Kubota H et al. Development of TaqMan-based quantitative PCR for sensitive and selective detection of toxigenic clostridium difficile in human stools. *PLoS One* 2014;9(10). doi:10.1371/journal.pone.0111684