

Supplementary Material:


Endpoint	African	African p	Hispanic	Hispanic p	Asian	Asian p	European
Prostate Cancer	2.43 [2.38-2.48]	<10 ⁻¹⁶	1.04 [1.0-1.08]	0.05	0.86 [0.76-0.95]	<10 ⁻²	Ref
Metastatic	3.01 [2.83-3.18]	<10 ⁻¹⁶	1.27 [1.13-1.41]	<10 ⁻⁴	1.02 [0.7-1.37]	0.88	Ref
Fatal	2.65 [2.37-2.96]	<10 ⁻¹⁶	1.15 [0.9-1.42]	0.24	0.58 [0.17-1.08]	0.15	Ref

Supplementary Table 1: Association of ancestry with three prostate cancer clinical endpoints. Hazard ratios (HRs) are shown from Cox-proportional hazards models with ancestry group as the only predictor variable. Numbers in brackets are 95% confidence intervals.


Endpoint	p Value	FH
Prostate Cancer	$<10^{-16}$	2.06 [2.0-2.12]
Metastatic	$<10^{-16}$	1.67 [1.5-1.83]
Fatal	$<10^{-11}$	1.84 [1.54-2.17]

Supplementary Table 2: Association of family history with three prostate cancer clinical endpoints. Hazard ratios (HRs) are shown from Cox-proportional hazards models with family history as the only predictor variable. This analysis was limited to the 374,455 participants who provided family history information in baseline survey data. Numbers in brackets are 95% confidence intervals.

A.



B.



Supplementary Figure 1. (A) PHS290 score density plot in prostate cancer controls for European, African, Hispanic and Asian ancestry groups. **(B)** PHS290 score density plot in prostate cancer cases for European, African, Hispanic and Asian ancestry groups. These plots do not account for age.