

1 Supplementary Table 1: Patient characteristics.

| Variables | Value | n (%) |
|--------------------------------|------------------|------------|
| Number of patients | | 491 |
| Age at time of surgery (years) | Median [min-max] | 69 [61-76] |
| | ≤65 | 188 (38.3) |
| | >65 | 303 (61.7) |
| Sex | Male | 329 (67.0) |
| | Female | 162 (33.0) |
| Histologic type | Differentiated | 270 (55.0) |
| | Undifferentiated | 221 (45.0) |
| Stage (TNM7 ^b) | IB | 116 (23.6) |
| | IIA | 86 (17.5) |
| | IIB | 116 (23.6) |
| | IIIA | 65 (13.2) |
| | IIIB | 64 (13.0) |
| | IIIC | 44 (9.0) |
| <i>Helicobacter pylori</i> | + | 175 (35.6) |
| | - | 316 (64.4) |
| Modality | Surgery + S-1 | 176 (35.8) |
| | Surgery only | 315 (64.1) |
| S-1 reduction | + | 82 (46.6) |
| | - | 94 (53.4) |

2


TMN7^b, TMN Classification of Malignant Tumours-7th edition

3

1 Supplementary Table 2: Concordance rate between IHC of MMR and MSI analysis status


| Immunohistochemistry of MMR | Microsatellite status | | |
|-----------------------------|-----------------------|----------------------|-----------|
| | MSI ⁺ (%) | MSI ⁻ (%) | Total (%) |
| dMMR | 1 (1.2) | 6 (7.1) | 7 (8.3) |
| pMMR | 8 (9.5) | 69 (82.1) | 77 (91.7) |
| Total | 9 (10.8) | 75 (89.2) | 84 |

2



1 Supplementary Figure 1: MSI analysis. Tumors showing allelic shifts in \geq one of five
 2 markers (BAT-26, NR-21, BAT-25, MONO-27, and NR-24) were defined as MSI⁺
 3 (hereafter referred to as MSI) and the rest were defined as MSI⁻. MSI is shown in black
 4 arrows. N; normal tissue, T; tumor tissue. BAT-26 is represented as a blue line. NR-21,
 5 BAT-25, and MONO-27 are represented as a green line. NR-24 is represented as a black
 6 line.

7



1 Supplementary Figure 2: IHC of PD-L1,CD4/CD8, and MMR proteins staining pattern
2 (A) Representative IHC staining of PD-L1, CD4, and CD8. Magnification, $\times 200$. (B)
3 Representative IHC staining of MMR proteins, including MLH1, MSH2, MSH6,
4 and PMS2, $\times 200$.
5

1 **Supplementary Methods**

2 FFPE tissue section slides were deparaffinized in 2 changes of xylene for 15 minutes each,
3 followed by rehydration in an ethanol gradient (100% ethanol, 95% ethanol, 70% ethanol,
4 and water). Antigen retrieval was performed in a 95°C water bath with Envision FLEX
5 low pH antigen retrieval solution (1x) (20 minutes pre-heat, 20 minutes heat, and 20
6 minutes cool).

7 8 **IHC of PD-L1**

9 Deparaffinization, rehydration, and antigen retrieval were performed on PT Link (PT100,
10 Dako, Santa Clara, CA, USA) using the EnVision™ FLEX Target Retrieval Solution, at
11 low pH 6.0 for 53 minutes at room temperature. Following FLEX peroxidase block for 5
12 minutes, specimens were incubated with primary mouse anti-PD-L1 monoclonal
13 antibody (M365329; Dako, Santa Clara, CA, USA), using a concentration of 1:50 for 60
14 minutes at room temperature. Specimens were then incubated with the EnVision™
15 FLEX+ Mouse LINKER for 30 minutes at room temperature, followed by incubation
16 with the EnVision™ FLEX hprseradish peroxidase (HRP) visualization reagent for 30
17 minutes at room temperature. The enzymatic conversion of the subsequently added 3,3'-
18 diaminobenzidine tetrahydrochloride (DAB) chromogen for 10 minutes at room
19 temperature followed by DAB enhancer for 5 minutes at room temperature resulted in
20 precipitation of a visible reaction product at the site of antigen. The specimens were then
21 counterstained with haematoxylin and coverslipped. All steps were followed by buffer
22 washes for 5 minutes each (EnVision™ FLEX Wash Buffer 20x).

23 24 **IHC of CD4/CD8**

25 An anti-CD4/CD8 primary antibody cocktail was added for 30 minutes. Slides were
26 rinsed twice in buffer and the dual anti-mouse/anti-rabbit secondary detection system
27 (MACH2 polymer) was added for 30 minutes. The slides were then rinsed twice in buffer
28 prior to detection with Betazoid DAB chromagen for 5 minutes (CD4+ cells), followed
29 by a water rinse, a buffer rinse, and Warp Red chromagen for 7 minutes (CD8+ cells).
30 Slides were counterstained in Mayer's Hematoxylin (Sigma-Aldrich, St. Louis, NI, USA)
31 and Scott's Tap Water Substitute, with a final rinse in water, and dried at 60°C for 15
32 minutes, followed by rinsing in xylene. Antigen retrieval was performed in a 95°C water
33 bath with Envision FLEX low pH antigen retrieval solution (1x) (20 minutes pre-heat, 20
34 minutes heat, and 20 minutes cool). The tissue sections were stained on a machine
35 (Autostainer Link 48, Dako, Santa Clara, CA, USA) following manufacturer's
36 instructions, with the following modification: DAB chromogen was applied once to

1 the tissue sections for 10 minutes. Slides were then counterstained in Mayer's
2 Hematoxylin and Scott's Tap Water Substitute, with a final rinse in water and allowed to
3 air dry.

4 5 **IHC of MMR**

6 MMR IHC was conducted on 3 µm-thick representative paraffin sections. The sections
7 were deparaffinized in xylene and dehydrated. Immunostaining was performed for each
8 protein. Endogenous peroxidase activity was inhibited by incubating the sections in 3%
9 H₂O₂ for 20 minutes. Antigen retrieval was required for all antibodies and consisted of
10 microwave treatment of sections for 20 minutes in citrate buffer (pH 7.0) in a microwave
11 oven at 300 W (H2500, Microwave Processor, Bio Rad, Hercules, CA, USA) as
12 previously described (54). The DAKO EnVision+ system [dextran polymers conjugated
13 with HRP; (DAKO, Santa Clara, CA, USA)] was used to examine immunohistochemical
14 expression of these markers. Negative control staining was conducted with a routine
15 method using the DAKO EnVision+ system (Dako, Santa Clara, CA, USA) according to
16 the manufacturer's instructions.