

# 1 **Taste loss as a distinct symptom of COVID-19: A systematic review** 2 **and meta-analysis**

3  
4 Mackenzie E. Hannum<sup>1</sup>, Riley J. Koch<sup>1</sup>, Vicente A. Ramirez<sup>1,2</sup>, Sarah S. Marks<sup>1</sup>, Aurora K.  
5 Toskala<sup>1</sup>, Riley D. Herriman<sup>1</sup>, Cailu Lin<sup>1</sup>, Paule V. Joseph<sup>3,4</sup> and Danielle R. Reed<sup>1\*</sup>

6  
7 <sup>1</sup>Monell Chemical Senses Center, 3500 Market St, Philadelphia PA 19104

8 <sup>2</sup>Department of Public Health, University of California Merced, Merced, CA 95348

9 <sup>3</sup>Division of Intramural Research, National Institute of Nursing Research, National Institutes of  
10 Health, Bethesda, MD, USA

11 <sup>4</sup>Division of Intramural Research, National Institute of Alcohol Abuse and Alcoholism, National  
12 Institutes of Health, Bethesda, MD, USA

## 13 **Abstract**

14 Chemosensory scientists have been skeptical that reports of COVID-19 taste loss are genuine, in  
15 part because before COVID-19, taste loss was rare and often confused with smell loss.

16 Therefore, to establish the predicted prevalence rate of taste loss in COVID-19 patients, we  
17 conducted a systematic review and meta-analysis of 376 papers published in 2020–2021, with  
18 241 meeting all inclusion criteria. Additionally, we explored how methodological differences  
19 (direct vs. self-report measures) may affect these estimates. We hypothesized that direct  
20 prevalence measures of taste loss would be the most valid because they avoid the taste/smell  
21 confusion of self-report. The meta-analysis showed that, among 138,897 COVID-19-positive  
22 patients, 39.2% reported taste dysfunction (95% CI: 35.34–43.12%), and the prevalence  
23 estimates were slightly but not significantly higher from studies using direct (n = 18) versus self-  
24 report (n = 223) methodologies (Q = 0.57, df = 1, p = 0.45). Generally, males reported lower  
25 rates of taste loss than did females and taste loss was highest in middle-aged groups. Thus, taste  
26 loss is a bona fide symptom COVID-19, meriting further research into the most appropriate  
27 direct methods to measure it and its underlying mechanisms.

28 **Key words:** gustatory dysfunction, taste loss, ageusia, COVID-19, coronavirus

29

## 30 **Introduction**

31           The novel coronavirus (COVID-19), a respiratory infection caused by severe acute  
32 respiratory syndrome coronavirus-2 (SARS-CoV-2), was first identified in Wuhan, China, and  
33 has since spread throughout the world. When the World Health Organization first declared this a  
34 pandemic in March 2020, researchers and clinicians were not yet aware that the virus affected  
35 individuals' senses of smell and taste, but these symptoms soon became apparent via patient  
36 reports. As a result of COVID-19, affected people can experience chemosensory dysfunction in a  
37 variety of ways, including complete loss of smell or taste (anosmia or ageusia, respectively),  
38 partial loss of smell or taste (hyposmia or hypogeusia), and/or a distorted sense of smell or taste  
39 (e.g., parosmia, dysgeusia). These chemosensory dysfunctions can be distressing to the affected  
40 individuals and can last for extended times, with some patients experiencing resolution within a  
41 few weeks to a month (Lee et al., 2020b; Gerkin et al., 2021) and others with symptoms for 6  
42 months or longer (Blomberg et al., 2021).

43           Previous meta-analyses examined smell and taste loss in COVID-19 patients, but often  
44 with a focus on onset and duration (Santos et al., 2021) or recovery (Boscutti et al., 2021) of  
45 chemosensory symptoms. Many focused only on smell loss (Hannum et al., 2020; Pang et al.,  
46 2020; Rocke et al., 2020) or general neurological symptoms (Abdullahi et al., 2020; Favas et al.,  
47 2020; Mair et al., 2021; Yassin et al., 2021). Very few continued to evaluate articles published in  
48 2021 and often capped reviewing articles 6-10 months after March 2020, when the pandemic  
49 was declared, limiting the number of articles included (ranging from 5 to 59 articles total).  
50 Therefore, we decided to conduct a more comprehensive analysis, spanning a year and a half, to  
51 ensure a fuller coverage of the available research.

52            Additionally, taste loss is often neglected in research compared to smell loss, as there is a  
53 common notion that taste loss is not as “real” as smell loss. Some claim taste loss is  
54 indistinguishable from smell loss (Le Bon et al., 2021) or is confused with smell loss (Deems et  
55 al., 1991), specifically with retronasal smell perception (Hintschich et al., 2020a). For the general  
56 population, loss of taste can be difficult to distinguish from smell loss. Therefore, it may be  
57 difficult to know, based on self-report measures alone, whether or not a participant truly lost  
58 their sense of taste.

59            Thus, many chemosensory researchers may attribute the taste loss phenomena seen in the  
60 current reports of COVID-19-positive patients to deficiencies of self-report or subjective  
61 measures of taste loss. Therefore, we conducted a systematic review and meta-analysis to  
62 estimate the true prevalence of taste loss in COVID-19 patients across a wide sample of studies  
63 (n=241) and to evaluate effects of major methodological differences in data collection. In  
64 particular, we compared overall findings on taste loss as determined by individual taste tests  
65 (herein referred to as direct tests) with self-reports without direct sensory testing. We  
66 hypothesized that direct methodologies would support the presence of taste loss as a distinct  
67 symptom, and direct measures might even be higher than self-report despite the possible inflation  
68 of self-reported taste loss which is exacerbated by smell loss.

69            Currently scientists are using both direct and self-report measures to examine  
70 chemosensory dysfunction, with self-report far more common due to the pandemic restrictions,  
71 e.g., sensory laboratories where direct testing is often conducted are closed. For taste, direct tests  
72 include standardized and non-standardized tests that contain various sweet, salty, and sometimes  
73 bitter and sour stimuli given to participants via solutions, drops, strips, or sprays (Cao et al.,  
74 2021; Singer-Cornelius et al., 2021). Non-standardized direct taste measures created to study

75 COVID-19-related taste dysfunction include solution-based tests, often prepared at home by  
76 participants (Vaira et al., 2020f). Self-report measures include interviews with researchers and  
77 clinicians, electronic health records as well as surveys administered over the phone, online, or in  
78 person.

79 To understand taste loss as a symptom of COVID-19, we conducted a large systematic  
80 review and meta-analysis, examining how it has been measured (direct vs. self-report) and how  
81 the measurement type can affect prevalence rates. We tested the hypothesis that direct measures  
82 are at least as sensitive as self-report measures and would confirm taste loss as a distinct  
83 symptom and not merely misattributed smell loss.

## 84 **Methods**

### 85 **Article Selection**

86 This systematic review and meta-analysis followed the Preferred Reporting Items for  
87 Systematic Reviews and Meta-Analyses guidelines (Moher et al., 2009). Articles were selected  
88 via searches on Pubmed/Medline and Google Scholar, using the keyword “COVID-19” with  
89 “taste”, “smell”, and/or “olfaction”, as well as “gustatory”. Literature retrieval began on May 15,  
90 2020, and concluded on June 1, 2021, resulting in 377 articles in total.

91 Initial screening of the articles included reading the titles and abstracts to assess their  
92 relevance. Articles with an abstract that reported chemosensory dysfunction in COVID-19-  
93 infected individuals were included in the systematic review ( $n = 376$ ). Next, at least two authors  
94 read the articles initially deemed relevant, to evaluate whether they fit the inclusion criteria:  
95 reporting positive COVID-19 tests, written in the English language, and lack of population bias.  
96 COVID-19 must have been confirmed via nasopharyngeal swab, reverse transcription  
97 polymerase chain reaction (RT-PCR), or assessment by physician or other medical personnel.

98 The articles were then evaluated on whether they reported taste loss data specifically. In total,  
99 135 articles were excluded based on such criteria as not evaluating taste loss, recruiting  
100 participants with chemosensory dysfunction, not testing for COVID-19, and presenting  
101 overlapping data (see Figure 1). In total, 241 articles were included in the final meta-analysis  
102 (corresponding citations described in “Included Articles” section at the end of the paper).

### 103 **Data Extraction**

104 We extracted from each article either the number or percentage of patients with taste  
105 dysfunction due to a SARS-CoV-2 infection. The prevalence of taste loss reported in each article  
106 was calculated by dividing the reported number of participants with taste loss as a symptom by  
107 the total number of COVID-19-positive participants. Additionally, measures of taste loss were  
108 labeled as “self-report” or “direct” to identify the method used to evaluate participants. Self-  
109 report measures included reported loss of taste via surveys, interviews, and electronic medical  
110 health records. Most articles (n = 223) used self-report methods. Table 1 summarizes the studies  
111 that used direct measures (n = 18), comprising actual taste tests administered either at home  
112 (Adamczyk et al., 2020; Hintschich et al., 2020a; Petrocelli et al., 2020b; Cao et al., 2021;  
113 Singer-Cornelius et al., 2021), at a testing facility (Altin et al., 2020; Bidkar et al., 2020a;  
114 Mazzatenta et al., 2020; Ramteke et al., 2020; Vaira et al., 2020a; Le Bon et al., 2021; Niklassen  
115 et al., 2021a; Salcan et al., 2021) or both in home and in a hospital environment (Vaira et al.,  
116 2020b, 2020c, 2020d, 2020f). Vaira et al. [2020e] had an unknown testing location. Many of the  
117 measures consisted of solution-based tests measuring four basic taste sensations: sweet, sour,  
118 salty, and bitter.

119           Although some authors reported exclusively smell loss or taste loss, many reported on  
120 both senses. Thus, it was necessary to include additional coding options for when authors  
121 reported the symptoms in tandem (e.g., “loss of smell or taste”). Therefore, articles were labeled  
122 as “taste only”, “smell and/or taste”, “smell and taste”, “smell or taste”, and “smell and/or taste;  
123 taste only” (in the event both values were reported, the numbers were summed), depending on  
124 how these symptoms were phrased in the article.

125           We also extracted demographic characteristics of each study, including the population  
126 mean and/or median age, sex (expressed as percentage of males in the population), and country  
127 of origin (for the geographic distribution of the study populations, see Supplementary Figure 1  
128 (S1)).

129           Four authors performed the initial reading of full texts and the data extraction from the  
130 studies (R.D.H., A.K.T., S.S.M., R.J.K.). Two authors confirmed this information and resolved  
131 any inconsistencies (M.E.H., D.R.R.). Differences were resolved through a discussion and  
132 renewed consensus on the proposed solution from all authors who read that specific article.

### 133 **Risk-of-Bias Assessment**

134           We used a risk-of-bias assessment from Hoy et al. (2012) to examine the articles selected  
135 for the meta-analysis. The assessment contained nine questions, outlined in Supplementary  
136 Materials (see Supplementary Table (S2)). Responses were scored as 1 (no) or 0 (yes), with  
137 summary scores of low (0-3), moderate (4-6), and high (7-9). Two authors completed the risk-of-  
138 bias assessment of each article using the checklist developed by Hoy et al. (2012), as described  
139 and adapted by Tong et al. (2020) (S.S.M. and A.K.T.). One author resolved any discrepancies  
140 (M.E.H.).

### 141 **Statistical Analysis**

142 The meta-analysis was conducted using the meta package in R (Schwarzer et al., 2019).  
143 Generalized linear mixed models were used for the meta-analysis, as recommended for the  
144 analysis of binary outcomes and proportions (Bakbergenuly and Kulinskaya, 2018; Schwarzer et  
145 al., 2019). Heterogeneity (e.g., between study variance) was assessed using Cochran's  $Q$ ,  $I^2$ , and  
146 tau squared ( $\tau^2$ ). We concluded there was evidence for heterogeneity when the p-value for  
147 Cochran's  $Q$  was less than 0.05 and if the  $I^2$  was greater than 50% (Higgins and Thompson,  
148 2002). Tau squared ( $\tau^2$ ), a measure of between study variance, was estimated using the  
149 maximum likelihood approach and has no associated test-statistic.

150 An overall pooled prevalence estimates were computed and reported for a random-effects  
151 model with the parameters described for all 241 studies. While both fixed and random-effect  
152 models were computed, excess the heterogeneity among our studies (see the Results section)  
153 suggested that unmeasured effects contribute to variance in our data more than would be  
154 expected from sampling error. The extreme diversity of taste loss as reported in individual  
155 studies (0 to 93.4%) suggested that the random-effects model was more appropriate.

156 Subgroup analysis was performed for studied employing direct methods ( $n = 18$ ) and  
157 self-report methods ( $n = 223$ ) to assess taste function in COVID-19-positive individuals.  
158 Additionally, the average age of participants and their sex (percentage of male subjects in each  
159 study) were included as covariates in univariate mixed-regression models. Subgroup analysis for  
160 age was arbitrarily categorized into five groupings: adolescents (0-18 years old), young adults  
161 (19-35 years old), middle-aged adults (36-50 years old), older adults (51-65 years old), and  
162 elderly adults (65+ years old). Finally, studies that used direct tests were separated, and subgroup  
163 analysis was performed for each type of collection methodology: solution based ( $N=12$ ), strip  
164 based ( $N=5$ ), and other ( $N=1$ ). For continuous variables (e.g., age), the transformed beta



165 coefficients are reported. The transformation for the generalized linear mixed models uses a logit  
166 transformation for each proportion, so the models are interpreted as the  $\log(\text{odds})$  or  $\log(p/1 - p)$ ,  
167 where  $p$  is equal to the prevalence for each study. Back-transformation into the prevalence  
168 estimates were done for subgroup analysis (e.g., age).

169 All statistical analyses were performed using R 4.0.5 (R Core, 2021) and RStudio  
170 1.4.1106 (RStudio Team, 2020). Visualization of the meta-analysis is displayed as an orchard  
171 plot adapted from Nakagawa et al. (2020). The R scripts and compiled data used for this analysis  
172 are available without restriction at GitHub (<https://github.com/vramirez4/COVID19-TasteLoss>).

## 173 **Results**

### 174 **Risk-of-Bias Assessment**

175 Each of the 241 articles included in this meta-analysis were reviewed for risk of bias.  
176 Among these articles, none had a high risk of bias, 142 studies had a moderate risk, and 99  
177 studies had a low risk (see Supplementary Table S2 for the full assessment).

### 178 **Prevalence of taste loss in COVID-19-positive patients**

179 Among the 241 studies, the sample sizes ranged from 11 to over 40,000 patients with  
180 COVID-19. The number of cases of taste loss per study ranged from 0 to 4,668, with raw  
181 prevalence estimates ranging from 0% to 93.4%. Collectively, the meta-analysis included  
182 138,785 patients who tested positive for COVID-19. Of these, 32,918 patients had some form of  
183 taste loss after infection with SARS-CoV-2. Heterogeneity among the prevalence estimates  
184 across all studies ( $n = 241$ ) yielded a significant Cochran's  $Q$  ( $Q = 29388.75$ , degrees of freedom  
185  $[df] = 240$ ,  $P < 0.001$ ), and an  $I^2$  estimate of 99.2%, and  $\tau^2$  estimate of 1.5. The pooled estimate  
186 for taste loss prevalence in COVID-19-positive patients following meta-analysis for the overall  
187 cohort was 36.90% (95% CI: 33.27–40.69%).

## 188 **Effect of methodology (direct vs. self-report) on prevalence estimate**

189 We employed a subgroup analysis to determine the effect of direct versus self-report  
190 approaches on taste loss prevalence (see Figure 2). Eighteen studies used direct methods to  
191 assess taste loss, comprising 2,240 COVID-19 patients, with 1,092 reported cases of taste loss.  
192 Per study, the prevalence of taste loss ranged from 0% to 84% among COVID-19-positive  
193 patients. For studies using direct approaches, the pooled estimate of the prevalence for the  
194 random-effect model was 42.0% (95% CI: 30.0-55.0%). Cochran's Q was significant ( $Q =$   
195  $187.93$ ,  $df = 17$ ,  $P < 0.001$ ), and an  $I^2$  of 91.0% was obtained, confirming heterogeneity of the  
196 data collected via direct measures. The  $\tau^2$  for the direct methodologies was 1.2.

197 A total of 223 studies used self-report methods (e.g., questionnaire, interview),  
198 comprising 136,545 subjects, with 31,826 cases of taste loss. The reported prevalence of taste  
199 loss ranged from 1% to 93% per study. The pooled estimate of the prevalence under the random-  
200 effect models was 36.53% (95% CI: 32.8%-40.5%). Similar to the direct subgroup, Cochran's Q  
201 was significant ( $Q = 29188.14$ ,  $df = 222$ ,  $P < 0.001$ ), and the  $I^2$  value was 99.2%, confirming that  
202 the studies differed in prevalence across the self-report studies. The  $\tau^2$  for self-reported studies  
203 was 1.5.

204 A test of heterogeneity between groups was employed using Cochran's Q, which  
205 revealed that the differences between methodologies were not statistically significant ( $Q = 0.66$ ,  
206  $df = 1$ ,  $p = 0.4157$ ) under the random-effects model. While our analysis showed that the  
207 prevalence of taste loss was higher when measured directly than by self-report, there was no  
208 significant effect of measurement method on the prevalence estimates of taste loss.

## 209 **Effect of age and sex on taste loss prevalence**

210 Additional analyses were undertaken to assess the effect of age and sex. Univariate mixed  
211 models for each covariate revealed that both age and sex had significant effects on the  
212 prevalence of taste loss. After categorizing each study by mean age group, we conducted the  
213 meta-analysis for those 210 studies that reported the ages of the participants (see Table 2).  
214 Among this subset of studied the pooled prevalence was 37.16% and demonstrated significant  
215 heterogeneity with ( $Q=33943.46$ ,  $df = 209$ ,  $p<0.001$ ),  $I^2 = 98.9$ , and  $\tau^2 = 1.5$ . The prevalence  
216 estimates per age category ranged from 11% in studies with average ages younger than 18 years  
217 to 44% in studies with average ages between 36 and 50 years. Heterogeneity in these studies was  
218 high ( $I^2 = 96.4-98.9\%$ ) both within groups ( $Q = 14106.77$ ,  $df = 205$ ,  $p<0.0001$ ) and between  
219 groups ( $Q = 32.19$ ,  $df = 4$ ,  $p<0.0001$ ). The results demonstrated that both the youngest and oldest  
220 age groups report the lowest prevalence of taste loss, while age groups between 18 and 65 years  
221 had pooled estimates ranging from 32% to 44%, with the highest in the middle-age (36-50 years)  
222 group. Similarly, it appeared that age accounted for some of the overall heterogeneity among  
223 some groups as measured by reductions in  $\tau^2$  among studies who contained mostly adolescent  
224 and elderly individuals.

225 The effects of sex were similarly examined. A univariate mixed model using percentage  
226 of males in each study as a covariate found an effect of  $\beta = -0.0296$  ( $p < 0.001$ ) for each percent  
227 increase. Overall, the higher the percentage of males in a study, the lower the prevalence of taste  
228 loss.

### 229 **Effect of type of direct approach on taste loss prevalence**

230 When we compared the types of direct report test (e.g., solution-based test, taste strip-  
231 based test), we found a significant difference in prevalence rates (see Table 3). We classified  
232 studies into three categories: taste strip testing ( $n=5$ ), taste solution testing ( $n = 12$ ), and “other”

233 (n = 1) for methods that do not employ either solution or strips. Pooled prevalence for solution-  
234 based tests was 54.25% (95% CI: 45.01-63.19%) and for taste strips was 24.63% (95% CI:  
235 13.10-41.46%). There was reduced heterogeneity in the subgroups compared to the overall meta-  
236 analysis: solution-based testing,  $I^2 = 88.8\%$ ; and strip-based testing,  $I^2 = 76.8\%$ . Measurements  
237 of  $\tau^2$ , were 0.3807 and 0.5991 for solution-based testing and strip-based testing respectively.  
238 There was significant heterogeneity between our pooled estimates ( $Q = 8.68$ ,  $df = 2$ ,  $p = 0.01$ ).  
239 Together, our results demonstrate that studies using solution-based taste tests, on average, result  
240 in higher prevalence of taste loss in COVID-19 patients than do studies using strips or other  
241 methods.

## 242 **Discussion**

243 Despite the occurrence of true taste loss in a variety of diseases such as cancer (Nolden et  
244 al., 2019), as well as in the general population (Rawal et al., 2016), taste loss has often been  
245 confused with smell loss (Le Bon et al., 2021). However, the current coronavirus pandemic  
246 suggests that in reality, taste loss is its own unique feature of the illness. The present meta-  
247 analysis found an overall taste loss prevalence of 37 % among 138,897 COVID-19-positive  
248 participants, which aligns with other meta-analyses of taste loss prevalence, ranging from 38%  
249 (Agyeman et al., 2020) to 49% (Hajikhani et al., 2020). This high prevalence is not due to  
250 confusion with smell loss because direct taste measures yield similar (or even slightly higher)  
251 prevalence than self-report. Therefore self-reports of taste loss appear to be valid among people  
252 with COVID-19 as they are among other groups (Jang et al., 2021).

253 The COVID-19 pandemic created an urgent need for direct taste measures suitable for the  
254 pandemic research environment, e.g., home testing, and researchers were innovative, but each  
255 developed their own method which makes it difficult to compare results (see Table 1). However,

256 despite the differences in methods, we can draw one general conclusion, which is that the form  
257 of the tastants matters (taste solutions are better than taste strips) but this general conclusion  
258 must be tentative given that the forms of delivery were not compared directly using the same  
259 approach, e.g., thresholds or identification.

260         The present meta-analysis found that around 4 in every 10 COVID-19 patients experience  
261 taste loss. We also found age and sex effects: females experienced higher rates of taste loss than  
262 males, aligning with other meta-analyses reporting a similar effect (von Bartheld et al., 2020;  
263 Amorim Dos Santos et al., 2021; Saniasiaya et al., 2021). Females may be more susceptible to  
264 taste loss because they are in general are more sensitive than males and have more sensory  
265 capacity to lose. Additionally, we found that COVID-19 associated taste loss peaks in middle age  
266 aligning with the general consensus across other COVID-19 meta-analyses (Agyeman et al.,  
267 2020; von Bartheld et al., 2020). Why the youngest and oldest groups report less taste loss than  
268 do middle-age adults is not currently known.

269         Although COVID-19 has intensified awareness of taste loss and furthered chemosensory  
270 research, scientists are still unsure of the biological mechanisms behind this symptom. The  
271 amount of SARS-CoV-2 virus in saliva is positively related to loss of taste: the more virus, the  
272 more taste loss (Huang et al., 2021; Taziki Balajelini et al., 2021), although this observation is  
273 controversial (Jain et al., 2020). Taste cells may be attacked directly by the virus because studies  
274 of expression patterns for ACE2, the receptor protein known to transport the SARS-CoV-2 virus  
275 into cells, and for TMPRSS2, the protein essential for processing the SARS-CoV-2 spike protein,  
276 showed both are expressed in the supporting cells of taste bud (Sakaguchi et al., 2020; Huang et  
277 al., 2021), including taste receptor cells themselves, at least in one patient (Doyle et al., 2021).  
278 There may also be direct effects on the brain that contribute to taste loss (Douaud et al., 2021).

## 279 **Limitations and Future Research**

280 In many of the included articles, clinicians and researchers collected self-reported taste  
281 loss information in tandem with smell loss (e.g., participants responding yes to “Loss of taste and  
282 smell” on a symptom screener), which can confound the results. Therefore, we explored any  
283 differences in how taste loss was collected, reflecting how it was reported in the articles (e.g.,  
284 “smell and taste” vs. “taste only”), and found no significant impact on the prevalence rate (see  
285 Supplementary Materials S3). This result indicates that taste loss is a common and pervasive  
286 symptom of COVID-19. Additionally, far more articles in this meta-analysis used self-report  
287 tests (n = 223) than direct tests (n = 18). This disparity may have prevented us from capturing  
288 significant differences between the two methods.

289 Nearly all of the articles included in this meta-analysis were nonspecific to different  
290 tastes, instead summarizing scores across multiple stimuli (e.g., sweet and salty) and reporting  
291 taste loss as a whole (though in one study participants self-reported taste-specific dysfunction:  
292 salt taste loss, 29.3%; sweet taste, 25.9%; general taste, 34.5% [El Kady et al., 2021]). However,  
293 specific taste sensitivities can be difficult to assess via self-report. Members of the general  
294 population, untrained in chemosensory science, may have difficulties identifying whether or not  
295 they truly lost a specific taste. Therefore, it is important to use direct measures to distinguish  
296 dysfunctions of specific tastes.

297 There are clear opportunities for advancements of standardized direct taste tests to  
298 measure taste loss. Of the 18 studies that used direct tests in general, only five used standardized  
299 tests, representing just 2.06% of the 242 studies examined in this systematic review: Taste Strips  
300 by Burghart Messtechnik (Hintschich et al., 2020a; Le Bon et al., 2021; Niklassen et al., 2021a;  
301 Singer-Cornelius et al., 2021) and the Brief Self-Administered Waterless Empirical Taste Test

302 (SA-WETT) by Sensonics International (Cao et al., 2021). Three other studies using direct tests  
303 were examined during the systematic review but were excluded for not meeting the inclusion  
304 criteria: reporting on a case study (Lee and Lee, 2020), recruiting patients with chemosensory  
305 dysfunction (Le Bon et al., 2020), and not reporting the required taste loss prevalence data (Huart  
306 et al., 2020). Among these three studies, Lee and Lee (2020) used a non-standardized direct test,  
307 and Huart et al. (2020) and Le Bon et al. (2020) used the Taste Strips – although this represents a  
308 missed opportunity to analyze more studies that used standardized tests, including these two  
309 articles would have increased the rate of standardized test use among all 241 studies to just  
310 2.89% (N=7).

311 As of September 2021, 220 million individuals have been infected with SARS-CoV- 2  
312 virus, with large number of recovered individuals with persistent symptoms included taste loss. It  
313 is well documented that disease-related or age-related or chemosensory loss have profound  
314 effects on an individual's quality of life. Unlike other disorders such as vision and hearing for  
315 which preventative and screening guidelines exist (United States Preventive Services Taskforce),  
316 they are not available for taste and smell disorders. The COVID-19 pandemic further highlighted  
317 this existing gap, lack of standardized measures and clinical guidelines for screening, assessing,  
318 and monitoring the taste system making it difficult for clinicians to track progression of disease.  
319 Assessment of taste function in patients with and without confirmed COVID-19 needs to become  
320 standard of practice for clinicians. This is particularly important for at least two reasons: 1)  
321 having baseline measures help clinicians assess trends over time, and 2) given the  
322 interrelatedness between the sense of smell and taste, objective measures of taste collected  
323 during clinical assessments may help dissociate whether it is a smell or taste problem. For  
324 patients who report changes in taste function during screening questionnaires, full testing with

325 standardized objective chemosensory tools should be performed. It is critical that clinicians are  
326 aware that most patients with chemosensory dysfunction complain of taste alterations, therefore,  
327 a closer inquiry of patient's reports regarding the specific taste quality (i.e., sweet, bitter, sour,  
328 and salt, fat) affected is important to further distinguish between taste and perception of flavor.

### 329 **Conclusion**

330         The COVID-19 pandemic demanded an urgent response from scientists and clinicians,  
331 who have been working to understand the novel virus and the symptoms it inflicts. Of the many  
332 unique features of this virus, smell and taste dysfunctions are among the most prominent. Yet as  
333 taste loss joined smell loss as a more prolific topic in scientific literature, many initially  
334 speculated that taste loss rates were overestimated due to confusion between taste and smell in  
335 self-reports. However, our meta-analysis found a prevalence rate for taste loss of 36.9% among  
336 138,897 COVID-19-positive individuals (95% CI: 33.27–40.69%), supported by direct methods,  
337 reflecting the validity of this distinct symptom. Dysfunction in the sense of taste was, and still is,  
338 a difficult reality for millions of people affected by the virus and merits further research to fully  
339 understand the mechanisms behind this phenomenon and how to properly assess and address it.  
340 Among the population of 138,897 COVID-19-positive individuals included in this meta-analysis,  
341 only 257 of them, across five separate studies, were assessed using standardized taste tests.  
342 Future research should include the development of fast and accurate taste tests, studies that  
343 measure taste and smell function separately to dissociate olfacto-gustatory interactions, as well  
344 as the employment of standardized taste tests in clinical settings to examine taste dysfunction. In  
345 addition, clinical trials are needed to elucidate frequency of screening and age at which to start  
346 and stop screening for chemosensory disorders. Finally, more mechanistic studies to understand



347 taste and smell disorders associated with COVID-19 to aid in developing new therapeutic  
348 options for patients with long-lasting impairment of their chemical senses.

### 349 **Included Articles**

350 In total, 241 were included in the present systematic review and meta-analysis (Gamper  
351 et al., 2012; Liu et al., 2016; Adamczyk et al., 2020; Adedeji et al., 2020; Adorni et al., 2020;  
352 Aggarwal et al., 2020; Al-Ani and Acharya, 2020; Altin et al., 2020; Andrews et al., 2020; Anna  
353 et al., 2020; Asai et al., 2020; Bastiani et al., 2020; Beltran-Corbellini et al., 2020; Bergquist et  
354 al., 2020; Biadsee et al., 2020; Bidkar et al., 2020b; Boscolo-Rizzo et al., 2020a, 2020b, 2020c,  
355 2021; Boudjema et al., 2020; Brandao Neto et al., 2020; Bulgurcu et al., 2020; Calica Utku et al.,  
356 2020; Carignan et al., 2020; Chary et al., 2020; Chen et al., 2020; Chiesa-Estomba et al., 2020;  
357 Cho et al., 2020; Chung et al., 2020; Cocco et al., 2020; Dawson et al., 2020; Dell'Era et al.,  
358 2020; De Maria et al., 2020; Durrani et al., 2020; Elimian et al., 2020; Farah Yusuf Mohamud et  
359 al., 2020; Fistera et al., 2020; Fontanet et al., 2020; Freni et al., 2020; Garg et al., 2020; Gelardi  
360 et al., 2020; Giacomelli et al., 2020; Gorzkowski et al., 2020; Gozen et al., 2020; Güner et al.,  
361 2020; Gudbjartsson et al., 2020; Guillen Martinez et al., 2020; Haehner et al., 2020; Hintschich  
362 et al., 2020b; Horvath et al., 2020; Iversen et al., 2020; Izquierdo-Domínguez et al., 2020; Jain et  
363 al., 2020; Jalessi et al., 2020; Kacem et al., 2020; Karadas et al., 2020; Kempker et al., 2020;  
364 Kim et al., 2020; Klopfenstein et al., 2020; Konstantinidis et al., 2020; Krishnasamy et al., 2020;  
365 Kronborg et al., 2020; Kumar et al., 2020, 2021a, 2021b; Lagi et al., 2020; Lan et al., 2020;  
366 Lapostolle et al., 2020; La Torre et al., 2020; Lechien et al., 2020b, 2020c, 2020a, 2021, 2021,  
367 2021; Lechner et al., 2020; Lee et al., 2020a, 2020b, 2021; Levinson et al., 2020; Liang et al.,  
368 2020; Liguori et al., 2020; Lindahl et al., 2020; Lombardi et al., 2020, 2021; Luers et al., 2020;  
369 Lv et al., 2020; Maechler et al., 2020; Magnavita et al., 2020; Makda et al., 2020; Mao et al.,

370 2020; Martin-Sanz et al., 2020; Mazzatenta et al., 2020; Meini et al., 2020; Menni et al., 2020;  
371 Mercante et al., 2020; Merkely et al., 2020; Merza et al., 2020; Moein et al., 2020; Morshed et  
372 al., 2020; Murat et al., 2020; Nakagawa et al., 2020, 2020; Nakakubo et al., 2020; Nakanishi et  
373 al., 2020; Noh et al., 2020; Otte et al., 2020; Ozelik Korkmaz et al., 2020; Paderno et al., 2020;  
374 Patel et al., 2020; Perlman et al., 2020; Petersen et al., 2020; Petrocelli et al., 2020b, 2020a;  
375 Pinna et al., 2020; Printza et al., 2020, 2021; Qiu et al., 2020; Rajkumar et al., 2020; Ramteke et  
376 al., 2020, 2020; Rojas-Lechuga et al., 2020; Roland et al., 2020; Romero-Sanchez et al., 2020;  
377 Rubel et al., 2020; Sakalli et al., 2020; Salepci et al., 2020; Sayin et al., 2020a, 2020b; Schirinzi  
378 et al., 2020; Schmithausen et al., 2020; Seo et al., 2020; Shoer et al., 2020; Sierpinski et al.,  
379 2020; Smith et al., 2020; Somekh et al., 2020; Speth et al., 2020; Spinato et al., 2020; Stavem et  
380 al., 2020; Sun et al., 2020; Teklu et al., 2020; Tham et al., 2020; Tsivgoulis et al., 2020; Tudrej et  
381 al., 2020; Vacchiano et al., 2020; Vaira et al., 2020e, 2020f, 2020b, 2020c, 2020d, 2020a, 2020b;  
382 Van Loon et al., 2020; Vena et al., 2020; Villarreal et al., 2020; Wagner et al., 2020; Waterfield  
383 et al., 2020; Wee et al., 2020; Weiss et al., 2020; Weng et al., 2020; Yan et al., 2020a, 2020b;  
384 Zayet et al., 2020a, 2020b; Zimmerman et al., 2020; Akinbami et al., 2021; Alharbi et al., 2021;  
385 AlShakhs et al., 2021; Amano et al., 2021; Arslan et al., 2021; Ashrafi et al., 2021; Bagnasco et  
386 al., 2021; Barillari et al., 2021; Besli et al., 2021; Bianco et al., 2021; Breyer et al., 2021;  
387 Callejon-Leblic et al., 2021; Cao et al., 2021; Carcamo Garcia et al., 2021; Concheiro-Guisan et  
388 al., 2021; Dini et al., 2021; Dixon et al., 2021; El Kady et al., 2021; Fisher et al., 2021; Fleischer  
389 et al., 2021; Galluzzi et al., 2021; Gianola et al., 2021; Gibbons et al., 2021; Gonzalez et al.,  
390 2021; Gupta et al., 2021a, 2021b; Gurrola et al., 2021; Hijazi et al., 2021; Ismail et al., 2021;  
391 Jethani et al., 2021; Kamel et al., 2021; Kandakure et al., 2021; Karaarslan et al., 2021; Karni et  
392 al., 2021; Kavaz et al., 2021; Ladoire et al., 2021; Lampl et al., 2021; Makaronidis et al., 2021;

393 Mannan et al., 2021; Moeller et al., 2021; Monti et al., 2021; Morlock et al., 2021; Niklassen et  
394 al., 2021b; Ninchritz-Becerra et al., 2021; Nishanth et al., 2021; Noviello et al., 2021; Oda et al.,  
395 2021; Omezli and Torul, 2021; O’Keefe et al., 2021; O’Sullivan et al., 2021; Ozcan et al., 2021;  
396 Parcha et al., 2021; Perula de Torres et al., 2021; Polat et al., 2021; Rodebaugh et al., 2021;  
397 Rousseau et al., 2021; Sahoo et al., 2021; Salcan et al., 2021; Savtale et al., 2021; Sbrana et al.,  
398 2021; Schwab et al., 2021; Sehanobish et al., 2021; Singer-Cornelius et al., 2021; Soh et al.,  
399 2021; Song et al., 2021a, 2021b; Tarifi et al., 2021; Thakur et al., 2021; Trachootham et al.,  
400 2021; İşlek and Balcı, 2021; van den Besselaar et al., 2021; Wierdsma et al., 2021; Yadav et al.,  
401 2021; Zejda et al., 2021; Zifko et al., 2021; Zou et al., 2021)

#### 402 **Funding**

403 Dr. Mackenzie Hannum is supported by NIH T32 funding (DC000014). Dr. Joseph is  
404 supported by the National Institute of Alcohol Abuse and Alcoholism under award number,  
405 Z01AA000135 and the National Institute of Nursing Research, the NIH Office of Workforce  
406 Diversity, National Institutes of Health Distinguished Scholar, and the Rockefeller University  
407 Heilbrunn Nurse Scholar Award. Dr. Reed is supported in part through NIH U01DC019578.

#### 408 **Acknowledgements**

409 We acknowledge Michael G. Tordoff for his assistance with the review of the literature,  
410 Sarah Lipson for her initial effort in reading the earlier articles, and Akane Kikuchi for ensuring  
411 no article was missed. Additionally, we thank Gary K. Beauchamp for his comments on the  
412 manuscript.

413

414 **Captions**

415 **Figure 1.** CONSORT flow diagram demonstrating the article selection process for this  
416 systematic review and meta-analysis.

417

418 **Table 1.** Overview of the direct approaches to assess taste loss in participants.

419 **Figure 2.** Orchard plot following the guidelines outlined by Nakagawa et al. The point estimate  
420 of the pooled prevalence (trunk) is represented by the bold blue or red dot. The confidence  
421 interval of the pooled prevalence estimate (branch) is represented by the bold black line, and the  
422 prediction interval (twig) is represented by the thin black line. Individual prevalence estimates  
423 from each study are represented by the scattered colored points (slightly transparent circles,  
424 called fruits). Each fruit is scaled by the precision of the point estimate of prevalence for each  
425 study, or the inverse of the standard error.


426

427 **Table 2.** Random-effect estimate of age group on COVID-19 taste loss prevalence using  
428 generalized linear mixed models.

429

430 **Table 3.** Random-effect estimate of direct testing type on COVID-19 taste loss prevalence using  
431 generalized linear mixed models.

432



433  
434

435 **Figure 1.** CONSORT flow diagram demonstrating the article selection process for this  
436 systematic review and meta-analysis.

437 **Table 1. Overview of the direct approaches to assess taste loss in participants.**

<i>Test</i>	<i>Article</i>	<i>Test Method</i>	<i>Test Objective</i>	<i>Taste Quality Measured</i>				
				<i>Salty</i>	<i>Sweet</i>	<i>Sour</i>	<i>Bitter</i>	<i>Umami</i>
Four-solution test	Vaira et al., 2020a–2020f Petrocelli et al., 2020 <sup>A</sup>	1 mL of each solution, plus deionized water as control, placed on the participant's tongue via cotton swab. Quarantined patients prepared their own solutions.	Identification	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
Four-solution test & Taste Strips	Altin et al., 2020 Salcan et al., 2021	Participants swallowed and identified the solution. Next, paper strips dipped into each solution and placed on the participant's tongue.	Identification Duration	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
Two-solution test (1)	Bidkar et al., 2020	Two drops (2 mL) of each solution placed on the participant's tongue via pipette.	Identification	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Two-solution test (2)	Mazzatenta et al., 2020	200 µl of each solution dropped onto participant's tongue.	Detection	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>
Taste Strips by BMG <sup>B</sup>	Hintschich et al., 2020 Singer-Cornelius et al., 2021 Le Bon et al., 2021 Niklassen et al., 2021 <sup>C</sup>	Taste strips (provided by Burghart Messtechnik GmbH) placed on the participant's tongue.	Identification and threshold concentration	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>

Taste sprays	Niklassen et al., 2021 <sup>C</sup>	Each solution sprayed onto participant's tongue.	Identification	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
Tastant capsules	Adamczyk et al., 2020	Each of ten 0.33 mL gelatin capsules (one tasteless and nine with tastant) dissolved on participant's tongue.	Description	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
Brief Self-Administered Waterless Empirical Taste Test (SA-WETT) <sup>B</sup>	Cao et al., 2021	27 disposable plastic strips that contain dried solutions of each taste (and some tasteless strips) self-placed on participant's tongue.	Identification	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
Chemosensory Test by India Protocol	Ramteke et al., 2020	Authors used coconut oil, chocolates, and flavored milk to test smell and taste function. No further details are provided.	N/A	N/A				

438 <sup>A</sup> These studies used a sweet solution concentration that is double what was used in the other four-solution tests.

439 <sup>B</sup> Validated test.

440 <sup>C</sup> Niklassen et al. (2021) used both Taste Strips and taste sprays with participants.

441


442

443

444

445

446



447  
448 **Figure 2.** Orchard plot of taste loss and COVID-19. The point estimate of the pooled prevalence  
449 (trunk) is represented by the bold turquoise or pink dot. The confidence interval of the pooled  
450 prevalence estimate (branch) is represented by the bold black line, and the prediction interval  
451 (twig) is represented by the thin black line. Individual prevalence estimates from each study are  
452 represented by the scattered colored points (slightly transparent circles). Each point is scaled by  
453 the precision of the point estimate of prevalence for each study, i.e., inverse of the standard error.  
454



455 **Table 2.** Random-effect estimate of age group on COVID-19 taste loss prevalence using  
456 generalized linear mixed models.  
457

<i>Age Category</i>	<i>k</i>	<i>Proportion</i>	<i>95%-CI</i>	<i>Q</i>	<i>I<sup>2</sup></i>	<i>Tau<sup>2</sup></i>
<i>Adolescent</i>	9	0.12	0.06-0.20	382.58	97.90%	0.828
<i>Young Adults</i>	20	0.32	0.1984-0.4640	1482.59	98.70%	1.961
<i>Middle Age</i>	118	0.44	0.3864-0.4954	10517.56	98.90%	1.4649
<i>Eldery</i>	8	0.18	0.0803-0.3570	193.64	96.40%	1.6031
<i>Older</i>	55	0.35	0.2882-0.4067	1522.64	96.50%	0.9307

458  
459  
460

461 **Table 3.** Random-effect estimate of direct testing type on COVID-19 taste loss prevalence using  
462 generalized linear mixed models.

463

<i>Direct Testing Type</i>	<i>k</i>	<i>Proportion</i>	<i>95%-CI</i>	<i>Q</i>	<i>I<sup>2</sup></i>	<i>Tau<sup>2</sup></i>
<i>Solution</i>	11	0.5602	0.4666-0.6497	87.62	88.60%	0.3566
<i>Strip</i>	5	0.2183	0.1510-0.3048	14.45	72.30%	0.1663
<i>Other</i>	2	0.0539	0.0000-0.9868	0	0.00%	17.9895

464

## References

- 465  
466
- 467 Abdullahi, A., Candan, S.A., Abba, M.A., Bello, A.H., Alshehri, M.A., Afamefuna Victor, E., Umar,  
468 N.A., and Kundakci, B. 2020. Neurological and Musculoskeletal Features of COVID-19: A  
469 Systematic Review and Meta-Analysis. *Front Neurol.* 11:687.
- 470 Adamczyk, K., Herman, M., Frączek, J., Piec, R., Szykuła-Piec, B., Zaczyński, A., Wójtowicz, R.,  
471 Bojanowski, K., Rusyan, E., Król, Z., et al. 2020. Sensitivity and specificity of prediction models  
472 based on gustatory disorders in diagnosing COVID-19 patients: a case-control study. *Infectious*  
473 *Diseases (except HIV/AIDS)*.
- 474 Adedeji, I.A., Abdu, Y.M., Bashir, M.F., Adamu, A.S., Gwarzo, G.D., Yaro, B.S., Musa, A.A.,  
475 Hassan, Z.I., Maigoro, A.M., and Jibrin, Y.B. 2020. Profile of children with COVID-19 infection: a  
476 cross sectional study from North-East Nigeria. *Pan Afr Med J.* 35:145.
- 477 Adorni, F., Prinelli, F., Bianchi, F., Giacomelli, A., Pagani, G., Bernacchia, D., Rusconi, S., Maggi,  
478 S., Trevisan, C., Noale, M., et al. 2020. Self-reported symptoms of SARS-CoV-2 infection in a  
479 non-hospitalized population: results from the large Italian web-based EPICoVID19 cross-  
480 sectional survey. *JMIR Public Health Surveill.*
- 481 Aggarwal, S., Garcia-Telles, N., Aggarwal, G., Lavie, C., Lippi, G., and Henry, B.M. 2020. Clinical  
482 features, laboratory characteristics, and outcomes of patients hospitalized with coronavirus  
483 disease 2019 (COVID-19): Early report from the United States. *Diagn Berl.* 7:91–96.
- 484 Agyeman, A.A., Chin, K.L., Landersdorfer, C.B., Liew, D., and Ofori-Asenso, R. 2020. Smell and  
485 Taste Dysfunction in Patients With COVID-19: A Systematic Review and Meta-analysis. *Mayo*  
486 *Clin Proc.* 95:1621–1631.
- 487 Akinbami, L.J., Petersen, L.R., Sami, S., Vuong, N., Lukacs, S.L., Mackey, L., Atas, J., and LaFleur,  
488 B.J. 2021. COVID-19 symptoms and SARS-CoV-2 antibody positivity in a large survey of first  
489 responders and healthcare personnel, May-July 2020. *Clin Infect Dis.*
- 490 Al-Ani, R.M., and Acharya, D. 2020. Prevalence of Anosmia and Ageusia in Patients with COVID-  
491 19 at a Primary Health Center, Doha, Qatar. *Indian J Otolaryngol Head Neck Surg.* 1–7.
- 492 Alharbi, H., You, S., and Katz, J. 2021. Should anosmia and dysgeusia be a concern for oral and  
493 maxillofacial surgeons during the COVID-19 pandemic? *Oral Maxillofac Surg.*
- 494 AlShakhs, A., Almomen, A., AlYaeesh, I., AlOmairin, A., AlMutairi, A.A., Alammam, Z., Almomen,  
495 H., and Almomen, Z. 2021. The Association of Smell and Taste Dysfunction with COVID19, And  
496 Their Functional Impacts. *Indian J Otolaryngol Head Neck Surg.* 1–6.
- 497 Altin, F., Cingi, C., Uzun, T., and Bal, C. 2020. Olfactory and gustatory abnormalities in COVID-19  
498 cases. *Eur Arch Otorhinolaryngol.* 277:2775–2781.

- 499 Amano, Y., Kage, H., Tanaka, G., Gono, W., Nakai, Y., Kurokawa, R., Inui, S., Okamoto, K.,  
500 Harada, S., Iwabuchi, M., et al. 2021. Diagnostic prediction of COVID-19 based on clinical and  
501 radiological findings in a relatively low COVID-19 prevalence area. *Respir Investig.*
- 502 Amorim Dos Santos, J., Normando, A.G.C., Carvalho da Silva, R.L., Acevedo, A.C., De Luca Canto,  
503 G., Sugaya, N., Santos-Silva, A.R., and Guerra, E.N.S. 2021. Oral Manifestations in Patients with  
504 COVID-19: A Living Systematic Review. *J Dent Res.* 100:141–154.
- 505 Andrews, P.J., Pendolino, A.L., Ottaviano, G., Scarpa, B., Grant, J., Gaudio, P., Bordin, A.,  
506 Marchese-Ragona, R., Leoni, D., Cattelan, A., et al. 2020. Olfactory and taste dysfunction among  
507 mild-to-moderate symptomatic COVID-19 positive health care workers: An international survey.  
508 *Laryngoscope Investig Otolaryngol.* 5:1019–1028.
- 509 Anna, F., Goyard, S., Lalanne, A.L., Nevo, F., Gransagne, M., Souque, P., Louis, D., Gillon, V.,  
510 Turbiez, I., Bidard, F.-C., et al. 2020. High seroprevalence but short-lived immune response to  
511 SARS-CoV-2 infection in Paris. *Eur J Immunol.* n/a.
- 512 Arslan, G., Aktürk, H., and Duman, M. 2021. Clinical Characteristics of Pediatric COVID-19 and  
513 Predictors of PCR Positivity. *Pediatr Int.*
- 514 Asai, N., Sakanashi, D., Nakamura, A., Kishino, T., Kato, H., Hagihara, M., Shiota, A., Koizumi, Y.,  
515 Yamagishi, Y., and Mikamo, H. 2020. Clinical manifestations and radiological features by chest  
516 computed tomographic findings of a novel coronavirus disease-19 pneumonia among 92  
517 patients in Japan. *J Microbiol Immunol Infect.*
- 518 Ashrafi, F., Ommi, D., Zali, A., Khani, S., Soheili, A., Arab-Ahmadi, M., Behnam, B., Nohesara, S.,  
519 Semnani, F., Fatemi, A., et al. 2021. Neurological Manifestations and their Correlated Factors in  
520 COVID-19 Patients; a Cross-Sectional Study. *Arch Acad Emerg Med.* 9:e34.
- 521 Bagnasco, D., Passalacqua, G., Braido, F., Tagliabue, E., Cosini, F., Filastro, M., Ioppi, A.,  
522 Carobbio, A., Mocellin, D., Riccio, A.M., et al. 2021. Quick Olfactory Sniffin' Sticks Test (Q-Sticks)  
523 for the detection of smell disorders in COVID-19 patients. *World Allergy Organ J.* 14:100497.
- 524 Bakbergenuly, I., and Kulinskaya, E. 2018. Meta-analysis of binary outcomes via generalized  
525 linear mixed models: a simulation study. *BMC Med Res Methodol.* 18:70.
- 526 Barillari, M.R., Bastiani, L., Lechien, J.R., Mannelli, G., Molteni, G., Cantarella, G., Coppola, N.,  
527 Costa, G., Trecca, E.M.C., Grillo, C., et al. 2021. A structural equation model to examine the  
528 clinical features of mild-to-moderate COVID-19: A multicenter Italian study. *J Med Virol.*  
529 93:983–994.
- 530 Bartheld, C.S. von, Hagen, M.M., and Butowt, R. 2020. Prevalence of Chemosensory  
531 Dysfunction in COVID-19 Patients: A Systematic Review and Meta-analysis Reveals Significant  
532 Ethnic Differences. *ACS Chem Neurosci.* 11:2944–2961.

- 533 Bastiani, L., Fortunato, L., Pieroni, S., Bianchi, F., Adorni, F., Prinelli, F., Giacomelli, A., Pagani, G.,  
534 Maggi, S., Trevisan, C., et al. 2020. EPICOV19: Psychometric assessment and validation of a  
535 short diagnostic scale for a rapid Covid-19 screening based on reported symptoms. *J Med*  
536 *Internet Res.*
- 537 Beltran-Corbellini, A., Chico-Garcia, J.L., Martinez-Poles, J., Rodriguez-Jorge, F., Natera-Villalba,  
538 E., Gomez-Corral, J., Gomez-Lopez, A., Monreal, E., Parra-Diaz, P., Cortes-Cuevas, J.L., et al.  
539 2020. Acute-onset smell and taste disorders in the context of COVID-19: a pilot multicentre  
540 polymerase chain reaction based case-control study. *Eur J Neurol.*
- 541 Bergquist, S.H., Partin, C., Roberts, D.L., O'Keefe, J.B., Tong, E.J., Zreloff, J., Jarrett, T.L., and  
542 Moore, M.A. 2020. Non-hospitalized Adults with COVID-19 Differ Noticeably from Hospitalized  
543 Adults in Their Demographic, Clinical, and Social Characteristics. *SN Compr Clin Med.* 1–9.
- 544 Besli, G.E., Öcal Demir, S., Girit, S., Arman, T., Duyu, M., and Arslanoglu, S. 2021. COVID-19 in  
545 children: A single center experience from Istanbul, Turkey. *Med J BAKIRKOY.* 17:64–71.
- 546 Besselaar, J.H. van den, Sikkema, R.S., Koene, F., Buul, L.W. van, Oude Munnink, B.B., Frénay, I.,  
547 Te Witt, R., Koopmans, M.P.G., Hertogh, C., and Buurman, B.M. 2021. Are presymptomatic  
548 SARS-CoV-2 infections in nursing home residents unrecognized symptomatic infections?  
549 Sequence and metadata from weekly testing in an extensive nursing home outbreak. *Age*  
550 *Ageing.*
- 551 Biadsee, A., Biadsee, A., Kassem, F., Dagan, O., Masarwa, S., and Ormianer, Z. 2020. Olfactory  
552 and Oral Manifestations of COVID-19: Sex-Related Symptoms-A Potential Pathway to Early  
553 Diagnosis. *Otolaryngol Head Neck Surg.* 194599820934380.
- 554 Bianco, M.R., Modica, D.M., Drago, G.D., Azzolina, A., Mattina, G., De Natale, M., Rossi, G.,  
555 Amata, M., Canzoneri, G., Manganaro, G., et al. 2021. Alteration of Smell and Taste in  
556 Asymptomatic and Symptomatic COVID-19 Patients in Sicily, Italy. *Ear Nose Throat J.* 100:182S-  
557 185S.
- 558 Bidkar, V., Mishra, M., Selvaraj, K., Joshi, P., H, S.B., Dabhekar, S., Prathipati, K.K., Rathod, B.S.,  
559 Shendre, P., and Gondode, P. 2020a. Testing Olfactory and Gustatory Dysfunctions among  
560 Quarantine COVID-19 Suspects. *Indian J Otolaryngol Head Neck Surg Off Publ Assoc Otolaryngol*  
561 *India.* 1–6.
- 562 Bidkar, V., Mishra, M., Selvaraj, K., Joshi, P., H, S.B., Dabhekar, S., Prathipati, K.K., Rathod, B.S.,  
563 Shendre, P., and Gondode, P. 2020b. Testing Olfactory and Gustatory Dysfunctions among  
564 Quarantine COVID-19 Suspects. *Indian J Otolaryngol Head Neck Surg.* 1–6.
- 565 Blomberg, B., Mohn, K.G.-I., Brokstad, K.A., Zhou, F., Linchausen, D.W., Hansen, B.-A., Lartey, S.,  
566 Onyango, T.B., Kuwelker, K., Sævik, M., et al. 2021. Long COVID in a prospective cohort of  
567 home-isolated patients. *Nat Med.* 1–7.

- 568 Boscolo-Rizzo, P., Borsetto, D., Fabbris, C., Spinato, G., Frezza, D., Menegaldo, A., Mularoni, F.,  
569 Gaudio, P., Cazzador, D., Marciani, S., et al. 2020a. Evolution of Altered Sense of Smell or  
570 Taste in Patients With Mildly Symptomatic COVID-19. *JAMA Otolaryngol Head Neck Surg*.
- 571 Boscolo-Rizzo, P., Borsetto, D., Spinato, G., Fabbris, C., Menegaldo, A., Gaudio, P., Nicolai, P.,  
572 Tirelli, G., Da Mosto, M.C., Rigoli, R., et al. 2020b. New onset of loss of smell or taste in  
573 household contacts of home-isolated SARS-CoV-2-positive subjects. *Eur Arch Otorhinolaryngol*.
- 574 Boscolo-Rizzo, P., Guida, F., Polesel, J., Marcuzzo, A.V., Antonucci, P., Capriotti, V., Sacchet, E.,  
575 Cragnolini, F., D'Alessandro, A., Zanelli, E., et al. 2021. Self-reported smell and taste recovery in  
576 COVID-19 patients: a one-year prospective study. *MedRxiv*. 2021.03.18.21253862.
- 577 Boscolo-Rizzo, P., Polesel, J., Spinato, G., Fabbris, C., Calvanese, L., Menegaldo, A., Borsetto, D.,  
578 and Hopkins, C. 2020c. Predominance of an altered sense of smell or taste among long-lasting  
579 symptoms in patients with mildly symptomatic COVID-19. *Rhinology*.
- 580 Boscutti, A., Delvecchio, G., Pighi, A., Cereda, G., Ciappolino, V., Bellani, M., Fusar-Poli, P., and  
581 Brambilla, P. 2021. OLFATORY AND GUSTATORY DYSFUNCTIONS IN SARS-CoV-2 INFECTION:  
582 *Brain Behav Immun - Health*. 100268.
- 583 Boudjema, S., Finance, J., Coulibaly, F., Meddeb, L., Tissot-Dupont, H., Michel, M., Lagier, J.C.,  
584 Million, M., Radulesco, T., Michel, J., et al. 2020. Olfactory and gustative disorders for the  
585 diagnosis of COVID-19. *Travel Med Infect Dis*. 37:101875.
- 586 Brandao Neto, D., Fornazieri, M.A., Dib, C., Di Francesco, R.C., Doty, R.L., Voegels, R.L., and  
587 Pinna, F.R. 2020. Chemosensory Dysfunction in COVID-19: Prevalences, Recovery Rates, and  
588 Clinical Associations on a Large Brazilian Sample. *Otolaryngol Head Neck Surg*.  
589 194599820954825.
- 590 Breyer, M.K., Breyer-Kohansal, R., Hartl, S., Kundi, M., Weseslindtner, L., Stiasny, K.,  
591 Puchhammer-Stockl, E., Schrott, A., Fodinger, M., Binder, M., et al. 2021. Low SARS-CoV-2  
592 seroprevalence in the Austrian capital after an early governmental lockdown. *Sci Rep*.  
593 11:10158.
- 594 Bulgurcu, S., Oztutgan, T., Baz, E., Yonem, A., Koc, N.G., Erkul, E., and Cekin, E. 2020.  
595 Assessment of Smell and Taste Disorders in COVID-19: A Cross-sectional Study. *J Craniofac Surg*.
- 596 Calica Utku, A., Budak, G., Karabay, O., Guclu, E., Okan, H.D., and Vatan, A. 2020. Main  
597 symptoms in patients presenting in the COVID-19 period. *Scott Med J*. 36933020949253.
- 598 Callejon-Leblic, M.A., Moreno-Luna, R., Del Cuvillo, A., Reyes-Tejero, I.M., Garcia-Villaran, M.A.,  
599 Santos-Pena, M., Maza-Solano, J.M., Martin-Jimenez, D.I., Palacios-Garcia, J.M., Fernandez-  
600 Velez, C., et al. 2021. Loss of Smell and Taste Can Accurately Predict COVID-19 Infection: A  
601 Machine-Learning Approach. *J Clin Med*. 10.

- 602 Cao, A.C., Nimmo, Z.M., Mirza, N., Cohen, N.A., Brody, R.M., and Doty, R.L. 2021. Objective  
603 screening for olfactory and gustatory dysfunction during the COVID-19 pandemic: A prospective  
604 study in healthcare workers using self-administered testing. *World J Otorhinolaryngol - Head*  
605 *Neck Surg.* S2095881121000159.
- 606 Carcamo Garcia, M.H., Garcia Choza, D.D., Salazar Linares, B.J., and Diaz, M.M. 2021.  
607 Neurological manifestations of patients with mild-to-moderate COVID-19 attending a public  
608 hospital in Lima, Peru. *ENeurologicalSci.* 23.
- 609 Carignan, A., Valiquette, L., Grenier, C., Musonera, J.B., Nkengurutse, D., Marcil-Heguy, A.,  
610 Vettese, K., Marcoux, D., Valiquette, C., Xiong, W.T., et al. 2020. Anosmia and dysgeusia  
611 associated with SARS-CoV-2 infection: an age-matched case-control study. *CMAJ.* 192:E702–  
612 E707.
- 613 Chary, E., Carsuzaa, F., Trijolet, J.P., Capitaine, A.L., Roncato-Saberan, M., Fouet, K., Cazenave-  
614 Roblot, F., Catroux, M., Allix-Beguec, C., and Dufour, X. 2020. Prevalence and Recovery From  
615 Olfactory and Gustatory Dysfunctions in Covid-19 Infection: A Prospective Multicenter Study.  
616 *Am J Rhinol Allergy.* 34:686–693.
- 617 Chen, A., Agarwal, A., Ravindran, N., To, C., Zhang, T., and Thuluvath, P.J. 2020. Are  
618 Gastrointestinal Symptoms Specific for COVID-19 Infection? A Prospective Case-Control Study  
619 from the United States. *Gastroenterology.*
- 620 Chiesa-Estomba, C.M., Lechien, J.R., Portillo-Mazal, P., Martinez, F., Cuauero-Sanchez, J., Calvo-  
621 Henriquez, C., and Saussez, S. 2020. Olfactory and gustatory dysfunctions in COVID-19. First  
622 reports of Latin-American ethnic patients. *Am J Otolaryngol.* 41:102605.
- 623 Cho, R.H., To, Z.W., Yeung, Z.W., Tso, E.Y., Fung, K.S., Chau, S.K., Leung, E.Y., Hui, T.S., Tsang,  
624 S.W., Kung, K.N., et al. 2020. COVID-19 Viral Load in the Severity of and Recovery from Olfactory  
625 and Gustatory Dysfunction. *Laryngoscope.*
- 626 Chung, T.W., Sridhar, S., Zhang, A.J., Chan, K.H., Li, H.L., Wong, F.K., Ng, M.Y., Tsang, R.K., Lee,  
627 A.C., Fan, Z., et al. 2020. Olfactory Dysfunction in Coronavirus Disease 2019 Patients:  
628 Observational Cohort Study and Systematic Review. *Open Forum Infect Dis.* 7:ofaa199.
- 629 Cocco, A., Amami, P., Desai, A., Voza, A., Ferreli, F., and Albanese, A. 2020. Neurological  
630 features in SARS-CoV-2-infected patients with smell and taste disorder. *J Neurol.*
- 631 Concheiro-Guisan, A., Fiel-Ozores, A., Novoa-Carballal, R., González-Duran, M.L., Portugués de  
632 la Red, M., Martínez-Reglero, C., Fernández-Pinilla, I., and González-Guijarro, I. 2021. Subtle  
633 olfactory dysfunction after SARS-CoV-2 virus infection in children. *Int J Pediatr*  
634 *Otorhinolaryngol.* 140.
- 635 Dawson, P., Rabold, E.M., Laws, R.L., Connors, E.E., Gharpure, R., Yin, S., Buono, S., Dasu, T.,  
636 Bhattacharyya, S., Westergaard, R.P., et al. 2020. Loss of Taste and Smell as Distinguishing  
637 Symptoms of COVID-19. *MedRxiv.* 2020.05.13.20101006.

- 638 De Maria, A., Varese, P., Dentone, C., Barisione, E., and Bassetti, M. 2020. High prevalence of  
639 olfactory and taste disorder during SARS-CoV-2 infection in outpatients. *J Med Virol*.
- 640 Deems, D.A., Doty, R.L., Settle, R.G., Moore-Gillon, V., Shaman, P., Mester, A.F., Kimmelman,  
641 C.P., Brightman, V.J., and Snow, J.B. 1991. Smell and taste disorders, a study of 750 patients  
642 from the University of Pennsylvania Smell and Taste Center. *Arch Otolaryngol Head Neck Surg*.  
643 117:519–528.
- 644 Dell’Era, V., Farri, F., Garzaro, G., Gatto, M., Aluffi Valletti, P., and Garzaro, M. 2020. Smell and  
645 taste disorders during COVID-19 outbreak: A cross-sectional study on 355 patients. *Head Neck*.
- 646 Dini, G., Montecucco, A., Rahmani, A., Barletta, C., Pellegrini, L., Debarbieri, N., Orsi, A.,  
647 Caligiuri, P., Varesano, S., Manca, A., et al. 2021. Clinical and epidemiological characteristics of  
648 COVID-19 during the early phase of the SARS-CoV-2 pandemic: a cross-sectional study among  
649 medical school physicians and residents employed in a regional reference teaching hospital in  
650 Northern Italy. *Int J Occup Med Env Health*.
- 651 Dixon, B.E., Wools-Kaloustian, K.K., Fadel, W.F., Duszynski, T.J., Yiannoutsos, C., Halverson, P.K.,  
652 and Menachemi, N. 2021. Symptoms and symptom clusters associated with SARS-CoV-2  
653 infection in community-based populations: Results from a statewide epidemiological study.  
654 *PLoS One*. 16:e0241875.
- 655 Douaud, G., Lee, S., Alfaro-Almagro, F., Arthofer, C., Wang, C., McCarthy, P., Lange, F.,  
656 Andersson, J.L.R., Griffanti, L., Duff, E., et al. 2021. Brain imaging before and after COVID-19 in  
657 UK Biobank.
- 658 Doyle, M.E., Appleton, A., Liu, Q.-R., Yao, Q., Mazucanti, C.H., and Egan, J.M. 2021. Human Type  
659 II Taste Cells Express Angiotensin-Converting Enzyme 2 and Are Infected by Severe Acute  
660 Respiratory Syndrome Coronavirus 2 (SARS-CoV-2). *Am J Pathol*. 191:1511–1519.
- 661 Durrani, M., Haq, I.U., Kalsoom, U., and Yousaf, A. 2020. Chest X-rays findings in COVID 19  
662 patients at a University Teaching Hospital - A descriptive study. *Pak J Med Sci*. 36:S22–S26.
- 663 El Kady, D.M., Gomaa, E.A., Abdella, W.S., Ashraf Hussien, R., Abd ElAziz, R.H., and Khater,  
664 A.G.A. 2021. Oral manifestations of COVID -19 patients: An online survey of the Egyptian  
665 population. *Clin Exp Dent Res*. cre2.429.
- 666 Elimian, K.O., Ochu, C.L., Ebhodaghe, B., Myles, P., Crawford, E.E., Igumbor, E., Ukponu, W.,  
667 Olayinka, A., Aruna, O., Dan-Nwafor, C., et al. 2020. Patient characteristics associated with  
668 COVID-19 positivity and fatality in Nigeria: retrospective cohort study. *BMJ Open*. 10:e044079.
- 669 Farah Yusuf Mohamud, M., Garad Mohamed, Y., Mohamed Ali, A., and Ali Adam, B. 2020. Loss  
670 of Taste and Smell are Common Clinical Characteristics of Patients with COVID-19 in Somalia: A  
671 Retrospective Double Centre Study. *Infect Drug Resist*. 13:2631–2635.



- 672 Favas, T.T., Dev, P., Chaurasia, R.N., Chakravarty, K., Mishra, R., Joshi, D., Mishra, V.N., Kumar,  
673 A., Singh, V.K., Pandey, M., et al. 2020. Neurological manifestations of COVID-19: a systematic  
674 review and meta-analysis of proportions. *Neurol Sci.* 1–34.
- 675 Fisher, D.L., Pavel, A., and Malnick, S. 2021. Rapid recovery of taste and smell in a patient with  
676 SARS-CoV-2 following convalescent plasma therapy. *QJM Int J Med.*
- 677 Fistera, D., Pabst, D., Hartl, A., Schaarschmidt, B.M., Umutlu, L., Dolff, S., Holzner, C., Kill, C., and  
678 Risse, J. 2020. Separating the wheat from the chaff-COVID-19 in a German emergency  
679 department: a case-control study. *Int J Emerg Med.* 13:44.
- 680 Fleischer, M., Kohrmann, M., Dolff, S., Szepanowski, F., Schmidt, K., Herbstreit, F., Gungor, C.,  
681 Stolte, B., Steiner, K.M., Stadtler, C., et al. 2021. Observational cohort study of neurological  
682 involvement among patients with SARS-CoV-2 infection. *Ther Adv Neurol Disord.*  
683 14:1756286421993701.
- 684 Fontanet, A., Tondeur, L., Madec, Y., Grant, R., Besombes, C., Jolly, N., Fernandes Pellerin, S.,  
685 Ungeheuer, M.-N., Cailleau, I., Kuhmel, L., et al. 2020. Cluster of COVID-19 in northern France: A  
686 retrospective closed cohort study. *MedRxiv.* 2020.04.18.20071134.
- 687 Freni, F., Meduri, A., Gazia, F., Nicastro, V., Galletti, C., Aragona, P., Galletti, C., Galletti, B., and  
688 Galletti, F. 2020. Symptomatology in head and neck district in coronavirus disease (COVID-19):  
689 A possible neuroinvasive action of SARS-CoV-2. *Am J Otolaryngol.* 41:102612.
- 690 Galluzzi, F., Rossi, V., Bosetti, C., and Garavello, W. 2021. Risk Factors for Olfactory and  
691 Gustatory Dysfunctions in Patients with SARS-CoV-2 Infection. *Neuroepidemiology.* 1–8.
- 692 Gamper, E.-M., Giesinger, J.M., Oberguggenberger, A., Kemmler, G., Wintner, L.M., Gattringer,  
693 K., Sperner-Unterweger, B., Holzner, B., and Zabernigg, A. 2012. Taste alterations in breast and  
694 gynaecological cancer patients receiving chemotherapy: Prevalence, course of severity, and  
695 quality of life correlates. *Acta Oncol.* 51:490–496.
- 696 Garg, R., Jain, R., Sodani, A., Chouksey, D., Dosi, R., Athale, S., Goyal, N., Rathi, P., Singh, H., and  
697 Telang, K. 2020. Neurological Symptoms as Initial Manifestation of Covid-19 - An Observational  
698 Study. *Ann Indian Acad Neurol.* 23:482–486.
- 699 Gelardi, M., Trecca, E., Cassano, M., and Ciprandi, G. 2020. Smell and taste dysfunction during  
700 the COVID-19 outbreak: a preliminary report. *Acta Biomed.* 91:230–231.
- 701 Gerkin, R.C., Ohla, K., Veldhuizen, M.G., Joseph, P.V., Kelly, C.E., Bakke, A.J., Steele, K.E.,  
702 Farruggia, M.C., Pellegrino, R., Pepino, M.Y., et al. 2021. Recent Smell Loss Is the Best Predictor  
703 of COVID-19 Among Individuals With Recent Respiratory Symptoms. *Chem Senses.* 46.
- 704 Giacomelli, A., Pezzati, L., Conti, F., Bernacchia, D., Siano, M., Oreni, L., Rusconi, S., Gervasoni,  
705 C., Ridolfo, A.L., Rizzardini, G., et al. 2020. Self-reported olfactory and taste disorders in SARS-  
706 CoV-2 patients: a cross-sectional study. *Clin Infect Dis.*

- 707 Gianola, S., Barger, S., Campanini, I., Corbetta, D., Gambazza, S., Innocenti, T., Meroni, R.,  
708 Castellini, G., and Turolla, A. 2021. The Spread of COVID-19 Among 15,000 Physical Therapists in  
709 Italy: A Cross-Sectional Study. *Phys Ther.*
- 710 Gibbons, C., Hussain, M., O’Keeffe, D.T., and Simpkin, A.J. 2021. An analysis of patient self-  
711 reported COVID-19 symptoms during the first wave of the pandemic in Ireland. *Ir J Med Sci.*
- 712 Gonzalez, C., Garcia-Huidobro, F.G., Lagos, A.E., Aliaga, R., Fuentes-Lopez, E., Diaz, L.A., Garcia-  
713 Salum, T., Salinas, E., Toro, A., Callejas, C.A., et al. 2021. Prospective assessment of smell and  
714 taste impairment in a South-American coronavirus disease 2019 (COVID-19) cohort: Association  
715 with the need for hospitalization and reversibility of dysfunction. *Int Forum Allergy Rhinol.*
- 716 Gorzkowski, V., Bevilacqua, S., Charmillon, A., Jankowski, R., Gallet, P., Rumeau, C., and Nguyen,  
717 D.T. 2020. Evolution of olfactory disorders in COVID-19 patients. *Laryngoscope.*
- 718 Gozen, E.D., Aliyeva, C., Tevetoglu, F., Karaali, R., Balkan, Yener, H.M., and Ozdogan, H.A. 2020.  
719 Evaluation of Olfactory Function With Objective Tests in COVID-19-Positive Patients: A Cross-  
720 Sectional Study. *Ear Nose Throat J.* 145561320975510.
- 721 Gudbjartsson, D.F., Helgason, A., Jonsson, H., Magnusson, O.T., Melsted, P., Norddahl, G.L.,  
722 Saemundsdottir, J., Sigurdsson, A., Sulem, P., Agustsdottir, A.B., et al. 2020. Spread of SARS-  
723 CoV-2 in the Icelandic Population. *N Engl J Med.* 382:2302–2315.
- 724 Guillen Martinez, A., Andreu Galvez, M., Rodriguez Sanz, S., Hernandez Ruiz, P., Garcia Morillas,  
725 A., and Esteban Sanchez, T. 2020. Incidence of smell and taste disorders and associated factors  
726 in patients with mild to moderate COVID-19. *Otolaryngol Pol.* 75:1–5.
- 727 Güner, H.R., Hasanoğlu, İ., Kayaaslan, B., Aypak, A., Kaya Kalem, A., Eser, F., Özdemir, B.,  
728 Sarıcaoğlu, E.M., Ayhan, M., Aybar Bilir, Y., et al. 2020. COVID-19 experience of the major  
729 pandemic response center in the capital: Results of the pandemic’s first month in Turkey. *Turk J*  
730 *Med Sci.*
- 731 Gupta, S., Kumbhat, P., and Seervi, M. 2021a. Olfactory and Gustatory Dysfunction in Covid-19:  
732 An Observational Study in a Tertiary Care Institute of Western Rajasthan. *Indian J Otolaryngol*  
733 *Head Neck Surg.* 1–5.
- 734 Gupta, V., Banavara Rajanna, L., Upadhyay, K., Bhatia, R., Madhav Reddy, N., Malik, D., and  
735 Srivastava, A. 2021b. Olfactory and Gustatory Dysfunction in COVID-19 Patients from Northern  
736 India: A Cross-Sectional Observational Study. *Indian J Otolaryngol Head Neck Surg.* 1–8.
- 737 Gurrola, J.G., Chang, J.L., Roland, L.T., Loftus, P.A., and Cheung, S.W. 2021. Short-term  
738 chemosensory distortions and phantoms in COVID-19. *Laryngoscope Invest Otolaryngol.*
- 739 Haehner, A., Draf, J., Draeger, S., With, K. de, and Hummel, T. 2020. Predictive value of sudden  
740 olfactory loss in the diagnosis of COVID-19. *MedRxiv.* 2020.04.27.20081356.

- 741 Hajikhani, B., Calcagno, T., Nasiri, M.J., Jamshidi, P., Dadashi, M., Goudarzi, M., Eshraghi, A.A.,  
742 and Mirsaeidi, M. 2020. Olfactory and gustatory dysfunction in COVID-19 patients: A  
743 meta-analysis study. *Physiol Rep.* 8:e14578.
- 744 Hannum, M., E., Ramirez, V., A., Lipson, S., J., Herriman, R., D., Toskala, A., K., Lin, C., Joseph, P.,  
745 V., and Reed, D., R. 2020. Objective Sensory Testing Methods Reveal a Higher Prevalence of  
746 Olfactory Loss in COVID-19-Positive Patients Compared to Subjective Methods: A Systematic  
747 Review and Meta-Analysis. *Chem Senses.* 45:856–874.
- 748 Higgins, J.P.T., and Thompson, S.G. 2002. Quantifying heterogeneity in a meta-analysis. *Stat*  
749 *Med.* 21:1539–1558.
- 750 Hijazi, L.O., Alaraifi, A.K., and Alsaab, F. 2021. Otolaryngology manifestations of COVID-19 in  
751 pediatric patients. *Int J Pediatr Otorhinolaryngol.* 144:110701.
- 752 Hintschich, C.A., Wenzel, J.J., Hummel, T., Hankir, M.K., Kühnel, T., Vielsmeier, V., and Bohr, C.  
753 2020a. Psychophysical tests reveal impaired olfaction but preserved gustation in COVID-19  
754 patients. *Int Forum Allergy Rhinol.* 10:1105–1107.
- 755 Hintschich, C.A., Wenzel, J.J., Hummel, T., Hankir, M.K., Kühnel, T., Vielsmeier, V., and Bohr, C.  
756 2020b. Psychophysical tests reveal impaired olfaction but preserved gustation in COVID-19  
757 patients. *Int Forum Allergy Rhinol.*
- 758 Horvath, L., Lim, J.W.J., Taylor, J.W., Saief, T., Stuart, R., Rimmer, J., and Michael, P. 2020. Smell  
759 and taste loss in COVID-19 patients: assessment outcomes in a Victorian population. *Acta*  
760 *Otolaryngol.* 1–5.
- 761 Hoy, D., Brooks, P., Woolf, A., Blyth, F., March, L., Bain, C., Baker, P., Smith, E., and Buchbinder,  
762 R. 2012. Assessing risk of bias in prevalence studies: modification of an existing tool and  
763 evidence of interrater agreement. *J Clin Epidemiol.* 65:934–939.
- 764 Huang, N., Pérez, P., Kato, T., Mikami, Y., Okuda, K., Gilmore, R.C., Conde, C.D., Gasmi, B., Stein,  
765 S., Beach, M., et al. 2021. SARS-CoV-2 infection of the oral cavity and saliva. *Nat Med.* 27:892–  
766 903.
- 767 Huart, C., Philpott, C., Konstantinidis, I., Altundag, A., Whitcroft, K.L., Trecca, E.M.C., Cassano,  
768 M., Rombaux, Ph., and Hummel, T. 2020. Comparison of COVID-19 and common cold  
769 chemosensory dysfunction. *Rhinology.* 58:623–625.
- 770 İşlek, A., and Balcı, M.K. 2021. Diagnostic Value of Butanol Threshold Test in COVID-19 Related  
771 Olfactory Dysfunction. *Indian J Otolaryngol Head Neck Surg.*
- 772 Ismail, F., Farag, A., Haq, S., and Kamal, M.A. 2021. Clinical Characteristics of the First 100  
773 Patients of COVID-19 in Tobruk, Libya: A Brief Report From Low-Resource Settings. *Disaster*  
774 *Med Public Health Prep.* 1–4.

- 775 Iversen, K., Bundgaard, H., Hasselbalch, R.B., Kristensen, J.H., Nielsen, P.B., Pries-Heje, M.,  
776 Knudsen, A.D., Christensen, C.E., Fogh, K., Norsk, J.B., et al. 2020. Risk of COVID-19 in health-  
777 care workers in Denmark: an observational cohort study. *Lancet Infect Dis*.
- 778 Izquierdo-Domínguez, A., Rojas-Lechuga, M.J., Chiesa-Estomba, C., Calvo-Henríquez, C.,  
779 Ninchritz-Becerra, E., Soriano-Reixach, M., Poletti-Serafini, D., Villarreal, I.M., Maza-Solano,  
780 J.M., Moreno-Luna, R., et al. 2020. Smell and taste dysfunctions in COVID-19 are associated  
781 with younger age in ambulatory settings - a multicenter cross-sectional study. *J Investig Allergol*  
782 *Clin Immunol. O*.
- 783 Jain, A., Kumar, L., Kaur, J., Baisla, T., Goyal, P., Pandey, A.K., Das, A., and Parashar, L. 2020.  
784 Olfactory and taste dysfunction in coronavirus disease 2019 patients: its prevalence and  
785 outcomes. *J Laryngol Otol.* 1–10.
- 786 Jalessi, M., Barati, M., Rohani, M., Amini, E., Ourang, A., Azad, Z., Hosseinzadeh, F., Cavallieri, F.,  
787 Ghadirpour, R., Valzania, F., et al. 2020. Frequency and outcome of olfactory impairment and  
788 sinonasal involvement in hospitalized patients with COVID-19. *Neurol Sci*.
- 789 Jang, S.S., Choi, J.S., Kim, J.H., Kim, N., and Ference, E.H. 2021. Discordance Between Subjective  
790 and Objective Measures of Smell and Taste in US Adults. *Otolaryngol--Head Neck Surg Off J Am*  
791 *Acad Otolaryngol-Head Neck Surg.* 1945998211018386.
- 792 Jethani, B., Gupta, M., Wadhvani, P., Thomas, R., Balakrishnan, T., Mathew, G., Mathur, M.,  
793 Rao, B.P., Shukla, D., Khullar, A., et al. 2021. Clinical Characteristics and Remedy Profiles of  
794 Patients with COVID-19: A Retrospective Cohort Study. *Homeopathy*.
- 795 Kacem, I., Gharbi, A., Harizi, C., Souissi, E., Safer, M., Nasri, A., Letaief, H., Akkari, M., Hechaichi,  
796 A., Mrabet, S., et al. 2020. Characteristics, onset, and evolution of neurological symptoms in  
797 patients with COVID-19. *Neurol Sci*.
- 798 Kamel, F.O., Magadmi, R.M., Alqutub, S.T., Badawi, M., Al-Sayes, F., Badawi, M., Madni, T.A.,  
799 Alhothali, A., Abozinadah, E.A., and Adam, S. 2021. Clinical and hematologic presentations of  
800 adults with COVID-19 patients in Jeddah: A case control study. *J Infect Public Health.* 14:709–  
801 716.
- 802 Kandakure, V.T., Valvi, H.R., Khokle, P., More, M.S., and Chouhan, R. 2021. Prevalence and  
803 Recovery from Newly Onset Anosmia and Ageusia in Covid 19 Patients at our Tertiary Care  
804 Centre. *Indian J Otolaryngol Head Neck Surg*.
- 805 Karaarslan, F., Demircioğlu Güneri, F., and Kardeş, S. 2021. Postdischarge rheumatic and  
806 musculoskeletal symptoms following hospitalization for COVID-19: prospective follow-up by  
807 phone interviews. *Rheumatol Int.* 41:1263–1271.
- 808 Karadas, O., Ozturk, B., and Sonkaya, A.R. 2020. A prospective clinical study of detailed  
809 neurological manifestations in patients with COVID-19. *Neurol Sci.* 41:1991–1995.

- 810 Karni, N., Klein, H., Asseo, K., Benjamini, Y., Israel, S., Nammary, M., Olshtain-Pops, K., Nir-Paz,  
811 R., Hershko, A., Muszkat, M., et al. 2021. Self-Rated Smell Ability Enables Highly Specific  
812 Predictors of COVID-19 Status: A Case-Control Study in Israel. *Open Forum Infect Dis.*  
813 8:ofaa589.
- 814 Kavaz, E., Tahir, E., Bilek, H.C., Kemal, O., Deveci, A., and Aksakal Tanyel, E. 2021. Clinical  
815 significance of smell and taste dysfunction and other related factors in COVID-19. *Eur Arch*  
816 *Otorhinolaryngol.*
- 817 Kempker, R.R., Kempker, J.A., Peters, M., Rebolledo, P.A., Carroll, K., Toomer, L., Wang, Y.F.W.,  
818 Ray, S.M., and Hunter, M. 2020. Loss of Smell and Taste Among Healthcare Personnel Screened  
819 for Coronavirus 2019. *Clin Infect Dis.*
- 820 Kim, G.U., Kim, M.J., Ra, S.H., Lee, J., Bae, S., Jung, J., and Kim, S.H. 2020. Clinical characteristics  
821 of asymptomatic and symptomatic patients with mild COVID-19. *Clin Microbiol Infect.* 26:948  
822 e1-948 e3.
- 823 Klopfenstein, T., Zahra, H., Kadiane-Oussou, N.J., Lepiller, Q., Royer, P.Y., Toko, L., Gendrin, V.,  
824 and Zayet, S. 2020. New loss of smell and taste: Uncommon symptoms in COVID-19 patients on  
825 Nord Franche-Comte cluster, France. *Int J Infect Dis.*
- 826 Konstantinidis, I., Delides, A., Tsakiropoulou, E., Maragoudakis, P., Sapounas, S., and Tsiodras, S.  
827 2020. Short-Term Follow-Up of Self-Isolated COVID-19 Patients with Smell and Taste  
828 Dysfunction in Greece: Two Phenotypes of Recovery. *ORL J Otorhinolaryngol Relat Spec.* 1–9.
- 829 Krishnasamy, N., Natarajan, M., Ramachandran, A., Vivian Thangaraj, J.W., Etherajan, T.,  
830 Rengarajan, J., Shanmugasundaram, M., Kandasamy, A., Ramamoorthy, R., Velusamy, A., et al.  
831 2020. Clinical Outcomes among Asymptomatic or Mildly Symptomatic COVID-19 Patients in an  
832 Isolation Facility in Chennai, India. *Am J Trop Med Hyg.*
- 833 Kronborg, T.M., Kimer, N., Junker, A.E., Werge, M.P., Gluud, L.L., and Ytting, H. 2020. Experience  
834 from a COVID-19 first-line referral clinic in Greater Copenhagen. *Dan Med J.* 67.
- 835 Kumar, A.A., Lee, S.W.Y., Lock, C., and Keong, N.C.H. 2021a. Geographical Variations in Host  
836 Predisposition to COVID-19 Related Anosmia, Ageusia, and Neurological Syndromes. *Front Med.*  
837 8.
- 838 Kumar, L., Kahlon, N., Jain, A., Kaur, J., Singh, M., and Pandey, A.K. 2021b. Loss of smell and  
839 taste in COVID-19 infection in adolescents. *Int J Pediatr Otorhinolaryngol.* 142:110626.
- 840 Kumar, N., Bhartiya, S., Desai, S., Mutha, A., Beldar, A., and Singh, T. 2020. Seroprevalence of  
841 Antibodies Against SARS-CoV-2 Among Health Care Workers in Mumbai, India. *Asia Pac J Public*  
842 *Health.* 1010539520977307.

- 843 La Torre, G., Massetti, A.P., Antonelli, G., Fimiani, C., Fantini, M., Marte, M., Faticoni, A., Previte,  
844 C.M., Turriziani, O., Pugliese, F., et al. 2020. Anosmia and Ageusia as Predictive Signs of COVID-  
845 19 in Healthcare Workers in Italy: A Prospective Case-Control Study. *J Clin Med*. 9.
- 846 Ladoire, S., Goussot, V., Redersdorff, E., Cueff, A., Ballot, E., Truntzer, C., Ayati, S., Bengrine-  
847 Lefevre, L., Bremaud, N., Coudert, B., et al. 2021. Seroprevalence of SARS-CoV-2 among the  
848 staff and patients of a French cancer centre after first lockdown: The canSEROcov study. *Eur J*  
849 *Cancer*. 148:359–370.
- 850 Lagi, F., Piccica, M., Graziani, L., Vellere, I., Botta, A., Tilli, M., Ottino, L., Borchi, B., Pozzi, M.,  
851 Bartalesi, F., et al. 2020. Early experience of an infectious and tropical diseases unit during the  
852 coronavirus disease (COVID-19) pandemic, Florence, Italy, February to March 2020. *Euro*  
853 *Surveill*. 25.
- 854 Lampl, B.M.J., Buczovsky, M., Martin, G., Schmied, H., Leitzmann, M., and Salzberger, B. 2021.  
855 Clinical and epidemiological data of COVID-19 from Regensburg, Germany: a retrospective  
856 analysis of 1084 consecutive cases. *Infection*.
- 857 Lan, F.Y., Filler, R., Mathew, S., Buley, J., Iliaki, E., Bruno-Murtha, L.A., Osgood, R., Christophi,  
858 C.A., Fernandez-Montero, A., and Kales, S.N. 2020. COVID-19 symptoms predictive of  
859 healthcare workers' SARS-CoV-2 PCR results. *PLoS One*. 15:e0235460.
- 860 Lapostolle, F., Schneider, E., Vianu, I., Dollet, G., Roche, B., Berdah, J., Michel, J., Goix, L.,  
861 Chanzy, E., Petrovic, T., et al. 2020. Clinical features of 1487 COVID-19 patients with outpatient  
862 management in the Greater Paris: the COVID-call study. *Intern Emerg Med*. 15:813–817.
- 863 Le Bon, S., Payen, L., Prunier, L., Steffens, Y., Horoi, M., Vaira, L.A., Hopkins, C., Lechien, J.R., and  
864 Saussez, S. 2021. Making scents of loss of taste in COVID-19: Is self-reported loss of taste due to  
865 olfactory dysfunction? A prospective study using psychophysical testing. *Int Forum Allergy*  
866 *Rhinol*. alr.22815.
- 867 Le Bon, S.-D., Pisarski, N., Verbeke, J., Prunier, L., Cavelier, G., Thill, M.-P., Rodriguez, A.,  
868 Dequanter, D., Lechien, J.R., Le Bon, O., et al. 2020. Psychophysical evaluation of chemosensory  
869 functions 5 weeks after olfactory loss due to COVID-19: a prospective cohort study on 72  
870 patients. *Eur Arch Otorhinolaryngol*.
- 871 Lechien, J.R., Cabaraux, P., Chiesa-Estomba, C.M., Khalife, M., Plzak, J., Hans, S., Martiny, D.,  
872 Calvo-Henriquez, C., Hopkins, C., and Saussez, S. 2020a. Objective olfactory testing in patients  
873 presenting with sudden onset olfactory dysfunction as the first manifestation of confirmed  
874 COVID-19 infection.
- 875 Lechien, J.R., Chiesa-Estomba, C.M., Beckers, E., Mustin, V., Ducarme, M., Journe, F., Marchant,  
876 A., Jouffe, L., Barillari, M.R., Cammaroto, G., et al. 2021. Prevalence and 6-month recovery of  
877 olfactory dysfunction: a multicentre study of 1363 COVID-19 patients. *J Intern Med*. n/a.

- 878 Lechien, J.R., Chiesa-Estomba, C.M., Hans, S., Barillari, M.R., Jouffe, L., and Saussez, S. 2020b.  
879 Loss of Smell and Taste in 2013 European Patients With Mild to Moderate COVID-19. *Ann Intern*  
880 *Med*.
- 881 Lechien, J.R., Chiesa-Estomba, C.M., Place, S., Van Laethem, Y., Cabaraux, P., Mat, Q., Huet, K.,  
882 Plzak, J., Horoi, M., Hans, S., et al. 2020c. Clinical and epidemiological characteristics of 1420  
883 European patients with mild-to-moderate coronavirus disease 2019. *J Intern Med*. 288:335–  
884 344.
- 885 Lechner, M., Liu, J., Counsell, N., Ta, N.H., Rocke, J., Anmolsingh, R., Eynon-Lewis, N., Paun, S.,  
886 Hopkins, C., Khwaja, S., et al. 2020. Course of symptoms for loss of sense of smell and taste over  
887 time in one thousand forty-one healthcare workers during the Covid-19 pandemic: Our  
888 experience. *Clin Otolaryngol*. n/a.
- 889 Lee, D.J., Daliyot, D., Wang, R., Lockwood, J., Das, P., Zimlichman, E., and Lee, J.M. 2021.  
890 Comparative Study of Chemosensory Dysfunction in COVID-19 in 2 Geographically Distinct  
891 Regions. *Ear Nose Throat J*. 1455613211000170.
- 892 Lee, D.J., Lockwood, J., Das, P., Wang, R., Grinspun, E., and Lee, J.M. 2020a. Self-reported  
893 anosmia and dysgeusia as key symptoms of coronavirus disease 2019. *CJEM*. 1–8.
- 894 Lee, J.-M., and Lee, S.J. 2020. Olfactory and Gustatory Dysfunction in a COVID-19 Patient with  
895 Ankylosing Spondylitis Treated with Etanercept: Case Report. *J Korean Med Sci*. 35.
- 896 Lee, Y., Min, P., Lee, S., and Kim, S.W. 2020b. Prevalence and Duration of Acute Loss of Smell or  
897 Taste in COVID-19 Patients. *J Korean Med Sci*. 35:e174.
- 898 Levinson, R., Elbaz, M., Ben-Ami, R., Shasha, D., Levinson, T., Choshen, G., Petrov, K., Gadoth, A.,  
899 and Paran, Y. 2020. Anosmia and dysgeusia in patients with mild SARS-CoV-2 infection.  
900 *MedRxiv*. 2020.04.11.20055483.
- 901 Liang, Y., Xu, J., Chu, M., Mai, J., Lai, N., Tang, W., Yang, T., Zhang, S., Guan, C., Zhong, F., et al.  
902 2020. Neurosensory dysfunction: A diagnostic marker of early COVID-19. *Int J Infect Dis*.  
903 98:347–352.
- 904 Liguori, C., Pierantozzi, M., Spanetta, M., Sarmati, L., Cesta, N., Iannetta, M., Ora, J., Mina, G.G.,  
905 Puxeddu, E., Balbi, O., et al. 2020. Subjective neurological symptoms frequently occur in  
906 patients with SARS-CoV2 infection. *Brain Behav Immun*. 88:11–16.
- 907 Lindahl, J.F., Hoffman, T., Esmailzadeh, M., Olsen, B., Winter, R., Amer, S., Molnar, C., Svalberg,  
908 A., and Lundkvist, A. 2020. High seroprevalence of SARS-CoV-2 in elderly care employees in  
909 Sweden. *Infect Ecol Epidemiol*. 10:1789036.
- 910 Liu, G., Zong, G., Doty, R.L., and Sun, Q. 2016. Prevalence and risk factors of taste and smell  
911 impairment in a nationwide representative sample of the US population: a cross-sectional  
912 study. *BMJ Open*. 6:e013246.

- 913 Lombardi, A., Consonni, D., Carugno, M., Bozzi, G., Mangioni, D., Muscatello, A., Castelli, V.,  
914 Palomba, E., Cantu, A.P., Ceriotti, F., et al. 2020. Characteristics of 1,573 healthcare workers  
915 who underwent nasopharyngeal swab for SARS-CoV-2 in Milano, Lombardy, Italy. *Clin Microbiol*  
916 *Infect.*
- 917 Lombardi, A., Mangioni, D., Consonni, D., Cariani, L., Bono, P., Cantu, A.P., Tiso, B., Carugno, M.,  
918 Muscatello, A., Lunghi, G., et al. 2021. Seroprevalence of anti-SARS-CoV-2 IgG among healthcare  
919 workers of a large university hospital in Milan, Lombardy, Italy: a cross-sectional study. *BMJ*  
920 *Open.* 11:e047216.
- 921 Luers, J.C., Rokohl, A.C., Loreck, N., Wawer Matos, P.A., Augustin, M., Dewald, F., Klein, F.,  
922 Lehmann, C., and Heindl, L.M. 2020. Olfactory and Gustatory Dysfunction in Coronavirus  
923 Disease 19 (COVID-19). *Clin Infect Dis.*
- 924 Lv, H., Zhang, W., Zhu, Z., Xiong, Q., Xiang, R., Wang, Y., Shi, W., Deng, Z., and Xu, Y. 2020.  
925 Prevalence and recovery time of olfactory and gustatory dysfunctions of hospitalized patients  
926 with COVID19 in Wuhan, China. *Int J Infect Dis.*
- 927 Maechler, F., Gertler, M., Hermes, J., Loon, W. van, Schwab, F., Piening, B., Rojansky, S.,  
928 Hommes, F., Kausch, F., Lindner, A.K., et al. 2020. Epidemiological and clinical characteristics of  
929 SARS-CoV-2 infections at a testing site in Berlin, Germany, March and April 2020 - A cross-  
930 sectional study. *Clin Microbiol Infect.*
- 931 Magnavita, N., Tripepi, G., and Di Prinzio, R.R. 2020. Symptoms in Health Care Workers during  
932 the COVID-19 Epidemic. A Cross-Sectional Survey. *Int J Env Res Public Health.* 17.
- 933 Mair, M., Singhavi, H., Pai, A., Singhavi, J., Gandhi, P., Conboy, P., Baker, A., and Das, S. 2021. A  
934 Meta-Analysis of 67 Studies with Presenting Symptoms and Laboratory Tests of COVID-19  
935 Patients. *The Laryngoscope.* 131:1254–1265.
- 936 Makaronidis, J., Firman, C., Magee, C.G., Mok, J., Balogun, N., Lechner, M., Carnemolla, A., and  
937 Batterham, R.L. 2021. Distorted chemosensory perception and female sex associate with  
938 persistent smell and/or taste loss in people with SARS-CoV-2 antibodies: a community based  
939 cohort study investigating clinical course and resolution of acute smell and/or taste loss in  
940 people with and without SARS-CoV-2 antibodies in London, UK. *BMC Infect Dis.* 21:221.
- 941 Makda, A., Kumar, S., Kumar, A., Kumar, V., and Rizwan, A. 2020. The Frequency of Neurological  
942 Symptoms in COVID-19 Patients at a Tertiary Care Hospital in Pakistan. *Cureus.* 12:e10360.
- 943 Mannan, A., Mehedi, H.M.H., Chy, N., Qayum, M.O., Akter, F., Rob, M.A., Biswas, P., Hossain, S.,  
944 and Ayub, M. 2021. A multi-centre, cross-sectional study on coronavirus disease 2019 in  
945 Bangladesh: clinical epidemiology and short-term outcomes in recovered individuals. *New*  
946 *Microbes New Infect.* 40:100838.



- 947 Mao, L., Wang, M., Chen, S., He, Q., Chang, J., Hong, C., Zhou, Y., Wang, D., Li, Y., Jin, H., et al.  
948 2020. Neurological Manifestations of Hospitalized Patients with COVID-19 in Wuhan, China: a  
949 retrospective case series study. *MedRxiv*. 2020.02.22.20026500.
- 950 Martin-Sanz, E., Riestra, J., Yebra, L., Larran, A., Mancino, F., Yanes-Diaz, J., Garrote, M.,  
951 Colmenero, M., Montiel, E., Molina, C., et al. 2020. Prospective study in 355 patients with  
952 suspected COVID-19 infection. Value of cough, subjective hyposmia, and hypogeusia.  
953 *Laryngoscope*.
- 954 Mazzatenta, A., Neri, G., D'Ardes, D., De Luca, C., Marinari, S., Porreca, E., Cipollone, F.,  
955 Vecchiet, J., Falcicchia, C., Panichi, V., et al. 2020. Smell and Taste in Severe CoViD-19: Self-  
956 Reported vs. Testing. *Front Med*. 7:753.
- 957 Meini, S., Suardi, L.R., Busoni, M., Roberts, A.T., and Fortini, A. 2020. Olfactory and gustatory  
958 dysfunctions in 100 patients hospitalized for COVID-19: sex differences and recovery time in  
959 real-life. *Eur Arch Otorhinolaryngol*.
- 960 Menni, C., Valdes, A.M., Freidin, M.B., Sudre, C.H., Nguyen, L.H., Drew, D.A., Ganesh, S.,  
961 Varsavsky, T., Cardoso, M.J., El-Sayed Moustafa, J.S., et al. 2020. Real-time tracking of self-  
962 reported symptoms to predict potential COVID-19. *Nat Med*.
- 963 Mercante, G., Ferreli, F., De Virgilio, A., Gaino, F., Di Bari, M., Colombo, G., Russo, E.,  
964 Costantino, A., Pirola, F., Cugini, G., et al. 2020. Prevalence of Taste and Smell Dysfunction in  
965 Coronavirus Disease 2019. *JAMA Otolaryngol Head Neck Surg*.
- 966 Merkely, B., Szabo, A.J., Kosztin, A., Berenyi, E., Sebestyén, A., Lengyel, C., Merkely, G., Karady,  
967 J., Varkonyi, I., Papp, C., et al. 2020. Novel coronavirus epidemic in the Hungarian population, a  
968 cross-sectional nationwide survey to support the exit policy in Hungary. *Geroscience*. 42:1063–  
969 1074.
- 970 Merza, M.A., Haleem Al Mezori, A.A., Mohammed, H.M., and Abdulah, D.M. 2020. COVID-19  
971 outbreak in Iraqi Kurdistan: The first report characterizing epidemiological, clinical, laboratory,  
972 and radiological findings of the disease. *Diabetes Metab Syndr*. 14:547–554.
- 973 Moein, S.T., Hashemian, S.M.R., Mansourafshar, B., Khorram-Tousi, A., Tabarsi, P., and Doty,  
974 R.L. 2020. Smell dysfunction: a biomarker for COVID-19. *Int Forum Allergy Rhinol*. n/a.
- 975 Moeller, A.L., Mills, E.H.A., Collatz Christensen, H., Gnesin, F., Blomberg, S., Zyllyftari, N., Jensen,  
976 B., Ringgren, K.B., Broccia, M.D., Boggild, H., et al. 2021. Symptom presentation of SARS-CoV-2-  
977 positive and negative patients: a nested case-control study among patients calling the  
978 emergency medical service and medical helpline. *BMJ Open*. 11:e044208.
- 979 Moher, D., Liberati, A., Tetzlaff, J., Altman, D.G., and Group, T.P. 2009. Preferred Reporting  
980 Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLOS Med*.  
981 6:e1000097.

- 982 Monti, G., Leggieri, C., Fominskiy, E., Scandroglio, A.M., Colombo, S., Tozzi, M., Moizo, E., Mucci,  
983 M., Crivellari, M., Pieri, M., et al. 2021. Two months quality of life of COVID-19 invasively  
984 ventilated survivors; an Italian single-center study. *Acta Anaesthesiol Scand*.
- 985 Morlock, R., Morlock, A., Downen, M., and Shah, S.N. 2021. COVID-19 prevalence and  
986 predictors in United States adults during peak stay-at-home orders. *PLoS One*. 16:e0245586.
- 987 Morshed, M.S., Mosabbir, A.A., Chowdhury, P., Ashadullah, S.M., and Hossain, M.S. 2020.  
988 Clinical manifestations of patients with Coronavirus Disease 2019 (COVID- 19) attending at  
989 hospitals in Bangladesh. *MedRxiv*.
- 990 Murat, S., Dogruoz Karatekin, B., Icagasioglu, A., Ulasoglu, C., İçten, S., and Incealtin, O. 2020.  
991 Clinical presentations of pain in patients with COVID-19 infection. *Ir J Med Sci* 1971 -.
- 992 Nakagawa, S., Lagisz, M., O’Dea, R.E., Rutkowska, J., Yang, Y., Noble, D.W., and Senior, A.M.  
993 2020. The Orchard Plot: Cultivating a Forest Plot for Use in Ecology, Evolution and Beyond.  
994 *Research Synthesis Methods*. 12:4–12.
- 995 Nakakubo, S., Suzuki, M., Kamada, K., Yamashita, Y., Nakamura, J., Horii, H., Sato, K.,  
996 Matsumoto, M., Abe, Y., Tsuji, K., et al. 2020. Proposal of COVID-19 Clinical Risk Score for the  
997 management of suspected COVID-19 cases: a case control study. *BMC Infect Dis*. 20:858.
- 998 Nakanishi, H., Suzuki, M., Maeda, H., Nakamura, Y., Ikegami, Y., Takenaka, Y., Mori, Y., Hasuo,  
999 T., and Hasegawa, C. 2020. Differential Diagnosis of COVID-19: Importance of Measuring Blood  
1000 Lymphocytes, Serum Electrolytes, and Olfactory and Taste Functions. *Tohoku J Exp Med*.  
1001 252:109–119.
- 1002 Niklassen, A.S., Draf, J., Huart, C., Hintschich, C., Bocksberger, S., Trecca, E.M.C., Klimek, L., Bon,  
1003 S.D.L., Altundag, A., and Hummel, T. 2021a. COVID-19: Recovery from Chemosensory  
1004 Dysfunction. A Multicentre study on Smell and Taste. *The Laryngoscope*. 131:1095–1100.
- 1005 Niklassen, A.S., Draf, J., Huart, C., Hintschich, C., Bocksberger, S., Trecca, E.M.C., Klimek, L., Le  
1006 Bon, S.D., Altundag, A., and Hummel, T. 2021b. COVID-19: Recovery from chemosensory  
1007 dysfunction. A multicentre study on smell and taste. *Laryngoscope*.
- 1008 Ninchritz-Becerra, E., Soriano-Reixach, M.M., Mayo-Yáñez, M., Calvo-Henríquez, C., Martínez-  
1009 Ruiz de Apodaca, P., Saga-Gutiérrez, C., Parente-Arias, P., Villareal, I.M., Viera-Artiles, J., Poletti-  
1010 Serafini, D., et al. 2021. Subjective evaluation of smell and taste dysfunction in patients with  
1011 mild COVID-19 in Spain. *Med Clínica Engl Ed*. 156:61–64.
- 1012 Nishanth, D., Jhuma, S., Nitesh, G., Ramesh chand, M., Charanjit, S., Gupta, D.K., and Sen, M.K.  
1013 2021. COVID-19 with and without anosmia/dysgeusia: A case-control study. *J Med Virol*. n/a.
- 1014 Noh, J.Y., Yoon, J.G., Seong, H., Choi, W.S., Sohn, J.W., Cheong, H.J., Kim, W.J., and Song, J.Y.  
1015 2020. Asymptomatic infection and atypical manifestations of COVID-19: Comparison of viral  
1016 shedding duration. *J Infect*.

- 1017 Nolden, A.A., Hwang, L.-D., Boltong, A., and Reed, D.R. 2019. Chemosensory Changes from  
1018 Cancer Treatment and Their Effects on Patients' Food Behavior: A Scoping Review. *Nutrients*.  
1019 11:2285.
- 1020 Noviello, D., Costantino, A., Muscatello, A., Bandera, A., Consonni, D., Vecchi, M., and Basilisco,  
1021 G. 2021. Functional gastrointestinal and somatoform symptoms five months after SARS-CoV-2  
1022 infection: A controlled cohort study. *Neurogastroenterol Motil.* e14187.
- 1023 Oda, Y., Shiraishi, S., Shimada, M., and Kurai, O. 2021. Clinical profiles and outcome of patients  
1024 with COVID-19 in a specialized hospital in Japan. *J Anesth.*
- 1025 O'Keefe, J.B., Tong, E.J., O'Keefe, G.D., and Tong, D.C. 2021. Description of symptom course in a  
1026 telemedicine monitoring clinic for acute symptomatic COVID-19: a retrospective cohort study.  
1027 *BMJ Open.* 11:e044154.
- 1028 Omezli, M.M., and Torul, D. 2021. Evaluation of the xerostomia, taste and smell impairments  
1029 after Covid-19. *Med Oral Patol Oral Cir Bucal.*
- 1030 O'Sullivan, G., Jacob, S., Barrett, P.M., and Gallagher, J. 2021. Covid-19 presentation among  
1031 symptomatic healthcare workers in Ireland. *Occup Med.*
- 1032 Otte, M.S., Eckel, H.N.C., Poluschkin, L., Klussmann, J.P., and Luers, J.C. 2020. Olfactory  
1033 dysfunction in patients after recovering from COVID-19. *Acta Otolaryngol.* 1–4.
- 1034 Ozcan, E., Yavuzer, S., Borku Uysal, B., Islamoglu, M.S., Ikitimur, H., Unal, O.F., Akpınar, Y.E.,  
1035 Seyhan, S., Koc, S., Yavuzer, H., et al. 2021. The Relationship Between Positivity for COVID-19  
1036 RT-PCR and Symptoms, Clinical Findings, and Mortality in Turkey. *Expert Rev Mol Diagn.*
- 1037 Ozcelik Korkmaz, M., Egilmez, O.K., Ozcelik, M.A., and Guven, M. 2020. Otolaryngological  
1038 manifestations of hospitalised patients with confirmed COVID-19 infection. *Eur Arch*  
1039 *Otorhinolaryngol.*
- 1040 Paderno, A., Schreiber, A., Grammatica, A., Raffetti, E., Tomasoni, M., Gualtieri, T., Taboni, S.,  
1041 Zorzi, S., Lombardi, D., Deganello, A., et al. 2020. Smell and taste alterations in Covid-19: a  
1042 cross-sectional analysis of different cohorts. *Int Forum Allergy Rhinol.*
- 1043 Pang, K.W., Chee, J., Subramaniam, S., and Ng, C.L. 2020. Frequency and Clinical Utility of  
1044 Olfactory Dysfunction in COVID-19: a Systematic Review and Meta-analysis. *Curr Allergy Asthma*  
1045 *Rep.* 20:76.
- 1046 Parcha, V., Booker, K.S., Kalra, R., Kuranz, S., Berra, L., Arora, G., and Arora, P. 2021. A  
1047 retrospective cohort study of 12,306 pediatric COVID-19 patients in the United States. *Sci Rep.*  
1048 11:10231.

- 1049 Patel, A., Charani, E., Ariyanayagam, D., Abdulaal, A., Denny, S.J., Mughal, N., and Moore, L.S.P.  
1050 2020. New-onset anosmia and ageusia in adult patients diagnosed with SARS-CoV-2 infection.  
1051 Clin Microbiol Infect.
- 1052 Perlman, A., Vodonos Zilberg, A., Bak, P., Dreyfuss, M., Leventer-Roberts, M., Vurembrand, Y.,  
1053 Jeffries, H.E., Fisher, E., Steuerman, Y., Namir, Y., et al. 2020. Digital and Remote Care among  
1054 71,619 Individuals Seeking COVID-19 Health Information and Services: A Retrospective Cohort  
1055 Study. J Med Internet Res.
- 1056 Perula de Torres, L.A., Gonzalez-Lama, J., Jimenez Garcia, C., Sanchez Montero, R., Rider  
1057 Garrido, F., Ortega Lopez, Y., Pajares Conde, D., Ramirez Baena, M., Parraga Martinez, I.,  
1058 Romero-Rodriguez, E., et al. 2021. Frequency and predictive validity of olfactory and taste  
1059 dysfunction in patients with SARS-CoV-2 infection. Med Clin Barc.
- 1060 Petersen, M.S., Kristiansen, M.F., Hanusson, K.D., Danielsen, M.E., Steig, B. á, Gaini, S., Strøm,  
1061 M., and Weihe, P. 2020. Long COVID in the Faroe Islands - a longitudinal study among non-  
1062 hospitalized patients. Clin Infect Dis.
- 1063 Petrocelli, M., Ruggiero, F., Baietti, A.M., Pandolfi, P., Salzano, G., Salzano, F.A., Lechien, J.R.,  
1064 Saussez, S., De Riu, G., and Vaira, L.A. 2020a. Remote psychophysical evaluation of olfactory  
1065 and gustatory functions in early-stage coronavirus disease 2019 patients: the Bologna  
1066 experience of 300 cases. J Laryngol Otol. 1–12.
- 1067 Petrocelli, M., Ruggiero, F., Baietti, A.M., Pandolfi, P., Salzano, G., Salzano, F.A., Lechien, J.R.,  
1068 Saussez, S., Riu, G.D., and Vaira, L.A. 2020b. Remote psychophysical evaluation of olfactory and  
1069 gustatory functions in early-stage coronavirus disease 2019 patients: the Bologna experience of  
1070 300 cases. J Laryngol Otol. 134:571–576.
- 1071 Pinna, P., Grewal, P., Hall, J.P., Tavarez, T., Dafer, R.M., Garg, R., Osteraas, N.D., Pellack, D.R.,  
1072 Asthana, A., Fegan, K., et al. 2020. Neurological manifestations and COVID-19: Experiences from  
1073 a tertiary care center at the Frontline. J Neurol Sci. 415:116969.
- 1074 Polat, B., Yilmaz, N.H., Altin, G., Atakcan, Z., and Mert, A. 2021. Olfactory and Gustatory  
1075 Dysfunctions in COVID-19 Patients: From a Different Perspective. J Craniofac Surg. Publish  
1076 Ahead of Print.
- 1077 Printza, A., Katotomichelakis, M., Metallidis, S., Panagopoulos, P., Sarafidou, A., Petrakis, V., and  
1078 Constantinidis, J. 2020. The clinical course of smell and taste loss in COVID-19 hospitalized  
1079 patients. Hippokratia. 24:66–71.
- 1080 Printza, A., Katotomichelakis, M., Valsamidis, K., Metallidis, S., Panagopoulos, P., Panopoulou,  
1081 M., Petrakis, V., and Constantinidis, J. 2021. Smell and Taste Loss Recovery Time in COVID-19  
1082 Patients and Disease Severity. J Clin Med. 10.
- 1083 Qiu, C., Cui, C., Hautefort, C., Haehner, A., Zhao, J., Yao, Q., Zeng, H., Nisenbaum, E.J., Liu, L.,  
1084 Zhao, Y., et al. 2020. Olfactory and Gustatory Dysfunction as an Early Identifier of COVID-19 in

- 1085 Adults and Children: An International Multicenter Study. *Otolaryngol Head Neck Surg.*  
1086 194599820934376.
- 1087 R Core, T. 2021. R: A language and environment for statistical computing.
- 1088 Rajkumar, I., Anand, K.H., Revathishree, K., Shoba, K., and Srinivasan, K. 2020. Contemporary  
1089 Analysis of Olfactory Dysfunction in Mild to Moderate Covid 19 Patients in A Tertiary Health  
1090 Care Centre. *Indian J Otolaryngol Head Neck Surg.* 1–5.
- 1091 Ramteke, S., Tikkas, R., Goel, M., Mandraha, S., and Shrivastava, J. 2020. Paediatric COVID-19:  
1092 Milder Presentation-A Silver Lining in Dark Cloud. *J Trop Pediatr.* 67.
- 1093 Rawal, S., Hoffman, H.J., Bainbridge, K.E., Huedo-Medina, T.B., and Duffy, V.B. 2016. Prevalence  
1094 and Risk Factors of Self-Reported Smell and Taste Alterations: Results from the 2011–2012 US  
1095 National Health and Nutrition Examination Survey (NHANES). *Chem Senses.* 41:69–76.
- 1096 Rocke, J., Hopkins, C., Philpott, C., and Kumar, N. 2020. Is loss of sense of smell a diagnostic  
1097 marker in COVID-19: A systematic review and meta-analysis. *Clin Otolaryngol.* 45:914–922.
- 1098 Rodebaugh, T.L., Frumkin, M.R., Reiersen, A.M., Lenze, E.J., Avidan, M.S., Miller, J.P., Piccirillo,  
1099 J.F., Zorumski, C.F., and Mattar, C. 2021. Acute Symptoms of Mild to Moderate COVID-19 Are  
1100 Highly Heterogeneous Across Individuals and Over Time. *Open Forum Infect Dis.* 8.
- 1101 Rojas-Lechuga, M.J., Izquierdo-Dominguez, A., Chiesa-Estomba, C., Calvo-Henriquez, C.,  
1102 Villarreal, I.M., Cuesta-Chasco, G., Bernal-Sprekelsen, M., Mullol, J., and Alobid, I. 2020.  
1103 Chemosensory dysfunction in COVID-19 out-patients. *Eur Arch Otorhinolaryngol.*
- 1104 Roland, L.T., Gurrola, J.G., Loftus, P.A., Cheung, S.W., and Chang, J.L. 2020. Smell and taste  
1105 symptom-based predictive model for COVID-19 diagnosis. *Int Forum Allergy Rhinol.*
- 1106 Romero-Sanchez, C.M., Diaz-Maroto, I., Fernandez-Diaz, E., Sanchez-Larsen, A., Layos-Romero,  
1107 A., Garcia-Garcia, J., Gonzalez, E., Redondo-Penas, I., Perona-Moratalla, A.B., Del Valle-Perez,  
1108 J.A., et al. 2020. Neurologic manifestations in hospitalized patients with COVID-19: The  
1109 ALBACOVID registry. *Neurology.* 95:e1060–e1070.
- 1110 Rousseau, M.C., Hully, M., Milh, M., Juzeau, D., Pollez, B., Peudener, S., Bahi Buisson, N.,  
1111 Gautheron, V., Chabrol, B., and Billette de Villemeur, T. 2021. Clinical characteristics of COVID-  
1112 19 infection in polyhandicapped persons in France. *Arch Pédiatrie.*
- 1113 RStudio Team. 2020. RStudio: Integrated Development for R. RStudio. Boston, MA: RStudio,  
1114 PBC.
- 1115 Rubel, K., Sharma, D., Campiti, V., Yedlicka, G., Burgin, S.J., Illing, E.A., Kroenke, K., and Ting, J.Y.  
1116 2020. COVID-19 Status Differentially Affects Olfaction: A Prospective Case-Control Study. *OTO*  
1117 *Open.* 4:2473974X20970176.

- 1118 Sahoo, P.R., Sahu, M., Surapaneni, P.S., Maiti, A., Vankamamidi, R., Panda, N., and Biswal, R.N.  
1119 2021. Evolution of olfactory and gustatory dysfunctions in COVID-19 patients in India. *Eur Arch*  
1120 *Otorhinolaryngol.*
- 1121 Sakaguchi, W., Kubota, N., Shimizu, T., Saruta, J., Fuchida, S., Kawata, A., Yamamoto, Y.,  
1122 Sugimoto, M., Yakeishi, M., and Tsukinoki, K. 2020. Existence of SARS-CoV-2 Entry Molecules in  
1123 the Oral Cavity. *Int J Mol Sci.* 21:6000.
- 1124 Sakalli, E., Temirbekov, D., Bayri, E., Alis, E.E., Erdurak, S.C., and Bayraktaroglu, M. 2020. Ear  
1125 nose throat-related symptoms with a focus on loss of smell and/or taste in COVID-19 patients.  
1126 *Am J Otolaryngol.* 41:102622.
- 1127 Salcan, İ., Karakeçili, F., Salcan, S., Ünver, E., Akyüz, S., Seçkin, E., and Cingi, C. 2021. Is taste and  
1128 smell impairment irreversible in COVID-19 patients? *Eur Arch Otorhinolaryngol.* 278:411–415.
- 1129 Salepci, E., Turk, B., Ozcan, S.N., Bektas, M.E., Aybal, A., Dokmetas, I., and Turgut, S. 2020.  
1130 Symptomatology of COVID-19 from the otorhinolaryngology perspective: a survey of 223 SARS-  
1131 CoV-2 RNA-positive patients. *Eur Arch Otorhinolaryngol.*
- 1132 Saniasiaya, J., Islam, M.A., and Abdullah, B. 2021. Prevalence and Characteristics of Taste  
1133 Disorders in Cases of COVID-19: A Meta-analysis of 29,349 Patients. *Otolaryngol Neck Surg.*  
1134 165:33–42.
- 1135 Santos, R.E.A., Silva, M.G. da, Monte Silva, M.C.B. do, Barbosa, D.A.M., Gomes, A.L. do V.,  
1136 Galindo, L.C.M., Silva Aragão, R. da, and Ferraz-Pereira, K.N. 2021. Onset and duration of  
1137 symptoms of loss of smell/taste in patients with COVID-19: A systematic review. *Am J*  
1138 *Otolaryngol.* 42:102889.
- 1139 Savtale, S., Hippargekar, P., Bhise, S., and Kothule, S. 2021. Prevalence of  
1140 Otorhinolaryngological Symptoms in Covid 19 Patients. *Indian J Otolaryngol Head Neck Surg.*
- 1141 Sayin, I., Yasar, K.K., and Yazici, Z.M. 2020a. Taste and Smell Impairment in COVID-19: An AAO-  
1142 HNS Anosmia Reporting Tool-Based Comparative Study. *Otolaryngol Head Neck Surg.*  
1143 194599820931820.
- 1144 Sayin, P., Altinay, M., Cinar, A.S., and Ozdemir, H.M. 2020b. Taste and Smell Impairment in  
1145 Critically Ill Patients With COVID-19: An Intensive Care Unit Study. *Ear Nose Throat J.*  
1146 145561320977464.
- 1147 Sbrana, M.F., Fornazier, M.A., Bruni-Cardoso, A., Avelino-Silva, V.I., Schechtman, D., Voegels,  
1148 R.L., Malnic, B., Glezer, I., and Rezende Pinna, F. de. 2021. Olfactory Dysfunction in Frontline  
1149 Health Care Professionals During COVID-19 Pandemic in Brazil. *Front Physiol.* 12:622987.
- 1150 Schirinzi, A., Cazzolla, A.P., Lovero, R., Lo Muzio, L., Testa, N.F., Ciavarella, D., Palmieri, G.,  
1151 Pozzessere, P., Procacci, V., Di Serio, F., et al. 2020. New Insights in Laboratory Testing for  
1152 COVID-19 Patients: Looking for the Role and Predictive Value of Human epididymis secretory

- 1153 protein 4 (HE4) and the Innate Immunity of the Oral Cavity and Respiratory Tract.  
1154 *Microorganisms*. 8.
- 1155 Schmithausen, R.M., Dohla, M., Schobetal, H., Diegmann, C., Schulte, B., Richter, E., Eis-  
1156 Hubinger, A.M., and Streeck, H. 2020. Characteristic Temporary Loss of Taste and Olfactory  
1157 Senses in SARS-CoV-2-positive-Individuals with Mild Symptoms. *Pathog Immun*. 5:117–120.
- 1158 Schwab, J., Jensen, C.D., and Fjaeldstad, A.W. 2021. Sustained Chemosensory Dysfunction  
1159 during the COVID-19 Pandemic. *ORL J Otorhinolaryngol Relat Spec*. 1–10.
- 1160 Schwarzer, G., Chemaitelly, H., Abu-Raddad, L.J., and Rücker, G. 2019. Seriously misleading  
1161 results using inverse of Freeman-Tukey double arcsine transformation in meta-analysis of single  
1162 proportions. *Res Synth Methods*. 10:476–483.
- 1163 Sehanobish, E., Barbi, M., Fong, V., Kravitz, M., Sanchez Tejera, D., Asad, M., Matsumura, C.,  
1164 Ferastroaru, D., O’Neill, M., Karagic, M., et al. 2021. COVID-19-Induced Anosmia and Ageusia  
1165 Are Associated With Younger Age and Lower Blood Eosinophil Counts. *Am J Rhinol Allergy*.
- 1166 Seo, M.Y., Seok, H., Hwang, S.J., Choi, H.K., Jeon, J.H., Sohn, J.W., Park, D.W., Lee, S.H., and  
1167 Choi, W.S. 2020. Trend of Olfactory and Gustatory Dysfunction in COVID-19 Patients in a  
1168 Quarantine Facility. *J Korean Med Sci*. 35:e375.
- 1169 Shoer, S., Karady, T., Keshet, A., Shilo, S., Rossman, H., Gavrieli, A., Meir, T., Lavon, A., Kolobkov,  
1170 D., Kalka, I., et al. 2020. Who should we test for COVID-19? A triage model built from national  
1171 symptom surveys. *MedRxiv*. 2020.05.18.20105569.
- 1172 Sierpinski, R., Pinkas, J., Jankowski, M., Zgliczynski, W.S., Wierzba, W., Gujski, M., and  
1173 Szumowski, L. 2020. Sex differences in the frequency of gastrointestinal symptoms and  
1174 olfactory or taste disorders in 1942 nonhospitalized patients with coronavirus disease 2019  
1175 (COVID-19). *Pol Arch Intern Med*. 130:501–505.
- 1176 Singer-Cornelius, T., Cornelius, J., Oberle, M., Metternich, F.U., and Brockmeier, S.J. 2021.  
1177 Objective gustatory and olfactory dysfunction in COVID-19 patients: a prospective cross-  
1178 sectional study. *Eur Arch Otorhinolaryngol*.
- 1179 Smith, D.S., Richey, E.A., and Brunetto, W.L. 2020. A Symptom-Based Rule for Diagnosis of  
1180 COVID-19. *SN Compr Clin Med*. 1–8.
- 1181 Soh, S.H.L., See, A., Teo, N.W.Y., Tan, H.K., Palaniappan, G., Lim, M.L.A., Kadir, H.B.A., and Toh,  
1182 S.T. 2021. Prevalence of olfactory and taste dysfunction in COVID-19 patients: a community  
1183 care facility study. *Eur Arch Otorhinolaryngol*.
- 1184 Somekh, I., Yakub Hanna, H., Heller, E., Bibi, H., and Somekh, E. 2020. Age-Dependent Sensory  
1185 Impairment in Covid-19 Infection and Its Correlation with Ace2 Expression. *Pediatr Infect J*.

- 1186 Song, J., Deng, Y., Wang, H., Wang, Z., Liao, B., Ma, J., He, C., Pan, L., Liu, Y., Alobid, I., et al.  
1187 2021a. Self-reported Taste and Smell Disorders in Patients with COVID-19: Distinct Features in  
1188 China. *Curr Med Sci.* 41:14–23.
- 1189 Song, S.W., Kim, D., Park, J.Y., and Lee, S. 2021b. Symptoms and Characteristics Which Require  
1190 Attention During COVID-19 Screening at a Port of Entry. *J Korean Med Sci.* 36:e14.
- 1191 Speth, M.M., Singer-Cornelius, T., Obere, M., Gengler, I., Brockmeier, S.J., and Sedaghat, A.R.  
1192 2020. Olfactory Dysfunction and Sinonasal Symptomatology in COVID-19: Prevalence, Severity,  
1193 Timing, and Associated Characteristics. *Otolaryngol Head Neck Surg.* 194599820929185.
- 1194 Spinato, G., Fabbris, C., Polesel, J., Cazzador, D., Borsetto, D., Hopkins, C., and Boscolo-Rizzo, P.  
1195 2020. Alterations in Smell or Taste in Mildly Symptomatic Outpatients With SARS-CoV-2  
1196 Infection. *JAMA.*
- 1197 Stavem, K., Ghanima, W., Olsen, M.K., Gilboe, H.M., and Einvik, G. 2020. Persistent symptoms  
1198 1.5-6 months after COVID-19 in non-hospitalised subjects: a population-based cohort study.  
1199 *Thorax.*
- 1200 Sun, J., He, W.-T., Wang, L., Lai, A., Ji, X., Zhai, X., Li, G., Suchard, M.A., Tian, J., Zhou, J., et al.  
1201 2020. COVID-19: Epidemiology, Evolution, and Cross-Disciplinary Perspectives. *Trends Mol Med.*  
1202 26:483–495.
- 1203 Tarifi, A., Al Shdaifat, A.A., Al-Shudifat, A.M., Azab, M., Ismail, J., Bashir, R., Amro, A., Altarifi, A.,  
1204 and Khader, Y. 2021. Clinical, SinoNasal, and Long-Term Smell and Taste Outcomes in Mildly  
1205 Symptomatic COVID-19 Patients. *Int J Clin Pr.* e14260.
- 1206 Taziki Balajelini, M.H., Rajabi, A., Mohammadi, M., Nikoo, H.R., Tabarraei, A., Mansouri, M., and  
1207 Hosseini, S.M. 2021. Virus load and incidence of olfactory, gustatory, respiratory,  
1208 gastrointestinal disorders in COVID-19 patients: A retrospective cohort study. *Clin Otolaryngol*  
1209 *Off J ENT-UK Off J Neth Soc Oto-Rhino-Laryngol Cervico-Facial Surg.*
- 1210 Teklu, S., Sultan, M., Azazh, A., Worku, A., Redae, B., Walelegn, M., Tefera, M., Argaw, R.,  
1211 Waganew, W., Yifru, S., et al. 2020. Clinical and Socio-demographic Profile of the First 33  
1212 COVID-19 Cases Treated at Dedicated Treatment Center in Ethiopia. *Ethiop J Health Sci.* 30:645–  
1213 652.
- 1214 Thakur, K., Sagayaraj, A., Prasad, K.C., and Gupta, A. 2021. Olfactory Dysfunction in COVID-19  
1215 Patients: Findings from a Tertiary Rural Centre. *Indian J Otolaryngol Head Neck Surg.* 1–7.
- 1216 Tham, A.C., Thein, T.-L., Lee, C.S., Tan, G.S.E., Manuis, C.M., Siow, J.K., Leo, Y.S., and Lim, M.Y.  
1217 2020. Olfactory taste disorder as a presenting symptom of COVID-19: a large single-center  
1218 Singapore study. *Eur Arch Otorhinolaryngol.*



- 1219 Tong, J.Y., Wong, A., Zhu, D., Fastenberg, J.H., and Tham, T. 2020. The Prevalence of Olfactory  
1220 and Gustatory Dysfunction in COVID-19 Patients: A Systematic Review and Meta-analysis.  
1221 *Otolaryngol Neck Surg.* 163:3–11.
- 1222 Trachootham, D., Thongyen, S., Lam-Ubol, A., Chotechuang, N., Pongpirul, W., and  
1223 Prasithsirikul, W. 2021. Simultaneously Complete but Not Partial Taste and Smell Losses were  
1224 Associated with SAR-CoV-2 Infection. *Int J Infect Dis.*
- 1225 Tsivgoulis, G., Fragkou, P.C., Delides, A., Karofylakis, E., Dimopoulou, D., Sfikakis, P.P., and  
1226 Tsiodras, S. 2020. Quantitative evaluation of olfactory dysfunction in hospitalized patients with  
1227 Coronavirus [2] (COVID-19). *J Neurol.* 267:2193–2195.
- 1228 Tudrej, B., Sebo, P., Lourdaux, J., Cuzin, C., Floquet, M., Haller, D.M., and Maisonneuve, H. 2020.  
1229 Self-Reported Loss of Smell and Taste in SARS-CoV-2 Patients: Primary Care Data to Guide  
1230 Future Early Detection Strategies. *J Gen Intern Med.*
- 1231 United States Preventive Services Taskforce. Recommended Topics for Vision and Hearing.
- 1232 Vacchiano, V., Riguzzi, P., Volpi, L., Tappata, M., Avoni, P., Rizzo, G., Guerra, L., Zaccaroni, S.,  
1233 Cortelli, P., Michelucci, R., et al. 2020. Early neurological manifestations of hospitalized COVID-  
1234 19 patients. *Neurol Sci.*
- 1235 Vaira, L.A., Deiana, G., Fois, A.G., Pirina, P., Madeddu, G., Vito, A.D., Babudieri, S., Petrocelli, M.,  
1236 Serra, A., Bussu, F., et al. 2020a. Objective evaluation of anosmia and ageusia in COVID-19  
1237 patients: Single-center experience on 72 cases. *Head Neck.* 42:1252–1258.
- 1238 Vaira, L.A., Hopkins, C., Petrocelli, M., Lechien, J.R., Chiesa-Estomba, C.M., Salzano, G.,  
1239 Cucurullo, M., Salzano, F.A., Saussez, S., Boscolo-Rizzo, P., et al. 2020b. Smell and taste recovery  
1240 in coronavirus disease 2019 patients: a 60-day objective and prospective study. *J Laryngol Otol.*  
1241 134:703–709.
- 1242 Vaira, L.A., Hopkins, C., Petrocelli, M., Lechien, J.R., Soma, D., Giovanditto, F., Rizzo, D., Salzano,  
1243 G., Piombino, P., Saussez, S., et al. 2020c. Do olfactory and gustatory psychophysical scores  
1244 have prognostic value in COVID-19 patients? A prospective study of 106 patients. *J Otolaryngol -*  
1245 *Head Neck Surg.* 49:56.
- 1246 Vaira, L.A., Hopkins, C., Salzano, G., Petrocelli, M., Melis, A., Cucurullo, M., Ferrari, M.,  
1247 Gagliardini, L., Pipolo, C., Deiana, G., et al. 2020d. Olfactory and gustatory function impairment  
1248 in COVID-19 patients: Italian objective multicenter-study. *Head Neck.* 42:1560–1569.
- 1249 Vaira, L.A., Lechien, J.R., Salzano, G., Salzano, F.A., Maglitto, F., Saussez, S., and De Riu, G.  
1250 2020e. Gustatory Dysfunction: A Highly Specific and Smell-Independent Symptom of COVID-19.  
1251 *Indian J Otolaryngol Head Neck Surg.*

- 1252 Vaira, L.A., Salzano, G., Petrocelli, M., Deiana, G., Salzano, F.A., and Riu, G.D. 2020f. Validation  
1253 of a self-administered olfactory and gustatory test for the remotely evaluation of COVID-19  
1254 patients in home quarantine. *Head Neck*. 42:1570–1576.
- 1255 Van Loon, N., Verbrugghe, M., Cartuyvels, R., and Ramaekers, D. 2020. Diagnosis of COVID-19  
1256 Based on Symptomatic Analysis of Hospital Healthcare Workers in Belgium: Observational Study  
1257 in a Large Belgian Tertiary Care Center during Early COVID-19 Outbreak. *J Occup Env Med*.
- 1258 Vena, A., Berruti, M., Adessi, A., Blumetti, P., Brignole, M., Colognato, R., Gaggioli, G., Giacobbe,  
1259 D.R., Bracci-Laudiero, L., Magnasco, L., et al. 2020. Prevalence of Antibodies to SARS-CoV-2 in  
1260 Italian Adults and Associated Risk Factors. *J Clin Med*. 9.
- 1261 Villarreal, I.M., Morato, M., Martinez-RuizCoello, M., Navarro, A., Garcia-Chilleron, R., Ruiz, A.,  
1262 Almeida, I.V. de, Mazon, L., and Plaza, G. 2020. Olfactory and taste disorders in healthcare  
1263 workers with COVID-19 infection. *Eur Arch Otorhinolaryngol*.
- 1264 Wagner, T., Shweta, F.N.U., Murugadoss, K., Awasthi, S., Venkatakrisnan, A.J., Bade, S.,  
1265 Puranik, A., Kang, M., Pickering, B.W., O’Horo, J.C., et al. 2020. Augmented Curation of Clinical  
1266 Notes from a Massive EHR System Reveals Symptoms of Impending COVID-19 Diagnosis.  
1267 *MedRxiv*. 2020.04.19.20067660.
- 1268 Waterfield, T., Watson, C., Moore, R., Ferris, K., Tonry, C., Watt, A., McGinn, C., Foster, S.,  
1269 Evans, J., Lyttle, M.D., et al. 2020. Seroprevalence of SARS-CoV-2 antibodies in children: a  
1270 prospective multicentre cohort study. *Arch Dis Child*.
- 1271 Wee, L.E., Chan, Y.F.Z., Teo, N.W.Y., Cherng, B.P.Z., Thien, S.Y., Wong, H.M., Wijaya, L., Toh, S.T.,  
1272 and Tan, T.T. 2020. The role of self-reported olfactory and gustatory dysfunction as a screening  
1273 criterion for suspected COVID-19. *Eur Arch Otorhinolaryngol*.
- 1274 Weiss, J.J., Attuquayefio, T.N., White, E.B., Li, F., Herz, R.S., White, T.L., Campbell, M., Geng, B.,  
1275 Datta, R., Wyllie, A.L., et al. 2020. Tracking Smell Loss to Identify Healthcare Workers with SARS-  
1276 CoV-2 Infection. *MedRxiv*. 2020.09.07.20188813.
- 1277 Weng, C.H., Saal, A., Butt, W.W.W., and Chan, P.A. 2020. Characteristics and clinical outcomes  
1278 of COVID-19 in Hispanic/Latino patients in a community setting: A retrospective cohort study. *J*  
1279 *Med Virol*.
- 1280 Wierdsma, N.J., Kruijenga, H.M., Konings, L.A., Krebbers, D., Jorissen, J.R., Joosten, M.I., Aken,  
1281 L.H. van, Tan, F.M., Bodegraven, A.A. van, Soeters, M.R., et al. 2021. Poor nutritional status, risk  
1282 of sarcopenia and nutrition related complaints are prevalent in COVID-19 patients during and  
1283 after hospital admission. *Clin Nutr ESPEN*. 43:369–376.
- 1284 Yadav, V., Bhagat, S., Sharma, D.K., Sibia, R.P.S., Pandav, R., and Sandhu, V.P. 2021. Olfactory  
1285 and Gustatory Dysfunction in COVID-19 Patients: A Tertiary Care Institute Experience in India.  
1286 *Indian J Otolaryngol Head Neck Surg*. 1–8.

- 1287 Yan, C.H., Faraji, F., Prajapati, D.P., Boone, C.E., and DeConde, A.S. 2020a. Association of  
1288 chemosensory dysfunction and Covid-19 in patients presenting with influenza-like symptoms.  
1289 Int Forum Allergy Rhinol. n/a.
- 1290 Yan, C.H., Faraji, F., Prajapati, D.P., Ostrander, B.T., and DeConde, A.S. 2020b. Self-reported  
1291 olfactory loss associates with outpatient clinical course in COVID-19. Int Forum Allergy Rhinol.
- 1292 Yassin, A., Nawaiseh, M., Shaban, A., Alsherbini, K., El-Salem, K., Soudah, O., and Abu-Rub, M.  
1293 2021. Neurological manifestations and complications of coronavirus disease 2019 (COVID-19): a  
1294 systematic review and meta-analysis. BMC Neurol. 21:138.
- 1295 Zayet, S., Kadiane-Oussou, N.J., Lepiller, Q., Zahra, H., Royer, P.Y., Toko, L., Gendrin, V., and  
1296 Klopfenstein, T. 2020a. Clinical features of COVID-19 and influenza: a comparative study on  
1297 Nord Franche-Comte cluster. Microbes Infect. 22:481–488.
- 1298 Zayet, S., Klopfenstein, T., Mercier, J., Kadiane-Oussou, N.J., Lan Cheong Wah, L., Royer, P.Y.,  
1299 Toko, L., and Gendrin, V. 2020b. Contribution of anosmia and dysgeusia for diagnostic of  
1300 COVID-19 in outpatients. Infection.
- 1301 Zejda, J.E., Brożek, G.M., Kowalska, M., Barański, K., Kaleta-Pilarska, A., Nowakowski, A., Xia, Y.,  
1302 and Buszman, P. 2021. Seroprevalence of Anti-SARS-CoV-2 Antibodies in a Random Sample of  
1303 Inhabitants of the Katowice Region, Poland. Int J Environ Res Public Health. 18.
- 1304 Zifko, U., Schmiedlechner, T., Saelens, J., Zifko, K., Wagner, M., Assadian, O., Grisold, W., and  
1305 Stingl, H. 2021. Covid-19: Involvement of the nervous system. Identifying neurological  
1306 predictors defining the course of the disease. J Neurol Sci. 425:117438.
- 1307 Zimmerman, R.K., Nowalk, M.P., Bear, T., Taber, R., Clarke, K.S., Sax, T.M., Eng, H., Clarke, L.G.,  
1308 and Balasubramani, G.K. 2020. Proposed clinical indicators for efficient screening and testing  
1309 for COVID-19 infection using Classification and Regression Trees (CART) analysis. Hum Vaccin  
1310 Immunother. 1–4.
- 1311 Zou, L., Yu, T., Zhang, Y., Dai, L., Zhang, Z., and Zhang, Z. 2021. Olfactory and Gustatory  
1312 Dysfunctions in Patients With COVID-19 in Wuhan, China.
- 1313  
1314  
1315  
1316  
1317  
1318

Initial screening through database search (Pubmed, Medline, Google Scholar)



Relevant articles meeting initial inclusion criteria (written in the English language, data available)  
N=376



Full text screened for prevalence rate of taste dysfunction  
N=241



Data extracted by two authors (one author confirmed)  
N=241



Articles included  
N=241

**Excluded, N=135**

Recruited patients with chemosensory dysfunction, N=24

Not tested for COVID-19, N=10

Did not report taste loss data, N=45

Papers with overlapping samples, N=12

Other, N=44

