Table 2. Sample distribution by country and population group, unweighted percentage (%) ar size, PHIA	nd

Population group	Eswatini (2016–2017)		Malawi (2015–2016)		Tanzania (2016–2017)		Zambia (2016)	
	%	n	%	n	%	n	%	n
Adults ages 15-59 years								
PLHIV-female †	20.1	1, 918	8.8	1,477	4.2	1192	8.8	1,688
PLHIV-male †	9.1	870	4.1	680	1.8	517	4.1	779
MSM			0.8	161	0.2	67	0.2	51
SW-female			1.0	200	2.7	803	0.4	76
SW-male			‡0.2	32	0.4	133	‡0.1	31
AGYW	19.7	2, 013	21.5	4,102	20.3	6,031	21.6	4,587
Totals	100.0	10,233	100.0	19,106	100.0	29,638	100.0	21,278
Children ages 0–17 years								
OVC	27.8	2, 573	19.1	4,471	20.3	7,388	22.0	6, 094
Total	100.0	9, 271	100.0	23,432	100.0	36,376	100.0	27,655

<sup>†</sup> Denominators for PLHIV percentages were 9,556 in Eswatini, 16,698 in Malawi, 28,347 in Tanzania and

<sup>19,113</sup> in Zambia, and excluded people who did not test for HIV or received indeterminate HIV test results.
--- Results had fewer than 25 adults identified during the survey and were suppressed.

<sup>‡</sup> Estimate based on 25–49 persons/observations and should be interpreted with caution.