


## Supplement material

**Supplement table 1:** Treatment regimes in TM subjects

treatment	number of subjects
Nebido 1000mg, every 8-12 weeks, i.m	5
testosterone cream 50mg/d	4
Nebido 1000mg + Moniq Gynial 75µg/d (desogestrel gel)	2
Testavan 46mg/d (testosterone gel)	1
testosterone cream 37.5mg/d + Moniq Gynial 75µg/d	1
testosterone cream 50mg/d + Moniq Gynial 75µg/d	1
Nebido 1000mg + decapeptyl depot (triptorelin), every 4-5 weeks	1

**Supplement table 2:** Absolute number of data points (of max. 15) at each measurement included in linear mixed effect model analyses for each region and neurotransmitter ratio

		GABA+/tCr		Glx/tCr	
		M1	M2	M1	M2
<b>hippocampus</b>	CW	15	12	15	15
	TM	13	13	13	14
<b>insula</b>	CW	12	13	14	15
	TM	12	12	14	15
<b>putamen</b>	CW	14	13	15	15
	TM	12	12	15	14
<b>pallidum</b>	CW	15	13	15	15
	TM	12	12	15	14
<b>thalamus</b>	CW	15	13	15	15
	TM	12	12	14	15



**Supplement figure 1:** LCModel fits of exemplary non-edited (EDIT OFF) and difference spectra (DIFF) of the hippocampus, insula and pallidum.