

**COVCOG 1: Factors predicting Cognitive Symptoms in Long COVID. A First Publication from the
COVID and Cognition Study.**

*Panyuan Guo**, *Alvaro Benito Ballesteros**, *Sabine P Yeung**, *Ruby Liu**, *Arka Saha**, *Lyn Curtis**,
*Muzaffer Kaser**, *Mark P Haggard**, *Lucy G Cheke**¹

Supplementary Materials

Supplementary Table 1. Distribution of profession and employment status before the pandemic in No COVID and COVID groups. Note that participants could choose multiple options.

	No COVID (n = 185)	COVID (n = 181)	Chi-square tests
Profession			
Not in paid work	66 (35.7%)	22 (12.2%)	$\chi^2(1) = 27.72, p < .001,$ $V = .275^*$
Healthcare setting	16 (8.6%)	40 (22.1%)	$\chi^2(1) = 12.77, p < .001,$ $V = .187^*$
Professional / Consultant / Academic	26 (14.1%)	29 (16%)	n.s.
Office Support / Administrative Work	17 (9.2%)	24 (13.3%)	n.s.
Education (Teaching / Childcare)	7 (3.8%)	19 (10.5%)	$\chi^2(1) = 6.25, p = .012,$ $V = .131$
Sales / Service	13 (7%)	8 (4.4%)	n.s.
Technician / Engineer / Mechanic	2 (1.1%)	7 (3.9%)	n.s.
Cleaner / Canteen Staff / Handyman	7 (3.8%)	1 (0.6%)	n.s.

Musician / Actor / Dancer and related performance work	3 (1.6%)	3 (1.7%)	n.s.
Construction / Agricultural / Forestry / Fishery worker	5 (2.7%)	1 (0.6%)	n.s.
Transport worker	4 (2.2%)	2 (1.1%)	n.s.
Craft / Art and related trade work	3 (1.6%)	1 (0.6%)	n.s.
Managerial	1 (0.5%)	1 (0.6%)	n.s.
Professional sports	-	1 (0.6%)	n.s.
Other	16 (14.1%)	25 (13.8%)	n.s.
Employment status before pandemic			
Full time	61 (33%)	103 (56.9%)	$\chi^2(1) = 21.19, p < .001,$ $V = .241^*$
Part time	39 (21.1%)	39 (21.5%)	n.s.
Part time in multiple jobs	6 (3.2%)	6 (3.3%)	n.s.
Other schedule (e.g., ad hoc / gig economy)	3 (1.6%)	7 (3.9%)	n.s.
Not working (unemployed / career break / retired)	42 (22.7%)	16 (8.8%)	$\chi^2(1) = 13.18, p < .001,$ $V = .190^*$
Unpaid carer (e.g., children / elders)	7 (3.8%)	7 (3.9%)	n.s.
Student (e.g., University)	50 (27%)	26 (14.4%)	$\chi^2(1) = 8.91, p = .003,$ $V = .156^*$

* denotes p values below Sidak-correct alpha (i.e. non-null)

Supplementary Table 2. Distribution of medical history in No COVID and COVID groups (top).

Descriptive statistics and analyses for health behaviours factors (1 = *Never* – 6 = *Several times daily*;

higher scores indicating higher frequency) in No COVID and COVID groups (bottom).

	No COVID (n = 185)	COVID (n = 181)	
Medical history: Frequency (%)			Chi-square tests
Asthma	35 (18.9%)	38 (21%)	n.s.
Depression	38 (20.5%)	34 (18.8%)	n.s.
Other mental health disorder	35 (18.9%)	28 (15.5%)	n.s.
Obesity	15 (8.1%)	23 (12.7%)	n.s.
High blood pressure	20 (10.8%)	19 (10.5%)	n.s.
History of migraines	7 (3.8%)	19 (10.5%)	$\chi^2(1) = 6.25, p = .012,$ $V = .131$
Inflammatory / Autoimmune	4 (2.2%)	18 (9.9%)	$\chi^2(1) = 9.81, p = .002,$ $V = .164^*$
Chronic fatigue syndrome / Myalgic encephalomyelitis (ME)	9 (4.9%)	8 (4.4%)	n.s.
Psychiatric/neurodevelopmental disorder	6 (3.2%)	7 (3.9%)	n.s.
Cardiovascular disease / Angina	4 (2.2%)	6 (3.3%)	n.s.
Diabetes (Type 2)	9 (4.9%)	2 (1.1%)	n.s.
Diabetes (Type 1)	4 (2.2%)	-	n.s.
Cancer	1 (0.5%)	2 (1.1%)	n.s.
A clotting disorder	1 (0.5%)	2 (1.1%)	n.s.

None of the above	79 (42.7%)	65 (35.9%)	n.s.
Health behaviours: Mean (SD)			Mann-Whitney U tests
Diet: Fruit and vegetables	4.66 (1.16)	5.04 (1.1)	COVID > No COVID: $U = 13525, p = .001^*$
Diet: Sugary food	3.69 (1.11)	3.4 (1.07)	No COVID > COVID: $U = 14168, p = .008^*$
Diet: Fatty food	3.26 (1.01)	3.06 (.86)	No COVID > COVID: $U = 14825, p = .041$
Physical activity	3.3 (1.24)	3.7 (1.33)	COVID > No COVID: $U = 13752, p = .002^*$
Alcohol	2.67 (1.23)	2.69 (1.02)	n.s.
Smoking	1.63 (1.54)	1.4 (1.19)	n.s.

* denotes p values below Sidak-correct alpha

Supplementary Table 3. Distribution of demographics in COVID subgroups: Recovered (R), Ongoing (Mild/Moderate (C+)) and Ongoing (Severe (C++)).

	Recovered (R) (n = 42)	Ongoing (Mild/Moderate) (C+) (n = 53)	Ongoing (Severe) (C++) (n = 66)	Chi-square tests
Weeks since infection				
3–10	15 (35.7%)	7 (13.2%)	4 (6.1%)	$\chi^2(6) = 19.64, p = .003,$ $V = .247$
11–20	1 (2.4%)	4 (7.5%)	5 (7.6%)	
21–30	12 (28.6%)	15 (28.3%)	17 (25.8%)	
31+	14 (33.3%)	27 (50.9%)	40 (60.6%)	
Sex				
Man	13 (31%)	16 (30.2%)	10 (15.2%)	n.s.
Woman	29 (69%)	36 (67.9%)	54 (81.8%)	
Other	-	1 (1.9%)	2 (3%)	
Age				
18-20	9 (21.4%)	2 (3.8%)	-	$\chi^2(10) = 53.41, p < .001,$ $V = .407$
21-30	14 (33.3%)	8 (15.1%)	2 (3%)	
31-40	10 (23.8%)	11 (20.8%)	14 (21.2%)	
41-50	3 (7.1%)	14 (26.4%)	16 (24.2%)	
51-60	4 (9.5%)	9 (17%)	26 (39.4%)	
61 or above	2 (4.8%)	9 (17%)	8 (12.1%)	
Education				
GCSE/equiv or below	7 (16.7%)	4 (7.5%)	2 (3%)	

A level/equiv	7 (16.7%)	4 (7.5%)	2 (3%)	
Attended college (no degree) / Technical training /	7 (16.7%)	13 (24.5%)	8 (12.1%)	
Assoc. degree				$\chi^2(10) = 20.03, p = .029,$
Bachelor's degree	8 (19%)	15 (28.3%)	26 (39.4%)	$V = .249$
Master's / Professional degree	11 (26.2%)	13 (24.5%)	24 (36.4%)	
Doctorate degree	2 (4.8%)	4 (7.5%)	4 (6.1%)	
Country				
UK	23 (54.8%)	34 (64.2%)	61 (92.4%)	$\chi^2(4) = 24.24, p < .001,$
North America	10 (23.8%)	12 (22.6%)	5 (7.6%)	$V = .274$
Other	9 (21.4%)	7 (13.2%)	-	
Ethnicity				
Northern European	29 (69%)	48 (90.6%)	62 (93.9%)	$\chi^2(2) = 14.68, p = .001,$ $V = .302$
Southern European / Latinx	6 (14.3%)	5 (9.4%)	5 (7.5%)	n.s.
African / Afro-Caribbean	4 (9.5%)	1 (1.9%)	2 (3%)	n.s.
Asian	4 (9.6%)	3 (5.7%)	-	n.s.
Other / Prefer not to say	3 (7.2%)	1 (1.9%)	1 (1.5%)	n.s.

Supplementary Table 4. Diagnoses during initial illness among those seeking medical attention (*N* = 109).

Diagnoses within initial illness period	Percentage of positive diagnosis (number)
Hypoxia	14.7% (16)
TIA (Transient ischemic attack)	0.9% (1)
Clotting / Blood clots	5.5% (6)
Encephalopathies	0.9% (1)
Aneurism	1.8% (2)
Inflammatory Syndrome	4.6% (5)
Encephalitis	1.8% (2)
Hematoma	0.9% (1)
Kidney failure	0.9% (1)
Encephalomyelitis	-
Respiratory arrest	-
Delirium	1.8% (2)
Cardiac arrest	1.8% (2)
Stroke	2.8% (3)
Shock	1.8% (2)

Supplementary Table 5. Descriptive statistics and analyses for initial symptom experience (recoded such that higher numbers indicate increased severity: 1 = *Not at all*, 3 = *Very severe*) in

COVID subgroups: Recovered (*R*), Ongoing (Mild/Moderate (*C+*)) and Ongoing (Severe (*C++*)).

	Recovered (<i>R</i>) (<i>n</i> = 42)	Ongoing (Mild/Moderate) (<i>C+</i>) (<i>n</i> = 53)	Ongoing (Severe) (<i>C++</i>) (<i>n</i> = 66)	Kruskal-Wallis H tests
Initial symptoms: Mean (SD)				
Fatigue	2.24 (.69)	2.58 (.57)	2.76 (.47)	$H(2) = 18.45, p < .001^*$
Muscle/body pains	1.81 (.67)	2.17 (.58)	2.44 (.61)	$H(2) = 22.63, p < .001^*$
Breathing issues	1.62 (.73)	2.04 (.73)	2.44 (.7)	$H(2) = 28.04, p < .001^*$
Headache	2 (.73)	2.04 (.68)	2.35 (.71)	$H(2) = 8.51, p = .014$
Brain fog	1.62 (.76)	2.17 (.7)	2.26 (.79)	$H(2) = 17.67, p < .001^*$
Chest pain/tightness	1.55 (.63)	2.13 (.65)	2.26 (.73)	$H(2) = 25.22, p < .001^*$
Limb weakness	1.43 (.59)	1.75 (.62)	2.17 (.65)	$H(2) = 31.14, p < .001^*$
Dizziness	1.79 (.72)	1.83 (.7)	2.15 (.69)	$H(2) = 9.01, p = .011$
Drowsiness	1.88 (.74)	2.19 (.71)	2.14 (.72)	n.s.
Cough	2.02 (.75)	1.77 (.75)	2.08 (.83)	n.s.
Loss of appetite	1.74 (.70)	1.7 (.75)	2.03 (.78)	$H(2) = 6.51, p = .039$
Fever	1.71 (.64)	1.7 (.77)	1.98 (.79)	n.s.
Abdominal pain	1.36 (.53)	1.58 (.66)	1.91 (.78)	$H(2) = 14.84, p = .001^*$
Altered consciousness	1.43 (.63)	1.66 (.62)	1.91 (.72)	$H(2) = 13.09, p = .001^*$
Loss of smell/taste	1.79 (.87)	1.74 (.88)	1.88 (.85)	n.s.
Sore throat	1.67 (.69)	1.75 (.65)	1.85 (.69)	n.s.

Nausea	1.38 (.58)	1.51 (.61)	1.85 (.69)	$H(2) = 15.03, p = .001^*$
Confusion	1.33 (.61)	1.58 (.69)	1.83 (.74)	$H(2) = 13.94, p = .001^*$
Eye-soreness	1.48 (.67)	1.6 (.63)	1.8 (.73)	$H(2) = 6.25, p = .044$
Diarrhea	1.5 (.67)	1.55 (.61)	1.76 (.77)	n.s.
Anxiety	1.62 (.7)	1.72 (.63)	1.68 (.71)	n.s.
Blurred vision	1.19 (.46)	1.32 (.55)	1.62 (.67)	$H(2) = 15.45, p < .001^*$
Disorientation	1.17 (.38)	1.47 (.58)	1.48 (.66)	$H(2) = 8.78, p = .012$
Acid reflux	1.21 (.47)	1.32 (.55)	1.45 (.66)	n.s.
Visual disturbances	1.19 (.4)	1.19 (.44)	1.36 (.65)	n.s.
Rash	1.14 (.42)	1.36 (.59)	1.32 (.53)	n.s.
Speech difficulty	1.14 (.35)	1.26 (.45)	1.32 (.53)	n.s.
Numbness	1.05 (.22)	1.23 (.47)	1.3 (.53)	$H(2) = 8.51, p = .014$
Vomiting	1.17 (.44)	1.08 (.33)	1.23 (.46)	n.s.
Hallucinations	1.07 (.26)	1.08 (.27)	1.23 (.55)	n.s.
Foot sores	1.07 (.34)	1.11 (.32)	1.2 (.4)	n.s.
Itchy welts	1.14 (.42)	1.25 (.43)	1.17 (.41)	n.s.
Delirium	1.14 (.35)	1.11 (.42)	1.15 (.47)	n.s.

* denotes p values below Sidak-correct alpha (i.e. non-null)

Supplementary Table 6. Descriptive statistics and analyses for initial symptom experience (recoded such that higher numbers indicate increased severity: 1 = *Not at all*, 3 = *Very severe*) in Unknown, Unconfirmed COVID, and Confirmed COVID groups.

	Unknown (n = 55)	Unconfirmed (n = 96)	Confirmed (n = 65)	Kruskal-Wallis H tests
Initial symptoms: Mean (SD)				
Fatigue	1.84 (.79)	2.58 (.63)	2.54 (.56)	$H(2) = 36.94, p < .001^*$
Muscle/body pains	1.58 (.63)	2.17 (.71)	2.22 (.6)	$H(2) = 30.09, p < .001^*$
Loss of smell/taste	1.22 (.57)	1.63 (.77)	2.08 (.92)	$H(2) = 30.92, p < .001^*$
Drowsiness	1.55 (.66)	2.11 (.72)	2.05 (.74)	$H(2) = 22, p < .001^*$
Brain fog	1.42 (.63)	2.08 (.77)	2.03 (.83)	$H(2) = 26.67, p < .001^*$
Headache	1.96 (.61)	2.25 (.68)	2.02 (.76)	$H(2) = 7.63, p = .022$
Breathing issues	1.51 (.66)	2.22 (.76)	1.91 (.81)	$H(2) = 27.87, p < .001^*$
Dizziness	1.53 (.6)	2 (.7)	1.88 (.74)	$H(2) = 15.78, p < .001^*$
Chest pain/tightness	1.4 (.56)	2.16 (.73)	1.85 (.71)	$H(2) = 36.05, p < .001^*$
Fever	1.49 (.64)	1.8 (.73)	1.85 (.8)	$H(2) = 7.96, p = .019$
Loss of appetite	1.45 (.66)	1.86 (.73)	1.82 (.81)	$H(2) = 11.72, p = .003$
Limb weakness	1.29 (.5)	1.86 (.71)	1.8 (.67)	$H(2) = 27.62, p < .001^*$
Cough	2.05 (.65)	2.09 (.78)	1.77 (.77)	$H(2) = 7.77, p = .021$
Altered consciousness	1.35 (.62)	1.73 (.7)	1.66 (.67)	$H(2) = 13.15, p = .001^*$
Anxiety	1.84 (.76)	1.69 (.62)	1.66 (.76)	n.s.
Eye-soreness	1.38 (.56)	1.69 (.65)	1.6 (.75)	$H(2) = 7.89, p = .019$
Sore throat	2.02 (.62)	1.9 (.67)	1.58 (.64)	$H(2) = 14.61, p = .001^*$
Diarrhea	1.36 (.49)	1.65 (.71)	1.58 (.68)	n.s.

Abdominal pain	1.4 (.6)	1.72 (.72)	1.57 (.71)	$H(2) = 7.54, p = .023$
Nausea	1.38 (.53)	1.67 (.66)	1.54 (.66)	$H(2) = 6.78, p = .034$
Confusion	1.25 (.52)	1.69 (.74)	1.52 (.67)	$H(2) = 14.15, p = .001^*$
Acid reflux	1.22 (.46)	1.29 (.54)	1.43 (.64)	n.s.
Disorientation	1.18 (.43)	1.45 (.61)	1.32 (.53)	$H(2) = 8.37, p = .015$
Blurred vision	1.2 (.52)	1.5 (.62)	1.28 (.57)	$H(2) = 15.03, p = .001^*$
Rash	1.18 (.51)	1.3 (.53)	1.26 (.54)	n.s.
Visual disturbances	1.13 (.34)	1.29 (.54)	1.22 (.52)	n.s.
Numbness	1.07 (.26)	1.23 (.47)	1.18 (.43)	n.s.
Speech difficulty	1.04 (.19)	1.3 (.51)	1.18 (.39)	$H(2) = 13.76, p = .001^*$
Itchy welts	1.18 (.55)	1.21 (.43)	1.15 (.4)	n.s.
Foot sores	1.04 (.27)	1.15 (.38)	1.12 (.33)	n.s.
Vomiting	1.15 (.41)	1.19 (.44)	1.12 (.38)	n.s.

** denotes p values below Sidak-correct alpha (i.e. non-null)*

Supplementary Table 7. Descriptive statistics and analyses for ongoing symptom

experience (recoded such that higher numbers indicate increased severity: 1 = *Not at all*, 5 = *Very severe and often*) in Unknown, Unconfirmed COVID, and Confirmed COVID groups.

	Unknown (n = 31)	Unconfirmed (n = 88)	Confirmed (n = 58)	Kruskal-Wallis H tests
Ongoing symptoms: Mean (SD)				
Fatigue	2.55 (1.5)	3.77 (1.35)	3.83 (1.39)	$H(2) = 16.86, p < .001^*$
Difficulty concentrating	2.03 (1.05)	3.64 (1.22)	3.52 (1.3)	$H(2) = 32.58, p < .001^*$
Brain fog	1.97 (1.33)	3.43 (1.45)	3.24 (1.49)	$H(2) = 20.73, p < .001^*$
Forgetfulness	2.35 (1.31)	3.27 (1.32)	3.19 (1.34)	$H(2) = 10.92, p = .004$
Tip of the Tongue (ToT)	1.65 (0.92)	3 (1.41)	3.16 (1.41)	$H(2) = 25.71, p < .001^*$
Difficulty sleeping	2.35 (1.33)	3.15 (1.41)	2.86 (1.16)	$H(2) = 8.09, p = .017$
Muscle/body pains	1.71 (0.78)	2.83 (1.39)	2.83 (1.48)	$H(2) = 16.32, p < .001^*$
Breathing issues	2 (1.29)	2.98 (1.47)	2.83 (1.4)	$H(2) = 11.07, p = .004$
Fast/irregular pulse	1.39 (0.88)	2.85 (1.41)	2.81 (1.5)	$H(2) = 29.07, p < .001^*$
Semantic disfluency	1.35 (0.88)	2.64 (1.42)	2.78 (1.49)	$H(2) = 26.08, p < .001^*$
Night waking	2.23 (1.33)	3.24 (1.38)	2.74 (1.33)	$H(2) = 13.18, p = .001^*$
Chest pain/tightness	1.84 (1.21)	2.89 (1.44)	2.69 (1.26)	$H(2) = 14.61, p = .001^*$
Vivid dreams	2.03 (1.3)	2.78 (1.15)	2.62 (1.32)	$H(2) = 9.12, p = .01$
Dizziness	1.9 (1.17)	2.67 (1.32)	2.57 (1.42)	$H(2) = 8.33, p = .016$
Drowsiness	2.06 (1.39)	2.35 (1.25)	2.53 (1.33)	n.s.
Headache	2.32 (1.05)	2.78 (1.33)	2.53 (1.31)	n.s.
Limb weakness	1.42 (0.67)	2.38 (1.34)	2.47 (1.47)	$H(2) = 14.53, p < .001^*$
Depression	3.03 (1.33)	2.51 (1.3)	2.43 (1.24)	n.s.

Excess thirst	1.77 (1.12)	2.43 (1.35)	2.41 (1.36)	$H(2) = 6.69, p = .035$
Loss of smell/taste	1.16 (0.52)	1.84 (1.21)	2.38 (1.62)	$H(2) = 16.25, p < .001^*$
Head pressure	1.68 (0.95)	2.4 (1.38)	2.31 (1.35)	$H(2) = 6.53, p = .038$
Anxiety	2.65 (1.52)	2.45 (1.33)	2.24 (1.26)	n.s.
Confusion	1.39 (0.67)	2.18 (1.3)	2.22 (1.36)	$H(2) = 9.89, p = .007$
Weight gain	1.81 (1.25)	2.1 (1.33)	2.16 (1.35)	n.s.
Abdominal pain	1.71 (1.07)	2.22 (1.16)	2 (1.14)	$H(2) = 6.44, p = .04$
Pins & needles	1.68 (1.01)	2.19 (1.26)	1.98 (1.16)	n.s.
Sore throat	1.94 (1.03)	2.33 (1.22)	1.93 (1.23)	$H(2) = 6.38, p = .041$
Loss of appetite	1.48 (0.57)	1.83 (1.08)	1.9 (1.28)	n.s.
Altered consciousness	1.65 (1.17)	1.85 (1.17)	1.9 (1.2)	n.s.
Diarrhea	1.52 (0.81)	1.91 (1.02)	1.9 (1.14)	n.s.
Acid reflux	1.39 (0.8)	1.68 (1.09)	1.88 (1.26)	n.s.
Eye-soreness	1.65 (1.08)	2.26 (1.19)	1.88 (1.31)	$H(2) = 11.27, p = .004$
Blurred vision	1.39 (0.76)	2.03 (1.24)	1.84 (1.27)	$H(2) = 8.14, p = .017$
Nausea	1.48 (0.85)	1.89 (0.95)	1.79 (1.21)	$H(2) = 6.32, p = .042$
Cough	2.16 (1.39)	2.31 (1.16)	1.79 (0.95)	$H(2) = 7.12, p = .028$
Hot flushes	1.26 (0.58)	1.84 (1.12)	1.74 (1.21)	$H(2) = 7.88, p = .019$
Weight loss	1.48 (0.89)	1.52 (0.99)	1.62 (1.09)	n.s.
Disorientation	1.29 (0.82)	1.57 (0.84)	1.48 (0.98)	n.s.
Visual disturbances	1.23 (0.62)	1.52 (0.98)	1.47 (0.98)	n.s.
Speech difficulty	1.19 (0.65)	1.66 (0.98)	1.43 (0.9)	$H(2) = 9.45, p = .009$
Numbness	1.23 (0.62)	1.6 (1.06)	1.43 (0.94)	n.s.
Rash	1.39 (0.96)	1.55 (1.04)	1.4 (0.79)	n.s.
Fever	1.13 (0.43)	1.42 (0.85)	1.33 (0.85)	n.s.

Incontinence	1.16 (0.58)	1.27 (0.6)	1.28 (0.7)	n.s.
Itchy welts	1.32 (0.91)	1.24 (0.66)	1.22 (0.53)	n.s.
Foot sores	1.06 (0.25)	1.22 (0.65)	1.17 (0.57)	n.s.
Vomiting	1.13 (0.43)	1.19 (0.48)	1.17 (0.53)	n.s.

** denotes p values below Sidak-correct alpha (i.e. non-null)*

Supplementary Table 8. Factors and loadings from the “since then” cognitive symptoms PCA.

	Cognitive Symptoms
Brain Fog	0.889
Forgetfulness	0.881
Tip of the Tongue (ToT)	0.879
Semantic Disfluency	0.874
Difficulty Concentrating	0.859

Supplementary Table 9. Distribution of experience with long COVID in COVID subgroups: Recovered (R), Ongoing (Mild/Moderate (C+)) and Ongoing (Severe (C++)).

	Now Recovered (R) (n = 27**)	Ongoing (Mild/Moderate) (C+) (n = 53)	Ongoing (Severe) (C++) (n = 66)	Chi-square tests
Identify as suffering from “Long COVID”				
Yes	3 (11.1%)	43 (81.1%)	62 (93.9%)	$\chi^2(4) = 85.75, p < .001,$ $V = .542$
No	16 (59.3%)	2 (3.8%)	-	
Other	8 (29.6%)	8 (15.1%)	4 (6.1%)	
Change of symptoms after initial illness				
No ongoing symptoms after initial recovery	5 (18.5%)	-	-	$\chi^2(6) = 37.52, p < .001,$ $V = .367$
Different symptoms at different times	8 (29.6%)	28 (52.8%)	39 (59.1%)	
Improvement in symptoms over time	5 (18.5%)	18 (34%)	9 (13.6%)	
Symptoms have been very consistent	3 (11.1%)	7 (13.2%)	17 (25.8%)	
I don't know / N/A	6 (22.2%)	-	1 (1.5%)	
Cycle of symptoms after initial illness				
Cycle every few days	3 (11.1%)	11 (20.8%)	14 (21.2%)	n.s.
Cycle every few weeks	3 (11.1%)	13 (24.5%)	19 (28.8%)	
Cycle monthly	2 (7.4%)	7 (13.2%)	9 (13.6%)	

No cycling	12 (44.4%)	19 (35.8%)	23 (34.8%)	
N/A	7 (25.9%)	3 (5.7%)	1 (1.5%)	
Impacts of long COVID				
Long period unable to work (due to illness)	2 (7.4%)	15 (28.3%)	50 (75.8%)	$\chi^2(2) = 46.42, p < .001, V = .564^*$
Difficulty coping day-to-day activities	6 (22.2%)	28 (52.8%)	48 (72.7%)	$\chi^2(2) = 20.23, p < .001, V = .372^*$
Difficulty getting medical professionals to take symptoms seriously	1 (3.7%)	21 (39.6%)	38 (57.6%)	$\chi^2(2) = 23.05, p < .001, V = .397^*$
Lost job due to illness	1 (3.7%)	9 (17%)	32 (48.5%)	$\chi^2(2) = 24.39, p < .001, V = .409^*$
Feeling that you have experienced a trauma	4 (14.8%)	21 (39.6%)	31 (47%)	$\chi^2(2) = 8.44, p = .015, V = .240$
Financial difficulty (as a result of illness)	1 (3.7%)	7 (13.2%)	14 (21.2%)	n.s.
None of the above	18 (66.7%)	9 (17%)	1 (1.5%)	$\chi^2(2) = 52.73, p < .001, V = .601^*$

* denotes p values below Sidak-correct alpha at .007 for the impacts of long COVID

**Excluding a small portion of participants who reported asymptomatic or feeling completely better very quickly from the Recovered subgroup ($n = 15$)

Supplementary Table 10. Descriptive statistics and analyses for ongoing symptom

experience (recoded such that higher numbers indicate increased severity: 1 = *Not at all*, 5 = *Very severe and often*) in COVID subgroups: Recovered (*R*), Ongoing (Mild/Moderate (*C+*)) and Ongoing (Severe (*C++*)).

	Now Recovered (<i>R</i>) (<i>n</i> = 27**)	Ongoing (Mild/Moderate) (<i>C+</i>) (<i>n</i> = 53)	Ongoing (Severe) (<i>C++</i>) (<i>n</i> = 66)	Kruskal-Wallis H tests
Ongoing symptoms: Mean (SD)				
Fatigue	2.33 (1.24)	3.57 (1.12)	4.58 (0.98)	$H(2) = 57.34, p < .001^*$
Difficulty concentrating	2.33 (1.36)	3.26 (1)	4.36 (0.78)	$H(2) = 52.98, p < .001^*$
Brain fog	1.85 (1.26)	3.17 (1.21)	4.12 (1.18)	$H(2) = 46.08, p < .001^*$
Forgetfulness	2.07 (1.24)	2.98 (1.08)	3.92 (1.13)	$H(2) = 39.24, p < .001^*$
Tip of the Tongue (ToT)	1.78 (1.16)	2.81 (1.18)	3.79 (1.23)	$H(2) = 41.03, p < .001^*$
Breathing issues	1.63 (1.15)	2.57 (1.07)	3.73 (1.32)	$H(2) = 46.08, p < .001^*$
Fast/irregular pulse	1.59 (0.8)	2.45 (1.25)	3.65 (1.31)	$H(2) = 45.11, p < .001^*$
Night waking	2 (1.27)	2.87 (1.35)	3.61 (1.16)	$H(2) = 27.25, p < .001^*$
Muscle/body pains	1.67 (0.83)	2.51 (1.15)	3.56 (1.41)	$H(2) = 37.66, p < .001^*$
Chest pain/tightness	1.70 (1.03)	2.43 (1.14)	3.56 (1.24)	$H(2) = 42.21, p < .001^*$
Semantic disfluency	1.59 (0.84)	2.19 (1.24)	3.55 (1.3)	$H(2) = 45.81, p < .001^*$
Difficulty sleeping	2.67 (1.44)	2.81 (1.21)	3.36 (1.3)	$H(2) = 7.47, p = .024$
Headache	1.96 (1.19)	2.23 (1.03)	3.35 (1.3)	$H(2) = 30.33, p < .001^*$
Dizziness	1.81 (1.18)	2.19 (1.04)	3.32 (1.33)	$H(2) = 31.94, p < .001^*$
Limb weakness	1.22 (0.42)	2.13 (1.09)	3.12 (1.45)	$H(2) = 40.54, p < .001^*$
Head pressure	1.67 (0.92)	1.94 (0.97)	2.98 (1.52)	$H(2) = 22.01, p < .001^*$

Vivid dreams	2.07 (1.17)	2.79 (1.1)	2.92 (1.26)	$H(2) = 9.94, p = .007$
Excess thirst	1.74 (1.16)	2.23 (1.2)	2.86 (1.40)	$H(2) = 15.74, p < .001^*$
Sore throat	1.33 (0.56)	1.91 (0.9)	2.73 (1.4)	$H(2) = 26.45, p < .001^*$
Drowsiness	1.78 (1.01)	2.4 (1.28)	2.71 (1.3)	$H(2) = 11.11, p = .004$
Confusion	1.33 (0.83)	2.02 (1.01)	2.7 (1.49)	$H(2) = 21.82, p < .001^*$
Pins & needles	1.41 (0.64)	1.75 (0.88)	2.68 (1.38)	$H(2) = 24.69, p < .001^*$
Abdominal pain	1.44 (0.7)	1.81 (0.74)	2.67 (1.33)	$H(2) = 25.12, p < .001^*$
Eye-soreness	1.48 (0.8)	1.94 (1.08)	2.5 (1.4)	$H(2) = 13.46, p = .001^*$
Blurred vision	1.19 (0.56)	1.68 (0.98)	2.5 (1.41)	$H(2) = 28.35, p < .001^*$
Cough	1.59 (0.84)	1.92 (1.04)	2.46 (1.16)	$H(2) = 14.55, p = .001^*$
Anxiety	2.19 (1.3)	2.38 (1.35)	2.44 (1.28)	n.s.
Depression	2.48 (1.37)	2.55 (1.2)	2.42 (1.31)	n.s.
Weight gain	1.56 (0.75)	2.06 (1.17)	2.41 (1.56)	n.s.
Loss of smell/taste	1.63 (1.25)	1.89 (1.2)	2.36 (1.57)	$H(2) = 6.27, p = .044$
Nausea	1.22 (0.51)	1.62 (0.74)	2.29 (1.25)	$H(2) = 22.88, p < .001^*$
Loss of appetite	1.59 (0.97)	1.51 (0.89)	2.23 (1.31)	$H(2) = 14.11, p = .001^*$
Diarrhoea	1.44 (0.58)	1.77 (0.89)	2.2 (1.26)	$H(2) = 8.28, p = .016$
Altered consciousness	1.22 (0.51)	1.81 (1.13)	2.18 (1.3)	$H(2) = 15.09, p = .001^*$
Hot flushes	1.33 (0.78)	1.68 (0.96)	2.09 (1.34)	$H(2) = 8.66, p = .013$
Acid reflux	1.33 (0.62)	1.68 (1.12)	2 (1.3)	n.s.
Speech difficulty	1.04 (0.19)	1.38 (0.63)	1.94 (1.19)	$H(2) = 18.85, p < .001^*$
Numbness	1.11 (0.32)	1.36 (0.79)	1.85 (1.24)	$H(2) = 11.93, p = .003$
Visual disturbances	1.15 (0.46)	1.31 (0.7)	1.8 (1.22)	$H(2) = 10.6, p = .005$
Disorientation	1.07 (0.27)	1.47 (0.75)	1.77 (1.08)	$H(2) = 11.79, p = .003$
Rash	1.07 (0.27)	1.42 (0.93)	1.71 (1.08)	$H(2) = 11.39, p = .003$

Fever	1.15 (0.46)	1.15 (0.41)	1.67 (1.11)	$H(2) = 11.63, p = .003$
Weight loss	1.52 (1.05)	1.51 (0.95)	1.62 (1.09)	n.s.
Incontinence	1.22 (0.58)	1.25 (0.55)	1.32 (0.73)	n.s.
Itchy welts on skin	1.07 (0.27)	1.25 (0.62)	1.29 (0.7)	n.s.
Foot sores	1.07 (0.39)	1.15 (0.5)	1.29 (0.76)	n.s.
Vomiting	1.11 (0.42)	1.09 (0.35)	1.26 (0.62)	n.s.

* denotes p values below Sidak-correct alpha (i.e. non-null)

**Excluding a small portion of participants who reported asymptomatic or feeling completely better very quickly from the Recovered subgroup ($n = 15$)

Analysis on Current symptoms

We examined symptoms reported as present or absent over the past 1–2 days ($n = 124$, see Supplementary Table 11). Twelve symptoms appeared in less than 10% participants so were not included in the analyses. Pearson’s chi-square tests with Sidak correction adjusting the alpha level to .0013 showed that the COVID subgroups significantly differed in their current experience of semantic disfluency only, with the symptom appearing more commonly in the $C++$ (Severe) subgroup than the other subgroups.

Supplementary Table 11. Distribution of current symptoms in COVID subgroups: Recovered (R), Ongoing (Mild/Moderate ($C+$)) and Ongoing (Severe ($C++$)).

	Recovered (R) ($n = 8^{**}$)	Ongoing (Mild/Moderate) ($C+$) ($n = 50^{**}$)	Ongoing (Severe) ($C++$) ($n = 66$)	Chi-square tests
Current symptoms: Frequency (%)				
Fatigue	5 (62.5%)	37 (74%)	57 (86.4%)	n.s.
Difficulty concentrating	5 (62.5%)	36 (72%)	56 (84.8%)	n.s.
Forgetfulness	3 (37.5%)	30 (60%)	52 (78.8%)	$\chi^2(2) = 8.48, p = .014,$ $V = .262$
Brain fog	2 (25%)	34 (68%)	50 (75.8%)	$\chi^2(2) = 8.72, p = .013,$ $V = .265$
Breathing issues	2 (25%)	24 (48%)	49 (74.2%)	$\chi^2(2) = 12.7, p = .002,$ $V = .32$
Tip of the Tongue (ToT)	1 (12.5%)	28 (56%)	46 (69.7%)	$\chi^2(2) = 10.47, p = .005,$ $V = .291$

Night waking	5 (62.5%)	28 (56%)	43 (65.2%)	n.s.
Chest pain/tightness	1 (12.5%)	24 (48%)	43 (65.2%)	$\chi^2(2) = 9.57, p = .008,$ $V = .278$
Muscle/body pains	2 (25%)	22 (44%)	43 (65.2%)	$\chi^2(2) = 8.03, p = .018,$ $V = .254$
Headache	2 (25%)	20 (40%)	40 (60.6%)	$\chi^2(2) = 6.97, p = .031,$ $V = .237$
Semantic disfluency	1 (12.5%)	13 (26%)	40 (60.6%)	$\chi^2(2) = 17.21, p < .001,$ $V = .373^*$
Dizziness	4 (50%)	14 (28%)	38 (57.6%)	$\chi^2(2) = 10.13, p = .006,$ $V = .286$
Fast/irregular pulse	2 (25%)	17 (34%)	35 (53%)	n.s.
Difficulty sleeping	5 (62.5%)	19 (38%)	32 (48.5%)	n.s.
Head pressure	1 (12.5%)	15 (30%)	32 (48.5%)	$\chi^2(2) = 6.57, p = .037,$ $V = .23$
Sore throat	-	14 (28%)	28 (42.4%)	$\chi^2(2) = 7.02, p = .03,$ $V = .238$
Eye-soreness	2 (25%)	15 (30%)	27 (40.9%)	n.s.
Limb weakness	1 (12.5%)	16 (32%)	27 (40.9%)	n.s.
Anxiety	3 (37.5%)	19 (38%)	25 (37.9%)	n.s.
Blurred vision	1 (12.5%)	7 (14%)	25 (37.9%)	$\chi^2(2) = 9.18, p = .01,$ $V = .272$
Vivid dreams	2 (25%)	18 (36%)	24 (36.4%)	n.s.
Depression	4 (50%)	22 (44%)	23 (34.8%)	n.s.
Weight gain	2 (25%)	13 (26%)	22 (33.3%)	n.s.

Pins & needles	2 (25%)	9 (18%)	22 (33.3%)	n.s.
Cough	2 (25%)	15 (30%)	21 (31.8%)	n.s.
Abdominal pain	-	9 (18%)	21 (31.8%)	n.s.
Loss of smell/taste	1 (12.5%)	14 (28%)	19 (28.8%)	n.s.
Confusion	2 (25%)	9 (18%)	19 (28.8%)	n.s.
Excess thirst	6 (75%)	17 (34%)	18 (27.3%)	$\chi^2(2) = 7.38, p = .025,$ $V = .244$
Hot flushes	2 (25%)	6 (12%)	18 (27.3%)	n.s.
Drowsiness	2 (25%)	14 (28%)	18 (27.3%)	n.s.
Nausea	-	8 (16%)	16 (24.2%)	n.s.
Loss of appetite	-	7 (14%)	13 (19.7%)	n.s.
Speech difficulty	-	1 (2%)	13 (19.7%)	$\chi^2(2) = 9.98, p = .007,$ $V = .284$
Diarrhoea	1 (12.5%)	11 (22%)	13 (19.7%)	n.s.
Altered consciousness	-	7 (14%)	12 (18.2%)	n.s.
Acid reflux	-	8 (16%)	12 (18.2%)	n.s.
Numbness	-	3 (6%)	11 (16.7%)	n.s.
Weight loss	1 (12.5%)	7 (14%)	9 (13.6%)	n.s.
Visual disturbances	-	4 (8%)	9 (13.6%)	n.s.

* denotes *p* values below Sidak-correct alpha (i.e. non-null)

**Further excluding a small portion of participants who reported asymptomatic or feeling completely better very quickly from the Recovered subgroup ($n = 15$) and an additional group of participants who reported not experiencing current symptoms ($n = 22$)

Supplementary Table 12. Factor loadings for current symptoms (binary present/absent data).

Symptom	Component			
	1	2	3	4
Breathing issues	0.647			
Chest pain/tightness	0.593			
Fatigue	0.557	0.302		
Weight gain	0.534			
Cough	0.533			
Muscle/body pains	0.513	0.3		
Sore throat	0.508			
Night waking	0.488		0.337	
Vivid dreams	0.386			
Excess thirst	0.368			
Fever	0.352			
Loss of smell/taste	0.294			
Rash	0.254			
Vomiting	0.263			
Hot flushes		0.585		
Abdominal pain		0.584		
Pins & needles		0.513		0.317
Eye-soreness		0.489		
Fast/irregular pulse	0.412	0.473		
Headache		0.472	0.397	
Face/lips swelling		0.455		
Dizziness		0.447	0.32	
Blurred vision		0.446		0.343
Nausea		0.44		
Foot sores		0.425		
Head pressure		0.413	0.374	
Itchy welts		0.296		
Diarrhea		0.291		
Weight loss			0.684	
Loss of appetite			0.637	
Drowsiness			0.557	

Speech difficulty		0.554	0.316
Confusion		0.553	
Depression	0.423	0.46	
Difficulty sleeping		0.418	
Anxiety	0.375	0.389	
Delirium			0.741
Disorientation			0.737
Hallucinations			0.696
Visual disturbances			0.51
Altered consciousness			0.446