

Supplemental Figure 1. Original cohort characteristics prior to exclusion, N = 352

	WHO ordinal scores <6 N =287	WHO ordinal scores 6-9 N =65	
	N (%)		p- values
Symptoms			
Cough	147 (51.2)	39 (60.0)	-
Dyspnea	150 (52.3)	45 (69.2)	
Rhinorrhea	13 (4.5)	5 (7.7)	
Anosmia	16 (5.6)	3 (4.6)	
Dysgeusia or hypogeusia	27 (9.4%)	9 (13.8)	
Sore throat	15 (5.2)	4 (6.2)	
Fever	116 (40.4)	36 (55.4)	
Hemoptysis	4 (1.4)	0 (0)	
Nausea and/or vomiting	66 (23)	10 (15.4)	
Diarrhea	54 (18.8)	20 (30.8)	
Myalgia and/or arthritis	47 (16.4)	11 (16.9)	
Rash	1 (0.3)	0 (0)	
Altered mental status	16 (5.6)	5 (7.7)	
Asymptomatic	54 (18.8)	4 (6.2)	
Comorbidities			
Cardiovascular	172 (59.9)	41 (63.1)	-
Diabetes mellitus	87 (30.3)	24 (36.9)	
Lung disease	56 (19.5)	8 (12.3)	
Renal	34 (11.8)	6 (9.2)	
Liver	23 (8)	10 (15.4)	
HIV	2 (0.7)	2 (3.1)	
Malignancy on chemotherapy	4 (1.4)	1 (1.5)	
Solid organ transplant	6 (2.1)	1 (1.5)	
Dementia	16 (5.6)	3 (4.6)	
Others	46 (16.0)	6 (9.2)	
No known PMH	35 (12.2)	8 (12.3)	
Campuses			
Angleton	22 (7.7)	4 (6.2)	-
League City	44 (15.3)	17 (26.2)	

Clear Lake	51 (17.8)	9 (13.8)	
Galveston	165 (57.5)	32 (49.2)	
Non UTMB transfers	4 (1.4)	2 (3.1)	
Missing values	1 (0.3)	1 (1.5)	
Demographics			
Sex			
Male	165 (57.5)	44 (67.7)	0.17
Female	122 (14.5)	21 (32.3)	
Code status			
Comfort care	8 (2.8)	10 (15.4)	-
DNI	18 (6.3)	6 (9.2)	
Full code	261 (90.9)	47 (72.3)	
Missing values	0 (0)	2 (3.1)	

Clinical variables **	Mean (+ SD)		p- values
BMI	31.4 (7.43)	33.9 (8.86)	0.0408
O2 saturation, %	95.6 (6.58)	93.0 (5.62)	0.00133
NLR	5.69 (5.10)	9.92 (11.2)	0.00505
CRP	4.46 (7.62)	11.5 (10.2)	< 0.001
D-dimer	1.99 (7.86)	3.62 (16.6)	0.449
qSOFA	0.286 (0.475)	0.569 (0.558)	<0.001
LDH	402 (418)	783 (559)	<0.001
	N (%)		p- values
Categorical LDH			
Normal value	197 (68.6)	19 (29.2)	<0.001
1-2 x ULN	77 (26.38)	30 (46.2)	
> 2 x ULN	13 (4.5)	14 (21.5)	

** 2 subjects in ordinal score 6-9 group had consistently missed values on clinical variables

Supplemental Figure 2. Initial regression model of cohort, N=329

Parameter	DF	Estimate	Standard Error	Wald Chi-Square	Pr > ChiSq
Intercept	1	229.0	99.3147	5.3166	0.0211
Admission Date	1	-0.0102	0.00451	5.1249	0.0236
Biological Sex	1	0.5877	1.7130	0.1177	0.7315
Age	1	-0.00041	0.0597	0.0000	0.9946
Biological Sex*Age	1	-0.0130	0.0257	0.2566	0.6124
BMI	1	-0.0468	0.1064	0.1936	0.6599
BMI*Age	1	0.00203	0.00166	1.5088	0.2193
Admission SpO2	1	-0.0963	0.0439	4.8183	0.0282
Neut./Lymph. Ratio	1	0.0762	0.0288	6.9776	0.0083
Procalcitonin	1	0.2200	0.5481	0.1612	0.6881
Admission D-Dimer	1	-0.0204	0.0137	2.2070	0.1374
LDH	1	1.1513	0.2995	14.7743	0.0001
CRP	1	0.2342	0.4337	0.2917	0.5891
qSOFA Score	1	0.8852	0.3196	7.6689	0.0056

*Note that complete case exclusion not applied for this table

Supplemental Figure 3. DNI Sensitivity Analysis

Odds Ratio Estimates			
Effect	Point Estimate	95% Wald Confidence Limits	
Age	1.044	1.020	1.069
BMI	1.082	1.034	1.133
Admission SpO2	0.907	0.834	0.987
Admission Date	0.989	0.981	0.998
Neut./Lymph. Ratio	1.078	1.023	1.137
Admission D-Dimer	0.979	0.954	1.005
LDH	3.379	2.043	5.588
qSOFA Score	2.240	1.218	4.118