

Supplementary appendix

NUCCIA MORICI

CONTENTS

COMMITTEES AND INVESTIGATORS.....	2
FIRST PROTOCOL VERSION (VERSION 1.0)	3
LAST PROTOCOL VERSION (VERSION 4.0)	27
SUMMARY OF MAIN PROTOCOL CHANGES	51
COMPLETE LIST OF STUDY OUTCOMES	54
CTPA COMPARISON BETWEEN GROUPS	59
CHARACTERISTICS OF SUBJECT WITH PULMUNARY EMBOLISM.....	60
ENROLLMENTS INFORMATIONS	62

COMMITTEES AND INVESTIGATORS

Steering Committee:

Nuccia Morici, Marco Cattaneo

Clinical Event Committee (CEC):

Gabriele Crimi, Stefano De Servi

Data Safety Monitoring Board:

Data monitoring was managed by a data manager and an independent researcher (Dr. Arianna Pani), specialized in pharmacology, which checked the events every two months.

Dr. Enrico Ammirati, Marco Valgimigli and Stefano Savonitto has only participated to the trial as consultant in the planning phases of the study.

FIRST PROTOCOL VERSION (VERSION 1.0)

Enoxaparin for thromboprophylaxis in hospitalized COVID-19 patients: comparison of 40 mg o.d. versus 40 mg b.i.d. A randomized Clinical Trial

Internal Reference Number / Short title: X-COVID 19

Ethics Ref:

Date and Version No: April 2020, Version 1.0

Chief Investigators:

Marco Cattaneo, Dipartimento di Scienze della Salute, Università Degli Studi di Milano, Milan, Italy; Unità di Medicina 2, ASST Santi Paolo e Carlo, Milan, Italy;
Nuccia Morici, ASST Grande Ospedale Metropolitano Niguarda, Milan, Italy.

Investigators:

Enrico Ammirati, Jacopo Oreglia, Alice Sacco, Claudia Marini, Massimo Puoti, Linda Guarnieri (ASST Grande Ospedale Metropolitano Niguarda, Milan, Italy), Gianfranco Aprigliano (Istituto Clinico Città Studi, Milan, Italy), Francesco Bedogni (IRCCS Policlinico San Donato, Milan, Italy), Gaetano Maria De Ferrari, Fabrizio D'Ascenzo (AOU Città della Salute e della Scienza, Presidio Molinette, Torino, Italy), Giovanni Forleo (Ospedale Sacco, Milan, Italy), Sergio Leonardi, Claudio Montalto (IRCCS Policlinico San Matteo, Pavia, Italy), Paolo Bonfanti (ASST Monza – Ospedale di Monza, Monza, Italy), Ilaria Battistoni, Matteo Francioni (AOU Ospedali Riuniti di Ancona, Ancona, Italy), Marco Metra, Pompilio Faggiano (ASST Spedali Civili di Brescia, Brescia, Italy), Annamaria Municinò (Ospedale Andrea Gallino, ASL 3 Genovese, Genova, Italy), Giulio Stefanini (IRCCS Istituto Clinico Humanitas, Rozzano, Milan, Italy), Stefano Savonitto (ASST Lecco, Ospedale A. Manzoni, Lecco, Italy)

Sponsor:

ASST Grande Ospedale Metropolitano Niguarda, Milan, Italy

Funder:

None

Chief Investigator Signature:

Marco Cattaneo, Nuccia Morici

Statistician Signature:

Gianfranco Alicandro

Confidentiality Statement: This document contains confidential information that must not be disclosed to anyone other than the Sponsor, the Investigator Team, HRA, host organisation, and members of the Research Ethics Committee, unless authorised to do so.

TABLE OF CONTENTS

1.	KEY TRIAL CONTACTS	6
2.	SYNOPSIS	7
3.	ABBREVIATIONS	9
4.	BACKGROUND AND RATIONALE	10
5.	OBJECTIVES AND OUTCOME MEASURES	11
6.	TRIAL DESIGN	12
7.	PARTICIPANT IDENTIFICATION	13
7.1	Trial participants	13
7.2	Inclusion Criteria	13
7.3	Exclusion Criteria	14
8	TRIAL PROCEDURES	14
8.1	Recruitment	15
8.2	Informed Consent	15
8.3	Screening and Eligibility Assessment	15
8.4	Randomisation, blinding and code-breaking	16
8.5	Baseline Assessments	16
8.6	Subsequent Visits	16
8.7	Discontinuation/Withdrawal of Participants from Trial Treatment	17
8.8	Definition of End of Trial	18
9.	COMPLIANCE WITH TRIAL TREATMENT	18
10.	ACCOUNTABILITY OF TRIAL TREATMENT	18
11.	CONCOMITANT MEDICATION	18
12.	POST-TRIAL TREATMENT	18
13.	SAFETY REPORTING	18
13.1.	Definitions	18
13.2.	Causality	19
13.3.	Procedures for Recording Adverse Events	19
13.4.	Reporting Procedures for Serious Adverse Events	20
13.5.	Expectedness	20
13.6.	SUSAR Reporting	20
13.7.	Safety Monitoring Committee	20
14.	STATISTICS	20
14.1.	Description of Statistical Methods	20
14.2.	Number of Participants	21
14.3.	Populations Analysis	21
14.4.	Criteria for the Termination of the Trial	22
14.5.	Procedure for Accounting for Missing, Unused, and Spurious Data	22

14.6. Inclusion in Analysis	22
14.7. Procedures for Reporting any Deviation(s) from the Original Statistical Plan	22
15. DATA MANAGEMENT	22
15.1. Source Data	22
15.2. Access to Data	23
15.3. Data Recording and Record Keeping	23
16. QUALITY ASSURANCE PROCEDURES	23
17. SERIOUS BREACHES	23
18. ETHICAL AND REGULATORY CONSIDERATIONS	23
18.1. Declaration of Helsinki	23
18.2. Guidelines for Good Clinical Practice	24
18.3. Approvals	24
18.4. Reporting	24
18.5. Participant Confidentiality	24
18.6. Expenses and Benefits	24
19. FINANCE AND INSURANCE	24
19.1. Funding	24
19.2. Insurance	24
20. PUBLICATION POLICY	25
21. REFERENCES	26

1. KEY TRIAL CONTACTS

Chief Investigator	Marco Cattaneo, Dipartimento di Scienze della Salute, Università Degli Studi di Milano, Milan, Italy; Unità di Medicina 2, ASST Santi Paolo e Carlo, Milan, Italy. Phone 0250323095 - 0281844638; fax 0250323089; e-mail marco.cattaneo@unimi.it Nuccia Morici, Unità di Cure Intensive Cardiologiche Dipartimento Cardio-toraco-vascolare, ASST Grande Ospedale Metropolitano Niguarda, Milan, Italy. Phone 0264442565, fax 0264442818, e-mail nuccia.morici@ospedaleniguarda.it
Sponsor	ASST Grande Ospedale Metropolitano Niguarda, Milan, Italy
Clinical Trials Unit	Unità di Medicina, ASST Santi Paolo e Carlo, Milan, Italy / Malattie Infettive, ASST Grande Ospedale Metropolitano Niguarda, Milan, Italy
Statisticians	Gianfranco Alicandro, e-mail gianfranco.alicandro@unimi.it
Committees	Comitato Etico Scientifico INMI Lazzaro Spallanzani (Ethical Committee) Steering Committee: Marco Cattaneo, Nuccia Morici, Enrico Ammirati, Marco Valgimigli, Stefano Savonitto.

2. SYNOPSIS

Trial Title	Enoxaparin for thromboprophylaxis in hospitalized COVID-19 patients: comparison of 40 mg o.d. versus 40 mg b.i.d. A randomized Clinical Trial
Internal ref. no. (or short title)	X-Covid 19
Promoter	ASST Grande Ospedale Metropolitano Niguarda
Principal Investigator	Marco Cattaneo / Nuccia Morici
EudraCT Number	2020-001708-41
Clinical Phase	Phase IIIb
Trial Design	Open-label Multicentre Prospective Controlled Randomized Trial in patients with SARS-CoV-2 infection. Patients will be randomized 1:1 to 40 mg subcutaneous enoxaparin o.d. versus 40 mg enoxaparin b.i.d within 12 hours after hospitalization.
Trial Participants <i>Inclusion criteria</i>	<ul style="list-style-type: none"> All-comers patients aged ≥ 18 years and admitted to hospital with laboratory-confirmed SARS-CoV-2 infection.
<i>Exclusion criteria</i>	<ul style="list-style-type: none"> Patients admitted directly to an intensive care unit; Estimated creatinine clearance < 15 ml/min/1.73m²; Patients needing anticoagulant for prior indication; Patients at high bleeding risk or experiencing clinically significant bleeding; Patients involved in clinical trial protocols restricting patients from concurrent studies; Any other significant disease or disorder which, in the opinion of the investigator, may either put the participants at risk because of participation in the trial, or may influence the result of the trial, or the participant's ability to participate in the trial.
Planned Sample Size	A sample of 2712 subjects is required to test the superiority hypothesis with 80% power according to a two-sided 0.05 level of significance that 40 mg enoxaparin b.i.d. will decrease the primary endpoint compared to 40 mg enoxaparin o.d. An event rate of 10% was hypothesized in the control arm and 7% in the treatment arm, corresponding to a 33% relative risk reduction.
Treatment duration	In-hospital (at least 30 days of therapy)
Follow up duration	30 days from the enrolment
Planned Trial Period	6 months
Primary Objective	To compare the effects of 40 mg subcutaneous enoxaparin o.d. versus 40 mg enoxaparin b.i.d on the incidence of venous thromboembolism (VTE) [a composite of incident asymptomatic and symptomatic deep vein thrombosis (DVT) diagnosed by serial compression ultrasonography (CUS), and symptomatic pulmonary embolism (PE) diagnosed by CT scan], in patients with SARS-CoV-2 infection.

<p>Secondary Objectives</p>	<ul style="list-style-type: none"> • To compare the effects of 40 mg subcutaneous enoxaparin o.d. versus 40 mg enoxaparin b.i.d on the incidence of in-hospital major complications, defined as the composite of death, VTE, use of mechanical ventilation, stroke, acute myocardial infarction and admission to an intensive care in patients with SARS-CoV-2. • To compare each single component of the primary endpoint between the two groups. • To compare maximum sequential organ failure assessment (SOFA) score between the two groups. • To compare C-reactive protein, D-dimer, IL-6 and hs-troponin levels (as % above the upper reference limit [URL]) among the two groups. • To compare the incidence of SARS-CoV-2-related Acute Respiratory Distress Syndrome (ARDS) between the two groups. • To compare length of hospital stay between the two groups. • To compare measures of right ventricular function at trans-thoracic echocardiography or CT between admission and follow-up, whenever available.
<p>Safety Endpoints</p>	<ul style="list-style-type: none"> • To compare the incidence of bleeding events according to the <i>ISTH bleeding scale</i> between the two groups. • To compare the incidence of bleeding events according to <i>BARC classification</i> between the two groups. Only BARC 3 and 5 will be considered for secondary endpoint. • To assess the incidence of heparin-induced thrombocytopenia in the two groups.

3. ABBREVIATIONS

AE	Adverse event
AR	Adverse reaction
BARC	Bleeding Academic Research Consortium
CI	Chief Investigator
CRF	Case Report Form
CT	Clinical Trials
CTA	Clinical Trials Authorisation
CTRG	Clinical Trials and Research Governance
DMC/DMSC	Data Monitoring Committee / Data Monitoring and Safety Committee
DSUR	Development Safety Update Report
GCP	Good Clinical Practice
GP	General Practitioner
IB	Investigator's Brochure
ICF	Informed Consent Form
ICH	International Conference on Harmonisation
IMP	Investigational Medicinal Product
IRB	Independent Review Board
PI	Principal Investigator
PIL	Participant/ Patient Information Leaflet
REC	Research Ethics Committee
SAE	Serious Adverse Event
SAR	Serious Adverse Reaction
SDV	Source Data Verification
SMPC	Summary of Medicinal Product Characteristics
SOP	Standard Operating Procedure
SOFA	Sequential Organ Failure Assessment
SUSAR	Suspected Unexpected Serious Adverse Reactions
TMF	Trial Master File

4. BACKGROUND AND RATIONALE

Critically ill medical patients are at heightened risk for venous thromboembolism (VTE) and, for this reason, they are treated with prophylactic doses of low molecular weight heparin (LMWH) (for instance, 40 mg enoxaparin). Patients with COVID-19, which is caused by SARS-CoV-2 infection, are apparently at particularly high risk of VTE, as a consequence of excessive activation of the hemostatic system which, in the most severe cases, could also be associated with the formation of microthrombi and clinically relevant disseminated intravascular coagulation.¹⁻⁴

Based on personal communications by medical doctors working mostly in Northern Italy, there are concerns about the efficacy of thromboprophylaxis with low-molecular-weight heparin (LMWH) at the standard doses in these patients, given their apparently very high risk of thromboembolic complications. Therefore, in many Northern Italian hospitals these patients are given LMWH at much higher doses than those recommended for thromboprophylaxis: in some hospitals, full anticoagulant doses (100 U/Kg b.i.d.) are being used even in the absence of evidence of ongoing thromboembolic disease. This practice does not take into due account the obvious risk of major, potentially fatal bleeding that is associated with high dosing, in the absence of any evidence of higher efficacy of high doses compared to standard prophylactic doses.

An additional rationale for the use of high dose LMWH in many hospitals is that hyper-activation of the hemostatic system in COVID-19 patients could lead to the formation of microthrombi in pulmonary vessels⁵, causing further respiratory distress, and in vessels of other vital organs, causing multi-organ failure (MOF). However, there is no demonstration that this therapeutic approach decreases the risk of microangiopathic thrombosis in these patients, as well as, for that matter, in other types of patients.

Given this scenario, we designed a randomized trial comparing standard prophylactic dose of subcutaneous enoxaparin (40 mg o.d.) with higher dose (40 mg b.i.d) with the aim of testing whether high-dose thromboprophylaxis is more effective than standard dose in preventing TEV in COVID-19 patients. This study would also allow us to test whether double-dose prophylaxis with enoxaparin

would favourably affect the natural history of the disease, by decreasing the incidence of organ dysfunction associated with the formation of microthrombi in vital organs.

5. OBJECTIVES AND OUTCOME MEASURES

Objectives	Outcome measures	Timepoint(s) of evaluation of these outcome measure (if applicable)
<p>Primary Objective</p> <ul style="list-style-type: none"> To compare the incidence of hospital-associated venous thromboembolism (HA-VTE) between the two groups (40 mg o.d. vs 40 mg b.i.d subcutaneous enoxaparin) 	<ul style="list-style-type: none"> Number (percentage) of patients who will experience the composite event. Relative risk will evaluate the association between the study outcome and treatment. 	<p>30-day</p>
<p>Secondary Objectives</p> <ul style="list-style-type: none"> To compare the incidence of in-hospital major complications, defined as the composite of death, HA-VTE, use of mechanical ventilation, major stroke, acute myocardial infarction and admission to an intensive care between the two groups. To compare each single component of the primary endpoint between the two groups. To compare maximum sequential organ failure assessment (SOFA) score between the two groups. To compare C-reactive protein, D-dimer, IL-6 and hs-troponin levels (as % above the upper reference limit [URL]) among the two groups. 	<ul style="list-style-type: none"> Number (percentage) of patients who will experience the event. Relative risk will evaluate the association between the study outcome and treatment. Number (percentage) of patients who will experience the event. Relative risk will evaluate the association between the study outcome and treatment. Maximum sequential organ failure assessment (SOFA) score summarized by median and interquartile range (IQR) or mean and standard deviation according to variable distribution. Laboratory parameters will be indicated as median and interquartile range (IQR) or mean and standard deviation according to variable distribution. 	<p>30-day</p> <p>30-day</p> <p>Discharge</p> <p>Discharge</p> <p>30-day</p> <p>Discharge</p> <p>30-day</p>

<ul style="list-style-type: none"> • To compare the incidence of SARS-CoV-2- related Acute Respiratory Distress Syndrome (ARDS) between the two groups. • To compare length of hospital stay • To compare measures of right ventricular function at trans-thoracic echocardiography or CT between admission and follow-up, whenever available. 	<ul style="list-style-type: none"> • <i>Number (percentage) of patients who will experience</i> the event. <i>Relative risk</i> will evaluate the association between the study outcome and treatment. • <i>Median and interquartile range (IQR) or mean and standard deviation</i> according to variable distribution. • Echocardiographic parameters will be summarized by <i>median and interquartile range (IQR) or mean and standard deviation</i> according to variable distribution. 	
<p>Safety endpoints</p> <ul style="list-style-type: none"> • To assess the occurrence of bleeding events, according to the two classifications selected • To assess the incidence of heparin-induced thrombocytopenia between the two groups. 	<ul style="list-style-type: none"> • <i>Number (percentage) of patients who will experience</i> the event. <i>Relative risk</i> will evaluate the association between the study outcome and treatment. • The incidence rate will be described across groups 	<p>During hospital stay</p> <p>During hospital stay</p>

6. TRIAL DESIGN

Prospective controlled randomized, open-label multi-center trial.

Patients with COVID-19 (any hospitalized patients) with laboratory-confirmed SARS-CoV-2 infection will be randomized 1:1 to 40 mg enoxaparin/day subcutaneous versus 40 mg enoxaparin b.i.d. within 12 hours after hospitalization. Standard dosage of Proton Pump Inhibitor (PPI) will be prescribed to each enrolled patient to reduce gastrointestinal (GI) bleeding risk. The chronic kidney disease epidemiology (CKD-EPI) formula will be used for eGFR calculation.

The study conduct will not change the usual clinical practice in terms of treatments and diagnostic procedures.

Patients will be monitored every 3 days with D-dimer, fibrinogen, complete blood count, lactate dehydrogenase (LDH), PT, APTT, C-reactive protein, procalcitonin only in the case of superinfection by a different agent, ferritin, creatine phosphokinase (CK), CK-MB, albumin, serum creatinine, aspartate transaminase (AST), alanine transaminase (ALT), high-sensitivity troponin, serum bilirubin and interleukine-6 along with other examinations according to clinical practice.

Patients will be monitored every seven days from the enrolment with CUS of the lower limbs for proximal DVT screening. For patients who display signs and symptoms of DVT, CUS will be performed as soon as possible in the same day. Proximal DVT, namely DVT of iliac, femoral and popliteal veins down to popliteal trifurcation, is defined by cross-sectional vein incompressibility or direct thrombus imaging with vein enlargement. Distal DVT might be reported if distal segments are explored.

For patients who display signs and symptoms of pulmonary embolism, a CT scan will be performed as soon as possible in the same day according to the indication of the attending physician.

Multidetector CT pulmonary angiography is the method of choice for imaging the pulmonary vasculature in patients with suspected pulmonary embolism, that is defined by contrast-filling defects. With regard to secondary objective, the following definitions apply.

All-cause mortality includes all deaths, regardless of whether the cause of death is determined.

Cardiovascular death is defined as death due to atherosclerotic coronary heart disease, cerebrovascular accident or (complication of) peripheral embolization, and includes deaths due to acute MI, stroke, sudden death, non-sudden death, unwitnessed death, and procedure-related deaths. Non-cardiovascular death includes deaths due to all other causes.

Myocardial Infarction denotes the presence of acute myocardial injury detected by abnormal cardiac biomarkers [Cardiac troponin I (cTnI) and T (cTnT) increased above the 99th percentile upper reference limit (URL)] in the setting of evidence of acute myocardial ischaemia.

Stroke is defined as any new neurological deficit lasting >24 hours. CT or MRI are required to confirm the stroke and to distinguish between ischemic and hemorrhagic stroke. Strokes will be also classified as fatal or nonfatal: nonfatal stroke will be also classified as disabling and non disabling.

The total duration of the study is 6 months for enrolment, follow-up, data collection, cleaning and analysis.

7. PARTICIPANT IDENTIFICATION

7.1 Trial participants

Patients directly admitted to hospital for suspected SARS-CoV-2 infection will be screened for randomization.

7.2 Inclusion Criteria

All the following inclusion criteria will be adopted:

All-comers patients aged ≥ 18 years and admitted to hospital with laboratory-confirmed SARS-CoV-2 infection.

7.3 Exclusion Criteria

The following patients will not be enrolled:

- Patients admitted directly to intensive care unit.
- Patients with estimated creatinine clearance < 15 ml/min/1.73m².
- Patients needing anticoagulant for prior indication.
- Patients at high bleeding risk or experiencing clinically significant bleeding;
- Patients involved in clinical trial protocols restricting patients from concurrent studies
- Patients with any other significant disease or disorder which, in the opinion of the Investigator, may either put the participants at risk because of participation in the trial, or may influence the result of the trial, or the participant's ability to participate in the trial.

8 TRIAL PROCEDURES

Eligibility will be evaluated by the attending physician after the first visit and after confirmation of the diagnosis. The patient, properly informed about the study, will be able to decide to sign the informed consent and the privacy form. The physician will obtain a detailed anamnesis and he/she will carefully check the clinical parameters needed to define tissue perfusion (as detailed by the inclusion criteria) along with body temperature. The following laboratory data will be checked: haemochrome, renal function, serum electrolytes, liver function, coagulation, blood gas analysis. Laboratory examinations will be sequentially evaluated according to the flow chart timeline (section 8.6). If the patient develops a condition needing an anticoagulation, an increase in enoxaparin dosage to 100 UI/kg (1 mg/kg) b.i.d. might be considered if indicated and in the absence of contraindications. Serial CUS will be performed, the first on admission and then every seven days. Bleeding events will be registered and graded according to the International Society on Thrombosis and Haemostasis (ISTH) bleeding scale and Bleeding Academic Research Consortium (BARC) scale. Also BARC 2 bleedings will be recorded.

Pre-specified subgroup analyses will be performed for 1) D-dimer categories (>1 <2 , 2, 3, 4, 5, 6 fold the ULN of the reference range); 2) estimated creatinine clearance ($<$ versus >60 ml/min/1.73 m²); 3) severity of the respiratory symptoms (mild versus moderate-severe defined as respiratory rate ≥ 30 breaths/min; arterial oxygen saturation $\leq 93\%$ at rest; PaO₂/FiO₂ ≤ 300 mmHg); 4) concomitant antiplatelet therapy.

The identification of adverse reactions will be continuous during hospitalisation. The data will be recorded in an eCRF.

8.1 Recruitment

Patients recruitment will be performed at the Cardiology Department, Malattie Infettive and Internal Medicine, ASST Grande Ospedale Metropolitano Niguarda, Milan, Italy; Medicina, Ospedale San Paolo, ASST Santi Paolo e Carlo, Milan, Italy; Istituto Clinico Città Studi, Milan, Italy; IRCCS Policlinico San Donato, Milan, Italy; AOU Città della Salute e della Scienza, Presidio Molinette, Torino, Italy; Ospedale Sacco, Milan, Italy; IRCCS Policlinico San Matteo, Pavia, Italy; ASST Monza – Ospedale di Monza, Monza, Italy; AOU Ospedali Riuniti di Ancona, Ancona, Italy; ASST Spedali Civili di Brescia, Brescia, Italy; Ospedale Andrea Gallino, ASL 3 Genovese, Genova, Italy; IRCCS Istituto Clinico Humanitas, Rozzano, Milan, Italy; ASST Lecco, Ospedale A. Manzoni, Lecco, Italy.

8.2 Informed Consent

The participant must personally sign and date the latest approved version of the Informed Consent form before any trial specific procedures.

Written and verbal versions of the Participant Information and Informed Consent will be presented to the participants detailing no less than: the exact nature of the trial; what it will involve the participant; the implications and constraints of the protocol; the known side effects and any risks involved in taking part. It will be clearly stated that the participant is free to withdraw from the trial at any time for any reason without prejudice to future care and with no obligation to give the reason for withdrawal.

The participant will be allowed as much time as wished, taking into consideration the urgency of treatments, to consider the information and the opportunity to ask the Investigator, his/her GP or other independent parties to decide whether he/she will participate in the trial. Written Informed Consent will then be obtained by means of participant dated signature and dated signature of the person who presented and obtained the Informed Consent. The person who obtained the consent must be suitably qualified and experienced, and authorized to do so by the Chief/Principal Investigator. The original signed Informed Consent will be given to the participant and a copy will be retained at the trial site.

8.3 Screening and Eligibility Assessment

As clinically indicated, the screening procedures will be performed throughout demographics, medical history, concomitant medication, physical examination, laboratory tests and radiological findings (chest CT scans are encouraged over chest X-rays).

The screening procedure will be immediately followed by recruitment at arrival in the ward (patients will not be randomized in the emergency department), if allowed on basal evaluation and patient informed consent signature.

8.4 Randomisation, blinding and code-breaking

This study will be a two-parallel group randomized, open-label, multi-center trial in hospitalized COVID-19 patients. Patients will be randomized by the attending physicians using an IVRS that randomizes at a ratio of 1:1 upon determination of eligibility.

A total sample of 2712 subjects is required to test the superiority hypothesis with 80% power according to a two-sided 0.5 level of significance that 40 mg enoxaparin subcutaneously bid/day (treatment arm) will decrease the primary endpoint compared to 40 mg enoxaparin/day (prophylaxis dose, that is considered the control arm as represents the standard of care for hospitalized patients that are bedridden). An absolute reduction of 3% is foreseen in the treatment arm (enoxaparin 40 mg b.i.d); an event rate of 10% was considered in the control arm (enoxaparin 40 mg o.d.) and 7% in the treatment arm.

The duration of the enrolment is 3 months from the obtaining of the approvals. The total duration of the study is 6 months for follow-up (1 months), data collection, cleaning and analysis (2 months). Randomisation will be done at the same visit as the baseline visit. There will not be a run-in period.

8.5 Baseline Assessments

Basal evaluation (V1)

Every patient will be informed about the objectives of our study and he/she will be asked for consent to the trial. At basal visit and following ones, the physician will perform: physical examination, hemodynamic parameters and blood gas analysis check; CUS. Echocardiogram might be performed if requested by the attending physician.

The occurrence of organ dysfunction and failure, from admission through days spent in intensive care, will be assessed by the Sequential Organ Failure Assessment (SOFA) score.

8.6 Subsequent Visits

After basal assessment, the following visits will be performed according to local clinical practice a check-up at discharge will be performed. Echocardiograms might be performed during hospitalization according to the indication of the attending physician. In case mechanical ventilation is required, ventilation modes and eventual variations of parameters will be recorded.

The schedule of clinical visits and assessment procedures is shown in the following flow chart.

Flow chart

Visit	V1	V2	V2	V3	V4
Time interval	Basal	To be checked each three days	To be checked each seven days	Discharge	30-day from the enrolment
Informed consent	X				
Time from onset symptoms to enrolment	X				
Physical examination	X				
Height	X				
Weight	X				
Anamnesis	X				X
BP (s/d/m)	X				
HR	X	X		X	
CUS	X		X		
Hb	X	X		X	
Creatinine	X	X		X	
Platelets	X	X		X	
Leucocytes	X	X		X	
INR	X	X		X	
Bilirubin	X	X		X	
AST	X	X		X	
ALT	X	X		X	
LDH	X	X		X	
CRP	X	X		X	
Troponin (HS)	X	X		X	
IL-6	X	X		X	
D-dimer	X	X		X	
Fibrinogen	X	X		X	
SOFA score	X	X			

8.7 Discontinuation/Withdrawal of Participants from Trial Treatment

Each participant has the right to withdraw from the trial at any time. In addition, the Investigator may discontinue a participant from the trial at any time if the Investigator considers it necessary for any reason including:

- Ineligibility (either arising during the trial or retrospectively having been overlooked at screening)
- Significant protocol deviation
- Significant non-compliance with treatment regimen or trial requirements
- An adverse event which requires discontinuation of trial treatment or results in inability to continue to comply with trial procedures
- Disease progression which requires discontinuation of trial treatment or results in inability to continue to comply with trial procedures
- Withdrawal of Consent

Withdrawal from the trial will result in exclusion of the data for that participant from analysis

The reason for withdrawal will be recorded in the eCRF.

If the participant is withdrawn due to an adverse event, the Investigator will arrange for follow-up visits or telephone calls until the adverse event has resolved or stabilised.

8.8 Definition of End of Trial

The end of trial is the date of the last visit/ telephone follow up/ home visit of the last participant.

9. COMPLIANCE WITH TRIAL TREATMENT

All patients will be treated in hospital ward, so compliance will be guaranteed by the investigators.

10. ACCOUNTABILITY OF TRIAL TREATMENT

The study does will involve drugs usually administrated in clinical practice.

11. CONCOMITANT MEDICATION

There are not any contraindicated medications. All used drugs, as per clinical indication, will be reported in the eCRF.

12. POST-TRIAL TREATMENT

At the end of the study period, all surviving patients will be treated as per current clinical practice in the participating Centers.

13. SAFETY REPORTING

13.1. Definitions

Adverse Event (AE)	Any untoward medical occurrence in a participant to whom a medicinal product or diagnostic-interventional procedure has been administered, including occurrences which are not necessarily caused by or related to that product or procedure.
Adverse Reaction (AR)	An untoward and unintended response in a participant to a medicinal product which is related to any dose administered to that participant. The phrase "response to an investigational medicinal product" means that a causal relationship between a trial medication and an AE is at least a reasonable possibility, i.e. the relationship cannot be ruled out. All cases judged by either the reporting medically qualified professional or the Sponsor as having a reasonable suspected causal relationship to the trial medication qualify as adverse reactions.
Serious Adverse Event (SAE)	A serious adverse event is any untoward medical occurrence that: <ul style="list-style-type: none">• Results in death

	<ul style="list-style-type: none"> • Is life-threatening • Requires inpatient hospitalisation or prolongation of existing hospitalisation • Results in persistent or significant disability/incapacity • Consists of a congenital anomaly or birth defect. <p>Other ‘important medical events’ may also be considered serious if they jeopardise the participant or require an intervention to prevent one of the above consequences.</p> <p>NOTE: The term "life-threatening" in the definition of "serious" refers to an event in which the participant was at risk of death at the time of the event; it does not refer to an event which hypothetically might have caused death if it were more severe.</p>
--	---

NB: to avoid confusion or misunderstanding of the difference between the terms “serious” and “severe”, the following note of clarification is provided: “Severe” is often used to describe intensity of a specific event, which may be of relatively minor medical significance. “Seriousness” is the regulatory definition supplied above.

13.2. Causality

The relationship of each adverse event to trial treatment must be determined by a medically qualified individual according to the following definitions:

Related: The adverse event follows a reasonable temporal sequence from trial treatment start. It cannot reasonably be attributed to any other cause.

Not Related: The adverse event is probably produced by the participant’s clinical state or by other modes of therapy administered to the participant.

13.3. Procedures for Recording Adverse Events

All AEs occurring during the trial / or until the following two months that are observed by the Investigator or reported by the participant, will be recorded on the eCRF, whether or not attributed to trial treatment.

The following information will be recorded: description, date of onset and end date, severity, assessment of relatedness to trial treatment, other suspect drug or device and action taken. Follow-up information should be provided as necessary.

The severity of events will be assessed on the following scale: 1 = mild, 2 = moderate, 3 = severe. AEs considered related to the trial treatment as judged by a medically qualified investigator will be followed either until resolution, or the event is considered stable.

It will be left to the Investigator’s clinical judgment to decide whether an AE is of enough severity to require the participant’s removal from treatment. A participant may also voluntarily withdraw from treatment due to what he or she perceives as an intolerable AE. If either of these occurs, the participant must undergo an end of trial assessment and be given appropriate care under medical supervision until symptoms cease, or the condition becomes stable.

13.4. Reporting Procedures for Serious Adverse Events

All SAEs (other than those defined in the protocol as not requiring reporting) must be reported on the SAE reporting form to Eudravigilance within 24 hours of the Site Study Team becoming aware of the event.

Review of SAEs must be timely, taking into account the reporting time for a potential SUSAR.

13.5. Expectedness

Expectedness will be determined according to the Investigators' Brochure/Summary of Product Characteristics.

13.6. SUSAR Reporting

All SUSARs will be reported by the CI to the relevant Competent Authority and to the REC and other parties as applicable. For fatal and life-threatening SUSARS, this will be done no later than 7 calendar days after the investigator team is first aware of the reaction. Any additional information will be reported within 15 calendar days. Treatment codes will be un-blinded for specific participants.

13.7. Safety Monitoring Committee

The aims of this committee include:

- To pick up any trends, such as increases in un/expected events, and take appropriate action
- To seek additional advice or information from investigators where required
- To evaluate the risk of the trial continuing and take appropriate action where necessary

14. STATISTICS

14.1. Description of Statistical Methods

All protocol-defined endpoints will be measured from the time of randomization. Descriptive statistics of baseline characteristics of patients randomized to the two groups will be reported. Continuous variables will be summarized by means, standard deviations, medians, inter-quartile ranges and minimum and maximum values. Categorical variables will be summarized by frequencies and percentages.

Data will be analysed according to the intention-to-treat principle. In addition, a per-protocol analysis of the primary endpoint, considering patients without major protocol violations (i.e. patients who crossed over), will be performed to evaluate the robustness of the data.

A test for difference of proportions will be carried out to compare in-hospital composite events among patients randomized to the two groups. A relative risk will be calculated to quantify the magnitude of the half anticoagulant effect.

For what regards continuous secondary endpoints, maximum SOFA score will be compared among the study groups through the t-test or Wilcoxon rank-sum test, according to the distribution of study endpoints. The same approach will be applied for laboratory examinations and times included in the secondary endpoints.

All tests will be two-sided and a p-value of 0.05 or less will be considered statistically significant. No adjustments for multiple testing will be applied in the analysis of secondary endpoints and in the subgroup analysis, therefore, inferences drawn from the reported p-values and 95% CIs will be considered exploratory. If a p-value is less than 0.001 it will be reported as “<0.001”.

14.2. Number of Participants

Summary Statements

The proportion for the primary endpoint is estimated to be 0.07 in the half-dose anticoagulant dose and 0.10 in the standard group. With a type I error α set to 0.05 and a power $1 - \beta$ set to 0.8, a total of 2712 subjects will be enrolled in the trial.

14.3. Populations Analysis

For this trial, the following populations will be defined and used in the analysis and/or presentation of the data.

Safety population: The safety population will be classified according to the actual treatment on enoxaparin 40 mg b.i.d. or control, received.

Intent-to-treat (ITT) population: Treatment classification for ITT analysis will be based on the randomized treatment.

Per-Protocol (PP) population: The PP population will be defined as all ITT patients who received assigned to enoxaparin 40 mg b.i.d. and will be without major protocol violations. Specifically, subjects may be excluded from PP population if any of the following criteria are met:

- Inclusion/Exclusion criteria violations;
-

The PP definition for each patient will be finalized before database lock and study unblinding.

The primary and secondary efficacy analyses will be based on the ITT population. Analyses based on the PP population will be considered supportive.

14.4. Criteria for the Termination of the Trial

Criteria refers to the above reported statistical plan (section 14.2).

14.5. Procedure for Accounting for Missing, Unused, and Spurious Data

All data will be checked for consistency and plausibility based on the nature of each collected variable before any statistical analysis. Data falling beyond limits of plausibility will be checked in the medical records and in the CRF to correct for possible typos or transcription errors. Changes in the database will be recorded together with date and name of the physician performing that correction and with the reason for correction.

Methods of multiple imputation will be considered if the missingness rate will be more than 20%.

14.6. Inclusion in Analysis

All patients will be included in the analysis

14.7. Procedures for Reporting any Deviation(s) from the Original Statistical Plan

All the statistical analysis procedures, all the possible deviations, and all the reporting operations will be performed according to the Statistical Principles for Clinical Trials (E9, ICH Harmonised Tripartite Guideline).

The statistical report will contain the log of any of the operation conducted on the database, plus, possible comments (preceded by a “%” symbol) if and when needed. Any deviation from the original statistical plan will be therefore reported in the log, together with the reason behind the deviation.

15. DATA MANAGEMENT

15.1. Source Data

Source documents are the first recorded data and from which participants’ eCRF data are obtained. These include, but are not limited to, hospital records (from which medical history and previous and concurrent medication may be summarised into the eCRF), clinical and office charts, laboratory and pharmacy records, diaries, microfiches, radiographs, and correspondence.

All documents will be stored safely in confidential conditions. On all trial-specific documents, other than the signed consent, the participant will be referred to by the trial participant number/code, not by name.

15.2. Access to Data

Direct access will be granted to authorised representatives from promoter and the regulatory authorities to permit trial-related monitoring, audits and inspections.

15.3. Data Recording and Record Keeping

All trial data will be entered on an electronic CRFs.

The participants will be identified by a unique trial specific number and/or code in any database. The name and any other identifying detail will NOT be included in any trial data electronic file.

16. QUALITY ASSURANCE PROCEDURES

The trial will be conducted in accordance to the current approved protocol, ICH GCP, relevant regulations and standard operating procedures.

17. SERIOUS BREACHES

The Medicines for Human Use (Clinical Trials) Regulations contain a requirement for the notification of "serious breaches" to the MHRA within 7 days of the Promoter becoming aware of the breach.

A serious breach is defined as "A breach of GCP or the trial protocol which is likely to affect to a significant degree

- (a) the safety or physical or mental integrity of the subjects of the trial; or
- (b) the scientific value of the trial".

In the event that a serious breach is suspected the Promoter must be contacted within 1 working day.

In collaboration with the C.I., the serious breach will be reviewed by the Sponsor and, if appropriate, the Sponsor will report it to the REC committee, Regulatory authority and the NHS host organisation within seven calendar days.

18. ETHICAL AND REGULATORY CONSIDERATIONS

18.1. Declaration of Helsinki

The Investigator will ensure that this trial is conducted in accordance with the principles of the Declaration of Helsinki.

18.2. Guidelines for Good Clinical Practice

The Investigator will ensure that this study is conducted in accordance with relevant regulations and with Good Clinical Practice.

18.3. Approvals

The protocol, informed consent form, participant information sheet and any proposed advertising material will be submitted to an appropriate Research Ethics Committee (REC) and regulatory authorities for written approval.

The Investigator will submit and, where necessary, obtain approval from the above parties for all substantial amendments to the original approved documents.

18.4. Reporting

The CI shall submit once a year throughout the clinical trial, or on request, an Annual Progress Report to the REC. In addition, an End of Trial notification and final report will be submitted to the REC.

18.5. Participant Confidentiality

The trial staff will ensure that the participants' anonymity is maintained. The participants will be identified only by initials and a participant ID number on the eCRF and any electronic database. All documents will be stored securely and only accessible by trial staff and authorised personnel. The trial will comply with the Data Protection Act, which requires data to be anonymised as soon as it is practical to do so.

18.6. Expenses and Benefits

They will not be intended payments to participants and any other benefits.

18.7. Other Ethical Considerations

Nothing to declare

19. FINANCE AND INSURANCE

19.1. Funding

None.

19.2. Insurance

The study compares two accepted treatment modalities of patients with COVID-19. As per DM 17.12.2004, art 2, comma 4, patients will be covered by hospital insurance. ASST Grande Ospedale Metropolitano will guarantee for patients' insurance.

20. PUBLICATION POLICY

The final manuscript of the study will be prepared by the Principal Investigators. The data will remain property of the Promoters.

21. REFERENCES

1. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020.
2. Fan BE, Chong VCL, Chan SSW, Lim GH, Lim KGE, Tan GB, Mucheli SS, Kuperan P, Ong KH. Hematologic parameters in patients with COVID-19 infection. *Am J Hematol*. 2020 Mar 4.
3. Tang N, Li D, Wang X, Sun Z. Abnormal coagulation parameters are associated with poor prognosis in patients with novel coronavirus pneumonia. *J Thromb Haemost*. 2020 Feb 19.
4. Driggin E, Madhavan MV, Bikdeli B, Chuich T, Laracy J, Bondi-Zoccai G, Brown TS, Nigoghossian C, Zidar DA, Haythe J, Brodie D, Beckman JA, Kirtane AJ, Stone GW, Krumholz HM, Parikh SA. Cardiovascular Considerations for Patients, Health Care Workers, and Health Systems During the Coronavirus Disease 2019 (COVID-19) Pandemic. *J Am Coll Cardiol*. 2020 Mar 18. pii: S0735-1097(20)34637-4.
5. Luo W, Yu H, Gou J, et al. Clinical Pathology of Critical Patient with Novel Coronavirus Pneumonia (COVID-19). *Preprints*. 2020, 2020020407.
6. Darzi AJ, Karam SG, Charide R, Etxeandia Ikobaltzeta I, Cushman M, Gould MK, Mbuagbaw L, Spencer F, Spyropoulos A, Streiff MB, Woller SC, Zakai N, Germini F, Rigoni M, Agarwal A, Morsi RZ, Iorio A, Akl E, Schunemann HJ. Prognostic factors for VTE and Bleeding in Hospitalized Medical Patients: a systematic review and meta-analysis. *Blood*. 2020 Feb 24. pii: blood.2019003603.
7. Mehran R, Rao SV, Bhatt DL, et al. Standardized Bleeding Definitions for Cardiovascular Clinical Trials. A Consensus Report from the Bleeding Academic Research Consortium. *Circulation*. 2011; 123:2736-2747.

LAST PROTOCOL VERSION (VERSION 4.0)

Enoxaparin for thromboprophylaxis in hospitalized COVID-19 patients: comparison of 40 mg o.d. versus 40 mg b.i.d. A randomized Clinical Trial

Internal Reference Number / Short title: X-COVID 19

Ethics Ref:

Date and Version No: September 2020, Version 4.0

Chief Investigators:	Marco Cattaneo, Dipartimento di Scienze della Salute, Università Degli Studi di Milano, Milan, Italy; Unità di Medicina 2, ASST Santi Paolo e Carlo, Milan, Italy; Nuccia Morici, ASST Grande Ospedale Metropolitano Niguarda, Milan, Italy.
Italian Investigators:	Paolo Tarsia, Claudia Marini, Massimo Puoti, Luca Bonacchini (ASST Grande Ospedale Metropolitano Niguarda, Milan, Italy), Paolo Righini (IRCCS Policlinico San Donato, Milan, Italy), Gaetano Maria De Ferrari, Fabrizio D'Ascenzo (AOU Città della Salute e della Scienza, Presidio Molinette, Torino, Italy), Sergio Leonardi, Claudio Montalto (IRCCS Policlinico San Matteo, Pavia, Italy), Paolo Bonfanti (ASST Monza – Ospedale di Monza, Monza, Italy), Ilaria Battistoni, Matteo Francioni (AOU Ospedali Riuniti di Ancona, Ancona, Italy, Annamaria Municinò, Alessandro Rollero (ASL 3 Genovese, Genova, Italy), Giulio Stefanini, Antonio Voza (IRCCS Istituto Clinico Humanitas, Rozzano, Milan, Italy), Stefano Savonitto (ASST Lecco, Ospedale A. Manzoni, Lecco, Italy), Italo Porto (IRCCS Ospedale Policlinico San Martino, Genova), Francesca Cortellaro (Ospedale San Carlo Borromeo, ASST Santi Paolo e Carlo, Milano, Italy), Donato Bettega (Ospedale Sacra Famiglia, Provincia Lombardo-Veneta Fatebenefratelli, Erba, Como, Italy), Ruggero Giuliani (Casa Circondariale San Vittore, ASST-Santi Paolo e Carlo, Milano), Daniele Grosseto (Ospedale Infermi, Azienda USL Romagna, Rimini, Italy); Roberto Pedretti, Maria Francesca Barmina (IRCCS Multimedica, Sesto San Giovanni, Milano); Francesco Saverio Serino (ASL 4 Veneto, Covid Hospital, Jesolo, Italy), Massimo Antonio Di Pietro, Michele Trezzi (AUSL Toscana Centro, Ospedale San Jacopo).
Russian Investigator	Makatsariya Alexander (I.M. Sechenov First Moscow State Medical University), Nikita Lomakin (Central Clinical Hospital of the Presidential Department of Russian Federation)
Sponsor:	ASST Grande Ospedale Metropolitano Niguarda, Milan, Italy
Funder:	Italfarmaco
Chief Investigator Signature:	Marco Cattaneo Nuccia Morici
Statistician Signature:	Laura Antolini Maria Grazia Valsecchi

Confidentiality Statement: This document contains confidential information that must not be disclosed to anyone other than the Sponsor, the Investigator Team, HRA, host organisation, and members of the Research Ethics Committee, unless authorised to do so.

TABLE OF CONTENTS

1.	KEY TRIAL CONTACTS	30
2.	SYNOPSIS	31
3.	ABBREVIATIONS	33
4.	BACKGROUND AND RATIONALE	33
5.	OBJECTIVES AND OUTCOME MEASURES	35
6.	TRIAL DESIGN	36
7.	PARTICIPANT IDENTIFICATION	37
7.1	Trial participants	37
7.2	Inclusion Criteria	37
7.3	Exclusion Criteria	38
8	TRIAL PROCEDURES	38
8.1	Recruitment	39
8.2	Informed Consent	39
8.3	Screening and Eligibility Assessment	39
8.4	Randomisation, blinding and code-breaking	40
8.5	Baseline Assessments	40
8.6	Subsequent Visits	40
8.7	Discontinuation/Withdrawal of Participants from Trial Treatment	41
8.8	Definition of End of Trial	42
9.	COMPLIANCE WITH TRIAL TREATMENT	42
10.	ACCOUNTABILITY OF TRIAL TREATMENT	42
11.	CONCOMITANT MEDICATION	42
12.	POST-TRIAL TREATMENT	42
13.	SAFETY REPORTING	42
13.1.	Definitions	42
13.2.	Causality	43
13.3.	Procedures for Recording Adverse Events	43
13.4.	Reporting Procedures for Serious Adverse Events	44
13.5.	Expectedness	44
13.6.	SUSAR Reporting	44
13.7.	Safety Monitoring Committee	44
14.	STATISTICS	45
14.1.	Description of Statistical Methods	45
14.2.	Number of Participants	45
14.3.	Populations Analysis	46
14.4.	Criteria for the Termination of the Trial	46
14.5.	Procedure for Accounting for Missing, Unused, and Spurious Data	46

14.6. Inclusion in Analysis	46	
14.7. Procedures for Reporting any Deviation(s) from the Original Statistical Plan		46
15. DATA MANAGEMENT	47	
15.1. Source Data	47	
15.2. Access to Data	47	
15.3. Data Recording and Record Keeping	47	
16. QUALITY ASSURANCE PROCEDURES	47	
17. SERIOUS BREACHES	47	
18. ETHICAL AND REGULATORY CONSIDERATIONS	48	
18.1. Declaration of Helsinki	48	
18.2. Guidelines for Good Clinical Practice	48	
18.3. Approvals	48	
18.4. Reporting	48	
18.5. Participant Confidentiality	48	
18.6. Expenses and Benefits	49	
19. FINANCE AND INSURANCE	49	
19.1. Funding	49	
19.2. Insurance	49	
20. PUBLICATION POLICY	49	
21. REFERENCES	50	

4. KEY TRIAL CONTACTS

Chief Investigator	<p>Marco Cattaneo, Dipartimento di Scienze della Salute, Università Degli Studi di Milano, Milan, Italy; Unità di Medicina 2, ASST Santi Paolo e Carlo, Milan, Italy. Phone 0250323095 - 0281844638; fax 0250323089; e-mail marco.cattaneo@unimi.it</p> <p>Nuccia Morici, Unità di Cure Intensive Cardiologiche Dipartimento Cardio-toraco-vascolare, ASST Grande Ospedale Metropolitano Niguarda, Milan, Italy. Phone 0264442565, fax 0264442818, e-mail nuccia.morici@ospedaleniguarda.it</p>
Sponsor	ASST Grande Ospedale Metropolitano Niguarda, Milan, Italy
Clinical Trials Unit	Unità di Medicina, ASST Santi Paolo e Carlo, Milan, Italy / Malattie Infettive, ASST Grande Ospedale Metropolitano Niguarda, Milan, Italy
Statisticians	Laura Antolini, e-mail laura.antolini@unimib.it
Committees	<p>Comitato Etico Scientifico INMI Lazzaro Spallanzani (Ethical Committee)</p> <p>Steering Committee: Marco Cattaneo, Nuccia Morici, Enrico Ammirati, Marco Valgimigli, Stefano Savonitto.</p>

5. SYNOPSIS

Trial Title	Enoxaparin for thromboprophylaxis in hospitalized COVID-19 patients: comparison of 40 mg o.d. versus 40 mg b.i.d. A randomized Clinical Trial
Internal ref. no. (or short title)	X-Covid 19
Promoter	ASST Grande Ospedale Metropolitano Niguarda
Principal Investigator	Marco Cattaneo / Nuccia Morici
EudraCT Number	2020-001708-41
Clinical Phase	Phase IIIb
Trial Design	Open-label Multicentre Prospective Controlled Randomized Trial in patients with SARS-CoV-2 infection. Patients will be randomized 1:1 to 40 mg subcutaneous enoxaparin o.d. versus 40 mg enoxaparin b.i.d
Trial Participants <i>Inclusion criteria</i>	<ul style="list-style-type: none"> All-comers patients aged ≥ 18 years and admitted to hospital with laboratory-confirmed SARS-CoV-2 infection.
<i>Exclusion criteria</i>	<ul style="list-style-type: none"> Patients admitted directly to an intensive care unit; Estimated creatinine clearance < 15 ml/min/1.73m²; Patients needing anticoagulant for prior indication; Patients treated with heparin at any increased dose compared to prophylactic regimen before enrolment Patients at high bleeding risk or experiencing clinically significant bleeding; Patients involved in competitive clinical trials exploring antithrombotic treatments; Any other significant disease or disorder which, in the opinion of the investigator, may either put the participants at risk because of participation in the trial, or may influence the result of the trial, or the participant's ability to participate in the trial.
Planned Sample Size	<p>A sample of 2712 subjects is required to test the superiority hypothesis with 80% power according to a two-sided 0.05 level of significance that 40 mg enoxaparin b.i.d. will decrease the primary endpoint compared to 40 mg enoxaparin o.d. An event rate of 10% was hypothesized in the control arm and 7% in the treatment arm, corresponding to a 33% relative risk reduction.</p> <p>The trial will be designed with one interim analysis. This means that a preliminary test to potentially stop the trial for efficacy is performed after 492 out of 2712 planned subjects (246 randomized to enoxaparin 40 mg od and 246 randomized to enoxaparin 40 mg bid) have completed the trial. If the trial is not stopped early, a final test is performed once 100 percent of the planned subjects will be enrolled. For the interim analysis the proportion for the primary endpoint is estimated to be 0.07 in the half-dose anticoagulant group and 0.17 in the standard group. With a type I error α set to 0.05 and a power $1 - \beta$ set to 0.8, a total of 492 subjects will be required.</p>
Treatment duration	In-hospital (at least 30 days of therapy)

Follow up duration	30 days from the enrolment
Planned Trial Period	13 months (Estimated end date May 2021)
Primary Objective	To compare the effects of 40 mg subcutaneous enoxaparin o.d. versus 40 mg enoxaparin b.i.d on the incidence of venous thromboembolism (VTE) [a composite of incident asymptomatic and symptomatic proximal deep vein thrombosis (DVT) diagnosed by serial compression ultrasonography (CUS), and symptomatic pulmonary embolism (PE) diagnosed by CT scan], in patients with SARS-CoV-2 infection.
Secondary Objectives	<ul style="list-style-type: none"> • To compare the effects of 40 mg subcutaneous enoxaparin o.d. versus 40 mg enoxaparin b.i.d on the incidence of in-hospital major complications, defined as the composite of death, VTE, use of mechanical ventilation, stroke, acute myocardial infarction and admission to an intensive care in patients with SARS-CoV-2. • To compare each single component of the primary endpoint between the two groups. • To compare maximum sequential organ failure assessment (SOFA) score between the two groups. • To compare C-reactive protein, D-dimer, IL-6 and hs-troponin levels (as % above the upper reference limit [URL]) among the two groups. • To compare the incidence of SARS-CoV-2-related Acute Respiratory Distress Syndrome (ARDS) between the two groups. • To compare length of hospital stay between the two groups. • To compare measures of right ventricular function at trans-thoracic echocardiography or CT between admission and follow-up, whenever available. • To compare the incidence of 30 days adverse events, defined as composite of death, stroke and myocardial infarction
Safety Endpoints	<ul style="list-style-type: none"> • To compare the incidence of bleeding events according to the <i>ISTH bleeding scale</i> between the two groups. • To compare the incidence of bleeding events according to <i>BARC classification</i> between the two groups. Only BARC 3 and 5 will be considered for secondary endpoint. • To assess the incidence of heparin-induced thrombocytopenia in the two groups.

6. ABBREVIATIONS

AE	Adverse event
AR	Adverse reaction
BARC	Bleeding Academic Research Consortium
CI	Chief Investigator
CRF	Case Report Form
CT	Clinical Trials
CTA	Clinical Trials Authorisation
CTRG	Clinical Trials and Research Governance
DMC/DMSC	Data Monitoring Committee / Data Monitoring and Safety Committee
DSUR	Development Safety Update Report
GCP	Good Clinical Practice
GP	General Practitioner
IB	Investigator's Brochure
ICF	Informed Consent Form
ICH	International Conference on Harmonisation
IMP	Investigational Medicinal Product
IRB	Independent Review Board
PI	Principal Investigator
PIL	Participant/ Patient Information Leaflet
REC	Research Ethics Committee
SAE	Serious Adverse Event
SAR	Serious Adverse Reaction
SDV	Source Data Verification
SMPC	Summary of Medicinal Product Characteristics
SOP	Standard Operating Procedure
SOFA	Sequential Organ Failure Assessment
SUSAR	Suspected Unexpected Serious Adverse Reactions
TMF	Trial Master File

8. BACKGROUND AND RATIONALE

Critically ill medical patients are at heightened risk for venous thromboembolism (VTE) and, for this reason, they are treated with prophylactic doses of low molecular weight heparin (LMWH) (for instance, 40 mg enoxaparin). Patients with COVID-19, which is caused by SARS-CoV-2 infection, are apparently at particularly high risk of VTE, as a consequence of excessive activation of the hemostatic system which, in the most severe cases, could also be associated with the formation of microthrombi and clinically relevant disseminated intravascular coagulation.¹⁻⁴

Based on personal communications by medical doctors working mostly in Northern Italy, there are concerns about the efficacy of thromboprophylaxis with low-molecular-weight heparin (LMWH) at the standard doses in these patients, given their apparently very high risk of thromboembolic complications. Therefore, in many Northern Italian hospitals these patients are given LMWH at much higher doses than those recommended for thromboprophylaxis: in some hospitals, full anticoagulant doses (100 U/Kg b.i.d.) are being used even in the absence of evidence of ongoing thromboembolic disease. This practice does not take into due account the obvious risk of major, potentially fatal bleeding that is associated with high dosing, in the absence of any evidence of higher efficacy of high doses compared to standard prophylactic doses.

An additional rationale for the use of high dose LMWH in many hospitals is that hyper-activation of the hemostatic system in COVID-19 patients could lead to the formation of microthrombi in pulmonary vessels⁵, causing further respiratory distress, and in vessels of other vital organs, causing multi-organ failure (MOF). However, there is no demonstration that this therapeutic approach decreases the risk of microangiopathic thrombosis in these patients, as well as, for that matter, in other types of patients.

Given this scenario, we designed a randomized trial comparing standard prophylactic dose of subcutaneous enoxaparin (40 mg o.d.) with higher dose (40 mg b.i.d) with the aim of testing whether high-dose thromboprophylaxis is more effective than standard dose in preventing TEV in COVID-19 patients. This study would also allow us to test whether double-dose prophylaxis with enoxaparin would favourably affect the natural history of the disease, by decreasing the incidence of organ dysfunction associated with the formation of microthrombi in vital organs.

9. OBJECTIVES AND OUTCOME MEASURES

Objectives	Outcome measures	Timepoint(s) of evaluation of these outcome measure (if applicable)
<p>Primary Objective</p> <ul style="list-style-type: none"> To compare the incidence of hospital-associated venous thromboembolism (HA-VTE) between the two groups (40 mg o.d. vs 40 mg b.i.d subcutaneous enoxaparin) 	<ul style="list-style-type: none"> <i>Number (percentage) of patients who will experience</i> the composite event. <i>Relative risk</i> will evaluate the association between the study outcome and treatment. 	Discharge
<p>Secondary Objectives</p> <ul style="list-style-type: none"> To compare the incidence of in-hospital major complications, defined as the composite of death, HA-VTE, use of mechanical ventilation, major stroke, acute myocardial infarction and admission to an intensive care between the two groups. To compare each single component of the primary endpoint between the two groups. To compare maximum sequential organ failure assessment (SOFA) score between the two groups. To compare C-reactive protein, D-dimer, IL-6 and hs-troponin levels (as % above the upper reference limit [URL]) among the two groups. To compare the incidence of SARS-CoV-2- related Acute Respiratory Distress Syndrome (ARDS) between the two groups. 	<ul style="list-style-type: none"> <i>Number (percentage) of patients who will experience</i> the event. <i>Relative risk</i> will evaluate the association between the study outcome and treatment. <i>Number (percentage) of patients who will experience</i> the event. <i>Relative risk</i> will evaluate the association between the study outcome and treatment. Maximum sequential organ failure assessment (SOFA) score summarized by <i>median and interquartile range (IQR) or mean and standard deviation</i> according to variable distribution. Laboratory parameters will be indicated as <i>median and interquartile range (IQR) or mean and standard deviation</i> according to variable distribution. <i>Number (percentage) of patients who will experience</i> the event. <i>Relative risk</i> will evaluate the association between the study outcome and treatment. 	<p>Discharge</p> <p>Discharge</p> <p>Discharge</p> <p>Discharge</p> <p>Discharge</p> <p>Discharge</p> <p>Discharge</p> <p>30 day</p>

<ul style="list-style-type: none"> • To compare length of hospital stay • To compare measures of right ventricular function at trans-thoracic echocardiography or CT between admission and follow-up, whenever available. • To compare the incidence of 30 days adverse events, defined as composite of death, stroke and myocardial infarction 	<ul style="list-style-type: none"> • Median and interquartile range (IQR) or mean and standard deviation according to variable distribution. • Echocardiographic parameters will be summarized by median and interquartile range (IQR) or mean and standard deviation according to variable distribution. • Number (percentage) of patients who will experience the event. Relative risk will evaluate the association between the study outcome and treatment. 	
<p>Safety endpoints</p> <ul style="list-style-type: none"> • To assess the occurrence of bleeding events, according to the two classifications selected • To assess the incidence of heparin-induced thrombocytopenia between the two groups. 	<ul style="list-style-type: none"> • Number (percentage) of patients who will experience the event. Relative risk will evaluate the association between the study outcome and treatment. • The incidence rate will be described across groups 	<p>During hospital stay</p> <p>During hospital stay</p>

10. TRIAL DESIGN

Prospective controlled randomized, open-label multi-center trial.

Patients with COVID-19 (any hospitalized patients) with laboratory-confirmed SARS-CoV-2 infection will be randomized 1:1 to 40 mg enoxaparin/day subcutaneous versus 40 mg enoxaparin b.i.d. Standard dosage of Proton Pump Inhibitor (PPI) will be prescribed to each enrolled patient to reduce gastrointestinal (GI) bleeding risk. The chronic kidney disease epidemiology (CKD-EPI) formula will be used for eGFR calculation. Enoxaparin treatment will continue until discharge. The study conduct will not change the usual clinical practice in terms of treatments and diagnostic procedures.

Patients will be monitored every 3 days with D-dimer, fibrinogen, complete blood count, lactate dehydrogenase (LDH), PT, APTT, C-reactive protein, procalcitonin only in the case of superinfection by a different agent, ferritin, creatine phosphokinase (CK), CK-MB, albumin, serum creatinine,

aspartate transaminase (AST), alanine transaminase (ALT), high-sensitivity troponin, serum bilirubin and interleukine-6 along with other examinations according to clinical practice.

Patients will be monitored every seven days from the enrolment with CUS of the lower limbs for proximal DVT screening. For patients who display signs and symptoms of DVT, CUS will be performed as soon as possible in the same day. Proximal DVT, namely DVT of iliac, femoral and popliteal veins down to popliteal trifurcation, is defined by cross-sectional vein incompressibility or direct thrombus imaging with vein enlargement. Distal DVT might be reported if distal segments are explored.

For patients who display signs and symptoms of pulmonary embolism, a CT scan will be performed as soon as possible in the same day according to the indication of the attending physician.

Multidetector CT pulmonary angiography is the method of choice for imaging the pulmonary vasculature in patients with suspected pulmonary embolism, that is defined by contrast-filling defects.

With regard to secondary objective, the following definitions apply.

All-cause mortality includes all deaths, regardless of whether the cause of death is determined.

Cardiovascular death is defined as death due to atherosclerotic coronary heart disease, cerebrovascular accident or (complication of) peripheral embolization, and includes deaths due to acute MI, stroke, sudden death, non-sudden death, unwitnessed death, and procedure-related deaths. Non-cardiovascular death includes deaths due to all other causes.

Myocardial Infarction denotes the presence of acute myocardial injury detected by abnormal cardiac biomarkers [Cardiac troponin I (cTnI) and T (cTnT) increased above the 99th percentile upper reference limit (URL)] in the setting of evidence of acute myocardial ischaemia.

Stroke is defined as any new neurological deficit lasting >24 hours. CT or MRI are required to confirm the stroke and to distinguish between ischemic and hemorrhagic stroke. Strokes will be also classified as fatal or nonfatal: nonfatal stroke will be also classified as disabling and non disabling.

The total duration of the study is 13 months for enrolment, follow-up, data collection, cleaning and analysis.

11. PARTICIPANT IDENTIFICATION

8.9 Trial participants

Patients directly admitted to hospital for suspected SARS-CoV-2 infection will be screened for randomization.

8.10 Inclusion Criteria

All the following inclusion criteria will be adopted:

All-comers patients aged ≥ 18 years and admitted to hospital with laboratory-confirmed SARS-CoV-2 infection.

8.11 Exclusion Criteria

The following patients will not be enrolled:

- Patients admitted directly to intensive care unit.
- Patients with estimated creatinine clearance $<15 \text{ ml/min/1.73m}^2$.
- Patients needing anticoagulant for prior indication.
- Patients treated with heparin at any increased dose compared to prophylactic regimen before enrolment
- Patients at high bleeding risk or experiencing clinically significant bleeding;
- Patients involved in competitive clinical trials exploring antithrombotic treatments;
- Patients with any other significant disease or disorder which, in the opinion of the Investigator, may either put the participants at risk because of participation in the trial, or may influence the result of the trial, or the participant's ability to participate in the trial.

9 TRIAL PROCEDURES

Eligibility will be evaluated by the attending physician after the first visit and after confirmation of the diagnosis. The patient, properly informed about the study, will be able to decide to sign the informed consent and the privacy form. The physician will obtain a detailed anamnesis and he/she will carefully check the clinical parameters needed to define tissue perfusion (as detailed by the inclusion criteria) along with body temperature. The following laboratory data will be checked: haemochrome, renal function, serum electrolytes, liver function, coagulation, blood gas analysis. Laboratory examinations will be sequentially evaluated according to the flow chart timeline (section 8.6). If the patient develops a condition needing an anticoagulation, an increase in enoxaparin dosage to 100 UI/kg (1 mg/kg) b.i.d. might be considered if indicated and in the absence of contraindications. Enoxaparin treatments will continue until discharge. Serial CUS will be performed, the first on admission and then every seven days. Bleeding events will be registered and graded according to the International Society on Thrombosis and Haemostasis (ISTH) bleeding scale and Bleeding Academic Research Consortium (BARC) scale. Also BARC 2 bleedings will be recorded. Pre-specified subgroup analyses will be performed for 1) D-dimer categories (>1 <2 , 2, 3, 4, 5, 6 fold the ULN of the reference range); 2) estimated creatinine clearance ($<$ versus $>60 \text{ ml/min/1.73 m}^2$); 3) severity of the respiratory symptoms (mild versus moderate-severe defined as respiratory rate ≥ 30 breaths/min; arterial oxygen saturation $\leq 93\%$ at rest; $\text{PaO}_2/\text{FiO}_2 \leq 300 \text{ mmHg}$); 4) concomitant antiplatelet therapy.

The identification of adverse reactions will be continuous during hospitalisation. The data will be recorded in an eCRF.

9.1 Recruitment

Patients recruitment in Italy will be performed at the Cardiology Department, Malattie Infettive and Internal Medicine, ASST Grande Ospedale Metropolitano Niguarda, Milan, Italy; Medicina, Ospedale San Paolo, ASST Santi Paolo e Carlo, Milan, Italy; IRCCS Policlinico San Donato, Milan, Italy; AOU Città della Salute e della Scienza, Presidio Molinette, Torino, Italy; IRCCS Policlinico San Matteo, Pavia, Italy; ASST Monza – Ospedale di Monza, Monza, Italy; AOU Ospedali Riuniti di Ancona, Ancona, Italy; ASL 3 Genovese, Genova, Italy; IRCCS Istituto Clinico Humanitas, Rozzano, Milan, Italy; ASST Lecco, Ospedale A. Manzoni, Lecco, Italy; IRCCS Ospedale Policlinico San Martino, Genova, Italy; Ospedale San Carlo Borromeo, ASST Santi Paolo e Carlo, Milan, Italy; Ospedale Sacra Famiglia, Provincia Lombardo-Veneta Fatebenefratelli, Erba, Como, Italy; Casa Circondariale San Vittore, ASST-Santi Paolo e Carlo, Milano, Italy; Ospedale Infermi, Azienda USL Romagna, Rimini, Italy; IRCCS Multimedica, Sesto San Giovanni (Milano), Italy; ASL 4 Veneto, COVID Hospital, Jesolo (Venezia); AUSL Toscana Centro, Ospedale San Jacopo, Pistoia.

Patients recruitment in Russia will be performed at the I.M. Sechenov First Moscow State Medical University, Moscow, Russia; Central Clinical Hospital of the Presidential Department of Russian Federation, Moscow, Russia

9.2 Informed Consent

The participant must personally sign and date the latest approved version of the Informed Consent form before any trial specific procedures.

Written and verbal versions of the Participant Information and Informed Consent will be presented to the participants detailing no less than: the exact nature of the trial; what it will involve the participant; the implications and constraints of the protocol; the known side effects and any risks involved in taking part. It will be clearly stated that the participant is free to withdraw from the trial at any time for any reason without prejudice to future care and with no obligation to give the reason for withdrawal.

The participant will be allowed as much time as wished, taking into consideration the urgency of treatments, to consider the information and the opportunity to ask the Investigator, his/her GP or other independent parties to decide whether he/she will participate in the trial. Written Informed Consent will then be obtained by means of participant dated signature and dated signature of the person who presented and obtained the Informed Consent. The person who obtained the consent must be suitably qualified and experienced, and authorized to do so by the Chief/Principal Investigator. The original signed Informed Consent will be given to the participant and a copy will be retained at the trial site.

9.3 Screening and Eligibility Assessment

As clinically indicated, the screening procedures will be performed throughout demographics, medical history, concomitant medication, physical examination, laboratory tests and radiological findings (chest CT scans are encouraged over chest X-rays).

The screening procedure will be immediately followed by recruitment at arrival in the ward (patients will not be randomized in the emergency department), if allowed on basal evaluation and patient informed consent signature.

9.4 Randomisation, blinding and code-breaking

This study will be a two-parallel group randomized, open-label, multi-center trial in hospitalized COVID-19 patients. Patients will be randomized by the attending physicians using an IVRS that randomizes at a ratio of 1:1 upon determination of eligibility.

A total sample of 2712 subjects is required to test the superiority hypothesis with 80% power according to a two-sided 0.5 level of significance that 40 mg enoxaparin subcutaneously bid/day (treatment arm) will decrease the primary endpoint compared to 40 mg enoxaparin/day (prophylaxis dose, that is considered the control arm as represents the standard of care for hospitalized patients that are bedridden). An absolute reduction of 3% is foreseen in the treatment arm (enoxaparin 40 mg b.i.d); an event rate of 10% was considered in the control arm (enoxaparin 40 mg o.d.) and 7% in the treatment arm.

The duration of the enrolment is 10 months from the obtaining of the approvals. The total duration of the study is 13 months for follow-up (1 months), data collection, cleaning and analysis (2 months). Randomisation will be done at the same visit as the baseline visit. There will not be a run-in period.

9.5 Baseline Assessments

Basal evaluation (V1)

Every patient will be informed about the objectives of our study and he/she will be asked for consent to the trial. At basal visit and following ones, the physician will perform: physical examination, hemodynamic parameters and blood gas analysis check; CUS. Echocardiogram might be performed if requested by the attending physician.

The occurrence of organ dysfunction and failure, from admission through days spent in intensive care, will be assessed by the Sequential Organ Failure Assessment (SOFA) score.

9.6 Subsequent Visits

After basal assessment, the following visits will be performed according to local clinical practice a check-up at discharge will be performed. Echocardiograms might be performed during hospitalization

according to the indication of the attending physician. In case mechanical ventilation is required, ventilation modes and eventual variations of parameters will be recorded.

If subject is discharged before 30 days from enrolment, a telephonic follow up will be performed at 30 days to assess potential adverse events occurred during that period. A composite of overall mortality, myocardial infarction and stroke will be considered.

The schedule of clinical visits and assessment procedures is shown in the following flow chart.

Flow chart

Visit	V1	V2	V2	V3	V4
Time interval	Basal	To be checked each three days	To be checked each seven days	Discharge	30-day from the enrolment
Informed consent	X				
Time from onset symptoms to enrolment	X				
Physical examination	X				
Height	X				
Weight	X				
Anamnesis	X				X
BP (s/d/m)	X				
HR	X	X		X	
CUS	X		X		
Hb	X	X		X	
Creatinine	X	X		X	
Platelets	X	X		X	
Leucocytes	X	X		X	
INR	X	X		X	
Bilirubin	X	X		X	
AST	X	X		X	
ALT	X	X		X	
LDH	X	X		X	
CRP	X	X		X	
Troponin (HS)	X	X		X	
IL-6	X	X		X	
D-dimer	X	X		X	
Fibrinogen	X	X		X	
SOFA score	X	X			

9.7 Discontinuation/Withdrawal of Participants from Trial Treatment

Each participant has the right to withdraw from the trial at any time. In addition, the Investigator may discontinue a participant from the trial at any time if the Investigator considers it necessary for any reason including:

- Ineligibility (either arising during the trial or retrospectively having been overlooked at screening)
- Significant protocol deviation
- Significant non-compliance with treatment regimen or trial requirements
- An adverse event which requires discontinuation of trial treatment or results in inability to continue to comply with trial procedures
- Disease progression which requires discontinuation of trial treatment or results in inability to continue to comply with trial procedures
- Withdrawal of Consent

Withdrawal from the trial will result in exclusion of the data for that participant from analysis

The reason for withdrawal will be recorded in the eCRF.

If the participant is withdrawn due to an adverse event, the Investigator will arrange for follow-up visits or telephone calls until the adverse event has resolved or stabilised.

9.8 Definition of End of Trial

The end of trial is the date of the last visit/ telephone follow up/ home visit of the last participant.

22. COMPLIANCE WITH TRIAL TREATMENT

All patients will be treated in hospital ward, so compliance will be guaranteed by the investigators.

23. ACCOUNTABILITY OF TRIAL TREATMENT

The study does will involve drugs usually administrated in clinical practice.

24. CONCOMITANT MEDICATION

There are not any contraindicated medications. All used drugs, as per clinical indication, will be reported in the eCRF.

25. POST-TRIAL TREATMENT

At the end of the study period, all surviving patients will be treated as per current clinical practice in the participating Centers.

26. SAFETY REPORTING

26.1. Definitions

Adverse Event (AE)	Any untoward medical occurrence in a participant to whom a medicinal product or diagnostic-interventional procedure has been
--------------------	--

	administered, including occurrences which are not necessarily caused by or related to that product or procedure.
Adverse Reaction (AR)	<p>An untoward and unintended response in a participant to a medicinal product which is related to any dose administered to that participant.</p> <p>The phrase "response to an investigational medicinal product" means that a causal relationship between a trial medication and an AE is at least a reasonable possibility, i.e. the relationship cannot be ruled out.</p> <p>All cases judged by either the reporting medically qualified professional or the Sponsor as having a reasonable suspected causal relationship to the trial medication qualify as adverse reactions.</p>
Serious Adverse Event (SAE)	<p>A serious adverse event is any untoward medical occurrence that:</p> <ul style="list-style-type: none"> • Results in death • Is life-threatening • Requires inpatient hospitalisation or prolongation of existing hospitalisation • Results in persistent or significant disability/incapacity • Consists of a congenital anomaly or birth defect. <p>Other 'important medical events' may also be considered serious if they jeopardise the participant or require an intervention to prevent one of the above consequences.</p> <p>NOTE: The term "life-threatening" in the definition of "serious" refers to an event in which the participant was at risk of death at the time of the event; it does not refer to an event which hypothetically might have caused death if it were more severe.</p>

NB: to avoid confusion or misunderstanding of the difference between the terms "serious" and "severe", the following note of clarification is provided: "Severe" is often used to describe intensity of a specific event, which may be of relatively minor medical significance. "Seriousness" is the regulatory definition supplied above.

26.2. Causality

The relationship of each adverse event to trial treatment must be determined by a medically qualified individual according to the following definitions:

Related: The adverse event follows a reasonable temporal sequence from trial treatment start. It cannot reasonably be attributed to any other cause.

Not Related: The adverse event is probably produced by the participant's clinical state or by other modes of therapy administered to the participant.

26.3. Procedures for Recording Adverse Events

All AEs occurring during the trial / or until the following two months that are observed by the Investigator or reported by the participant, will be recorded on the eCRF, whether or not attributed to trial treatment.

The following information will be recorded: description, date of onset and end date, severity, assessment of relatedness to trial treatment, other suspect drug or device and action taken. Follow-up information should be provided as necessary.

The severity of events will be assessed on the following scale: 1 = mild, 2 = moderate, 3 = severe. AEs considered related to the trial treatment as judged by a medically qualified investigator will be followed either until resolution, or the event is considered stable.

It will be left to the Investigator's clinical judgment to decide whether an AE is of enough severity to require the participant's removal from treatment. A participant may also voluntarily withdraw from treatment due to what he or she perceives as an intolerable AE. If either of these occurs, the participant must undergo an end of trial assessment and be given appropriate care under medical supervision until symptoms cease, or the condition becomes stable.

26.4. Reporting Procedures for Serious Adverse Events

All SAEs (other than those defined in the protocol as not requiring reporting) must be reported on the SAE reporting form to Eudravigilance within 24 hours of the Site Study Team becoming aware of the event.

Review of SAEs must be timely, taking into account the reporting time for a potential SUSAR.

26.5. Expectedness

Expectedness will be determined according to the Investigators' Brochure/Summary of Product Characteristics.

26.6. SUSAR Reporting

All SUSARs will be reported by the CI to the relevant Competent Authority and to the REC and other parties as applicable. For fatal and life-threatening SUSARS, this will be done no later than 7 calendar days after the investigator team is first aware of the reaction. Any additional information will be reported within 15 calendar days. Treatment codes will be un-blinded for specific participants.

26.7. Safety Monitoring Committee

The aims of this committee include:

- To pick up any trends, such as increases in un/expected events, and take appropriate action
- To seek additional advice or information from investigators where required
- To evaluate the risk of the trial continuing and take appropriate action where necessary

27. STATISTICS

27.1. Description of Statistical Methods

All protocol-defined endpoints will be measured from the time of randomization. Descriptive statistics of baseline characteristics of patients randomized to the two groups will be reported. Continuous variables will be summarized by means, standard deviations, medians, inter-quartile ranges and minimum and maximum values. Categorical variables will be summarized by frequencies and percentages.

Data will be analysed according to the intention-to-treat principle. In addition, a per-protocol analysis of the primary endpoint, considering patients without major protocol violations (i.e. patients who crossed over), will be performed to evaluate the robustness of the data.

A test for difference of proportions will be carried out to compare in-hospital composite events among patients randomized to the two groups. A relative risk will be calculated to quantify the magnitude of the half anticoagulant effect.

For what regards continuous secondary endpoints, maximum SOFA score will be compared among the study groups through the t-test or Wilcoxon rank-sum test, according to the distribution of study endpoints. The same approach will be applied for laboratory examinations and times included in the secondary endpoints.

All tests will be two-sided and a p-value of 0.05 or less will be considered statistically significant. No adjustments for multiple testing will be applied in the analysis of secondary endpoints and in the subgroup analysis, therefore, inferences drawn from the reported p-values and 95% CIs will be considered exploratory. If a p-value is less than 0.001 it will be reported as “<0.001”.

27.2. Number of Participants

Summary Statements

The proportion for the primary endpoint is estimated to be 0.07 in the half-dose anticoagulant dose and 0.10 in the standard group. With a type I error α set to 0.05 and a power $1 - \beta$ set to 0.8, a total of 2712 subjects will be enrolled in the trial. The trial will be designed with one interim analysis. This means that a preliminary test to potentially stop the trial for efficacy is performed after 492 out of 2712 planned subjects (246 randomized to enoxaparine 40 mg od and 246 randomized to enoxaparine 40 mg bid) have completed the trial. If the trial is not stopped early, a final test is performed once 100 percent of the planned subjects will be enrolled. For the interim analysis the proportion for the primary endpoint is estimated to be 0.07 in the half-dose anticoagulant group and 0.17 in the standard group. With a type I error α set to 0.05 and a power $1 - \beta$ set to 0.8, a total of 492 subjects will be required.

27.3. Populations Analysis

For this trial, the following populations will be defined and used in the analysis and/or presentation of the data.

Safety population: The safety population will be classified according to the actual treatment on enoxaparin 40 mg b.i.d. or control, received.

Intent-to-treat (ITT) population: Treatment classification for ITT analysis will be based on the randomized treatment.

Per-Protocol (PP) population: The PP population will be defined as all ITT patients who received assigned to enoxaparin 40 mg b.i.d. and will be without major protocol violations. Specifically, subjects may be excluded from PP population if any of the following criteria are met:

- Inclusion/Exclusion criteria violations;
-

The PP definition for each patient will be finalized before database lock and study unblinding.

The primary and secondary efficacy analyses will be based on the ITT population. Analyses based on the PP population will be considered supportive.

27.4. Criteria for the Termination of the Trial

Criteria refers to the above reported statistical plan (section 14.2).

27.5. Procedure for Accounting for Missing, Unused, and Spurious Data

All data will be checked for consistency and plausibility based on the nature of each collected variable before any statistical analysis. Data falling beyond limits of plausibility will be checked in the medical records and in the CRF to correct for possible typos or transcription errors. Changes in the database will be recorded together with date and name of the physician performing that correction and with the reason for correction.

Methods of multiple imputation will be considered if the missingness rate will be more than 20%.

27.6. Inclusion in Analysis

All patients will be included in the analysis

27.7. Procedures for Reporting any Deviation(s) from the Original Statistical Plan

All the statistical analysis procedures, all the possible deviations, and all the reporting operations will be performed according to the Statistical Principles for Clinical Trials (E9, ICH Harmonised Tripartite Guideline).

The statistical report will contain the log of any of the operation conducted on the database, plus, possible comments (preceded by a “%” symbol) if and when needed. Any deviation from the original statistical plan will be therefore reported in the log, together with the reason behind the deviation.

28. DATA MANAGEMENT

28.1. Source Data

Source documents are the first recorded data and from which participants’ eCRF data are obtained. These include, but are not limited to, hospital records (from which medical history and previous and concurrent medication may be summarised into the eCRF), clinical and office charts, laboratory and pharmacy records, diaries, microfiches, radiographs, and correspondence.

All documents will be stored safely in confidential conditions. On all trial-specific documents, other than the signed consent, the participant will be referred to by the trial participant number/code, not by name.

28.2. Access to Data

Direct access will be granted to authorised representatives from promoter and the regulatory authorities to permit trial-related monitoring, audits and inspections.

28.3. Data Recording and Record Keeping

All trial data will be entered on an electronic CRFs.

The participants will be identified by a unique trial specific number and/or code in any database. The name and any other identifying detail will NOT be included in any trial data electronic file.

29. QUALITY ASSURANCE PROCEDURES

The trial will be conducted in accordance to the current approved protocol, ICH GCP, relevant regulations and standard operating procedures.

30. SERIOUS BREACHES

The Medicines for Human Use (Clinical Trials) Regulations contain a requirement for the notification of "serious breaches" to the MHRA within 7 days of the Promoter becoming aware of the breach. A serious breach is defined as "A breach of GCP or the trial protocol which is likely to affect to a significant degree

- (a) the safety or physical or mental integrity of the subjects of the trial; or
- (b) the scientific value of the trial".

In the event that a serious breach is suspected the Promoter must be contacted within 1 working day. In collaboration with the C.I., the serious breach will be reviewed by the Sponsor and, if appropriate, the Sponsor will report it to the REC committee, Regulatory authority and the NHS host organisation within seven calendar days.

31. ETHICAL AND REGULATORY CONSIDERATIONS

31.1. Declaration of Helsinki

The Investigator will ensure that this trial is conducted in accordance with the principles of the Declaration of Helsinki.

31.2. Guidelines for Good Clinical Practice

The Investigator will ensure that this study is conducted in accordance with relevant regulations and with Good Clinical Practice.

31.3. Approvals

The protocol, informed consent form, participant information sheet and any proposed advertising material will be submitted to an appropriate Research Ethics Committee (REC) and regulatory authorities for written approval.

The Investigator will submit and, where necessary, obtain approval from the above parties for all substantial amendments to the original approved documents.

31.4. Reporting

The CI shall submit once a year throughout the clinical trial, or on request, an Annual Progress Report to the REC. In addition, an End of Trial notification and final report will be submitted to the REC.

31.5. Participant Confidentiality

The trial staff will ensure that the participants' anonymity is maintained. The participants will be identified only by initials and a participant ID number on the eCRF and any electronic database. All documents will be stored securely and only accessible by trial staff and authorised personnel. The trial

will comply with the Data Protection Act, which requires data to be anonymised as soon as it is practical to do so.

31.6. Expenses and Benefits

They will not be intended payments to participants and any other benefits.

31.7. Other Ethical Considerations

Nothing to declare

32. FINANCE AND INSURANCE

32.1. Funding

Italfarmaco will provide funding for clinical study.

32.2. Insurance

The study compares two accepted treatment modalities of patients with COVID-19.

As per DL 8 aprile 2020, n. 23, art 40, comma 5, “Disposizioni urgenti materia di sperimentazione dei medicinali per l'emergenza epidemiologica da COVID”, for COVID-related No-Profit studies, no insurance is needed in Italy.

For Russian centers, no insurance is needed per local laws.

33. PUBLICATION POLICY

The final manuscript of the study will be prepared by the Principal Investigators.

The data will remain property of the Promoters.

34. REFERENCES

1. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020.
2. Fan BE, Chong VCL, Chan SSW, Lim GH, Lim KGE, Tan GB, Mucheli SS, Kuperan P, Ong KH. Hematologic parameters in patients with COVID-19 infection. *Am J Hematol*. 2020 Mar 4.
3. Tang N, Li D, Wang X, Sun Z. Abnormal coagulation parameters are associated with poor prognosis in patients with novel coronavirus pneumonia. *J Thromb Haemost*. 2020 Feb 19.
4. Driggin E, Madhavan MV, Bikdeli B, Chuich T, Laracy J, Bondi-Zoccai G, Brown TS, Nigoghossian C, Zidar DA, Haythe J, Brodie D, Beckman JA, Kirtane AJ, Stone GW, Krumholz HM, Parikh SA. Cardiovascular Considerations for Patients, Health Care Workers, and Health Systems During the Coronavirus Disease 2019 (COVID-19) Pandemic. *J Am Coll Cardiol*. 2020 Mar 18. pii: S0735-1097(20)34637-4.
5. Luo W, Yu H, Gou J, et al. Clinical Pathology of Critical Patient with Novel Coronavirus Pneumonia (COVID-19). *Preprints*. 2020, 2020020407.
6. Darzi AJ, Karam SG, Charide R, Etxeandia Ikobaltzeta I, Cushman M, Gould MK, Mbuagbaw L, Spencer F, Spyropoulos A, Streiff MB, Woller SC, Zakai N, Germini F, Rigoni M, Agarwal A, Morsi RZ, Iorio A, Akl E, Schunemann HJ. Prognostic factors for VTE and Bleeding in Hospitalized Medical Patients: a systematic review and meta-analysis. *Blood*. 2020 Feb 24. pii: blood.2019003603.
7. Mehran R, Rao SV, Bhatt DL, et al. Standardized Bleeding Definitions for Cardiovascular Clinical Trials. A Consensus Report from the Bleeding Academic Research Consortium. *Circulation*. 2011; 123:2736-2747.

SUMMARY OF MAIN PROTOCOL CHANGES

The original version of the protocol was 1.0 and was dated April 2020. Below the rationale and the summary of changes introduced with the various amendments.

Amendment 1 to the protocol – Version 2.0 – May 2020:

Amendment Rationale:

- Updated the primary endpoint:
Primary endpoint was misleading, missing the specification that only proximal deep vein thrombosis has to be considered in the composite analysis
- Changes to trial design:
Removed the need to randomize within 12 hours of hospital admission to allow for better recruitments.
- Modified exclusion criteria:
Modified the criteria for subjects participating in other clinical trials to exclude only those enrolled in competitive clinical trials with antithrombotic agents; all subjects with higher than prophylactic doses of heparin were also excluded.
- Added the interim analysis:
To allow an early evaluation of trial efficacy and safety
- Updated investigator list and recruiting centers
To allow an increased rate of enrollment by adding new centers to the trial

Summary of changes:

- Updated the primary objective (section 2)
- Modification to trial design (section 6)
- Modified the exclusion criteria (section 7.3)
- Updated the list of participating centers (section 8.1)
- Added an ad-interim analysis (section 14.2)

Amendment 2 to the protocol – Version 3.0 – June 2020:

Amendment Rationale:

- The trial was modified in order to become an international multicenter trial:
Change was proposed to permit a better enrollment since Russia was undergoing a high incidence of infections
- Added the references to the sponsor allowing the internationalization of the study:

Sponsor's funding was needed to allow opening centers in Russia by stipulating collaborations with local CRO

- Updates on the insurance policies:
Russians' centers required the stipulation of a local insurance policy
- Updated investigator list and recruiting centers
To allow an increased rate of enrollment by adding new centers to the trial

Summary of changes:

- Modified the recruitment to include Russian centers (section 8.1)
- Updated the list of participating centers (section 8.1)
- Updated the funding references (Heading and section 19.1)
- Updated the insurance policies (section 19.2)

Amendment 3 of the protocol – Version 4.0 – September 2020:

Amendment Rationale:

- Modified trial design:
Updated wordings for enoxaparin treatments that were ambiguous: added the specification that treatment will continue until discharge
- Added a secondary endpoint:
“To compare the incidence of 30 days adverse events, defined as composite of death, stroke and myocardial infarction” , considering the increasing reported association between COVID disease and cardiovascular events.
- Modified the endpoint evaluation timepoints:
Timepoints were modified from a 30-day time to Discharge to cover the usual period managed with thromboprophylaxis in medical patients.
- Increased the duration of the study to 13 months:
Change was needed due to slow enrollment rates
- Updated information on Follow Up procedures:
The follow up procedures were better described, specifying that a telephonic 30-day follow up evaluation is needed for subjects discharged before that timepoint
- Updates on the insurance policies:
The need for an insurance was not needed in Russia for COVID related studies, thus it was removed
- Updated investigator list and recruiting centers

To allow an increased rate of enrollment by adding new centers to the trial

Summary of changes:

- Added a new secondary endpoint (section 2)
- Modified endpoints timepoints (section 5)
- Updated trial design (section 6 and section 8)
- Updated the list of participating centres (section 8.1)
- Modified the end study date and duration (section 6 and section 8.4)
- Clarified follow-up procedures (section 8.6)
- Updated the insurance policies (section 19.2)

COMPLETE LIST OF STUDY OUTCOMES

Primary Objective:

- To compare the effects of 40 mg subcutaneous enoxaparin o.d. versus 40 mg enoxaparin b.i.d. on the incidence of venous thromboembolism (VTE) (a composite of incident asymptomatic and symptomatic proximal deep vein thrombosis (DVT) diagnosed by serial compression ultrasonography (CUS), and symptomatic pulmonary embolism (PE) diagnosed by CT scan), in patients with SARS-CoV-2 infection.

Secondary Objectives:

- To compare the effects of 40 mg subcutaneous enoxaparin o.d. versus 40 mg enoxaparin b.i.d. on the incidence of in-hospital major complications, defined as the composite of death, VTE, use of mechanical ventilation, stroke, acute myocardial infarction and admission to an intensive care in patients with SARS-CoV-2.
- To compare each single component of the primary endpoint between the two groups.
- To compare maximum sequential organ failure assessment (SOFA) score between the two groups.
- To compare C-Reactive Protein, D-dimer, IL-6 and HS-troponin levels (as % above the upper reference limit [URL]) among the two groups.
- To compare the incidence of SARS-CoV-2-related acute respiratory distress syndrome (ARDS) between the two groups.
- To compare length of hospital, stay between the two groups.
- To compare measures of right ventricular function at trans-thoracic echocardiography or CT between admission and follow-up, whenever available.
- To compare the incidence of 30 days adverse events, defined as composite of death, stroke and myocardial infarction.

Safety Endpoints:

- To compare the incidence of bleeding events according to the *ISTH bleeding scale* between the two groups.
- To compare the incidence of bleeding events according to *BARC classification* between the two groups. Only BARC 3 and 5 will be considered for secondary endpoint.
- To assess the incidence of heparin-induced thrombocytopenia in the two groups.

Definitions of Outcome Events

- **All-cause death:** Death due to any causes in the first 30 days. The steering committee made the decision not to adjudicate the cause of death whenever inaccuracies were possible and no systematic autopsies were performed.
- **Deep venous thrombosis:** Incident symptomatic and asymptomatic proximal deep vein thrombosis diagnosed in a major vein of the legs using compression ultrasound.

- **Pulmonary embolism:** Any pulmonary embolism found at autopsy or diagnosed on echocardiography, V/Q scan, CT angiography or invasive pulmonary angiography. ASH 2020 Guidelines were followed for management.(1)
- **Ischemic stroke:** An episode of neurological dysfunction caused by focal cerebral, spinal, or retinal infarction.(2)
- **Type 1 myocardial infarction (T1MI):** It was defined as: “An acute myocardial injury with clinical evidence of acute myocardial ischaemia and with detection of a rise and/or fall of cTn values with at least one value above the 99th percentile URL and at least one of the following: Symptoms of myocardial ischaemia; New ischaemic ECG changes; Development of pathological Q waves; Imaging evidence of new loss of viable myocardium or new regional wall motion abnormality in a pattern consistent with an ischaemic aetiology; Identification of a coronary thrombus by angiography or autopsy (not for types 2 or 3 MIs)”(3)
- **ARDS:** An acute diffuse, inflammatory lung injury, leading to increased pulmonary vascular permeability, increased lung weight, and loss of aerated lung tissue. It was defined, following Berlin 2012 definition, as the acute onset of hypoxemia with bilateral opacities on chest imaging, with no evidence of left atrial hypertension.(4)
- **Heparin-induced thrombocytopenia:** An adverse drug reaction mediated by platelet-activating antibodies that target complexes of platelet factor 4 and heparin. Diagnosed following ASH 2018 Guidelines (5)
- **BARC Scale (6)**
 - Type 0: no bleeding
 - Type 1: bleeding that is not actionable and does not cause the patient to seek unscheduled performance of studies, hospitalization, or treatment by a healthcare professional; may include episodes leading to self-discontinuation of medical therapy by the patient without consulting a healthcare professional.
 - Type 2: any overt, actionable sign of hemorrhage (eg, more bleeding than would be expected for a clinical circumstance, including bleeding found by imaging alone) that does not fit the criteria for type 3, 4, or 5 but does meet at least one of the following criteria: (1) requiring nonsurgical, medical intervention by a healthcare professional, (2) leading to hospitalization or increased level of care, or (3) prompting evaluation.
 - Type 3a: Overt bleeding plus hemoglobin drop of 3 to <5 g/dL (provided hemoglobin drop is related to bleed); Any transfusion with overt bleeding.
 - Type 3b: Overt bleeding plus hemoglobin drop ≥ 5 g/dL (provided hemoglobin drop is related to bleed); Cardiac tamponade; Bleeding requiring surgical intervention for control (excluding dental/nasal/skin/hemorrhoid); Bleeding requiring intravenous vasoactive agents.

Type 3c: Intracranial hemorrhage (does not include microbleeds or hemorrhagic transformation, does include intraspinal); Subcategories confirmed by autopsy or imaging or lumbar puncture; Intraocular bleed compromising vision.

Type 4: CABG-related bleeding: Perioperative intracranial bleeding within 48 h; Reoperation after closure of sternotomy for the purpose of controlling bleeding; Transfusion of ≥ 5 U whole blood or packed red blood cells within a 48-h period; Chest tube output ≥ 2 L within a 24-h period.

Type 5: fatal bleeding

Type 5a: Probable fatal bleeding; no autopsy or imaging confirmation but clinically suspicious

Type 5b: Definite fatal bleeding; overt bleeding or autopsy or imaging confirmation

- **ISTH definition of major bleeding in non-surgical patients:**

Fatal bleeding, and/or

Symptomatic bleeding in a critical area or organ, such as intracranial, intraspinal, intraocular, retroperitoneal, intra-articular or pericardial, or intramuscular with compartment syndrome, and/or

Bleeding causing a fall in haemoglobin level of 20 g L^{-1} (1.24 mmol L^{-1}) or more, or leading to transfusion of two or more units of whole blood or red cells. (7)

- **ISTH definition of major bleeding in surgical patients:**

Fatal bleeding and/or

Bleeding that is symptomatic and occurs in a critical area or organ, such as intracranial, intraspinal, intraocular, retroperitoneal, pericardial, in a non-operated joint, or intramuscular with compartment syndrome, assessed in consultation with the surgeon and/or

Extrasurgical site bleeding causing a fall in haemoglobin level of 2 g/dL (1.24 mmol/L) or more, or leading to transfusion of two or more units of whole blood or red cells, with temporal association within 24–48 h to the bleeding, and/or

Surgical site bleeding that requires a second intervention (open, arthroscopic, endovascular) or a hemarthrosis of sufficient size as to interfere with rehabilitation by delaying mobilization or delayed wound healing, resulting in prolonged hospitalization or a deep wound infection, and/or

Surgical site bleeding that is unexpected and prolonged and/ or sufficiently large to cause hemodynamic instability, as assessed by the surgeon. There should be an associate fall in haemoglobin level of at least 2 g/dL (1.24 mmol/L), or transfusion, indicated by the bleeding, of at least two units of whole blood or red cells, with temporal association within 24 h to the bleeding.

The period for collection of these data is from start of surgery until five half-lives after the last dose of the drug with the longest half-life and with the longest treatment period (in case of unequal active treatment durations).

The population is those who have received at least one dose of the study drug. (8)

- **SOFA Score:** The Sequential Organ Failure Assessment Score, evaluating organ dysfunction in critically ill patients.

Respiration (P/F), Coagulation (Platelets), Liver (Bilirubin), Cardiovascular (hypotension and amine use) , Central Nervous System (Glasgow Coma Scale) and Renal (Creatinine or urine output) were assessed.(9,10)

Variables/score	0	1	2	3	4
PaO ₂ /FiO ₂ (mmHg)	>400	≤400	≤300	≤200	≤100
Platelets (x10 ³ /μl)	>150	≤150	≤100	≤50	≤20
Bilirubin (mg/dl)	<1.2	1.2-1.9	2-5.9	6-11.9	>12
Cardiovascular (Hg/kg/min)	-	MAP<70	Dop≤5	Dop>5 (Epi≤0.1)	Epi>0.1
Glasgow Coma Scale	15	13-14	10-12	6-9	<6
Creatinine (mg/dl)	<1.2	1.2-1.9	2-3.4	3.5-4.9	>5

MAP, Mean Arterial Pressure; Dop, Dopamine; Epi, Epinephrine.


- **Borg Scale:** (11) a 6-to-20-grade scale for ratings of perceived exertion.
 - 6: No exertion at all
 - 7: Very, very light
 - 9: Very light
 - 11: Fairly light
 - 13: Somewhat hard
 - 15: Hard
 - 17: Very hard
 - 19: Very, very hard

REFERENCES


1. Ortel TL, Neumann I, Ageno W, Beyth R, Clark NP, Cuker A, et al. American Society of Hematology 2020 guidelines for management of venous thromboembolism: treatment of deep vein thrombosis and pulmonary embolism. *Blood Advances*. 2 ottobre 2020;4(19):4693–738.
2. Sacco RL, Kasner SE, Broderick JP, Caplan LR, Connors JJ (Buddy), Culebras A, et al. An Updated Definition of Stroke for the 21st Century. *Stroke*. 1 luglio 2013;44(7):2064–89.
3. Thygesen K, Alpert JS, Jaffe AS, Chaitman BR, Bax JJ, Morrow DA, et al. Fourth universal definition of myocardial infarction (2018). *European Heart Journal*. 14 gennaio 2019;40(3):237–69.
4. The ARDS Definition Task Force*. Acute Respiratory Distress Syndrome: The Berlin Definition. *JAMA*. 20 giugno 2012;307(23):2526–33.
5. Cuker A, Arepally GM, Chong BH, Cines DB, Greinacher A, Gruel Y, et al. American Society of Hematology 2018 guidelines for management of venous thromboembolism: heparin-induced thrombocytopenia. *Blood Advances*. 27 novembre 2018;2(22):3360–92.
6. Mehran R, Rao SV, Bhatt DL, Gibson CM, Caixeta A, Eikelboom J, et al. Standardized bleeding definitions for cardiovascular clinical trials: a consensus report from the Bleeding Academic Research Consortium. *Circulation*. 14 giugno 2011;123(23):2736–47.
7. Schulman S, Kearon C. Definition of major bleeding in clinical investigations of antihemostatic medicinal products in non-surgical patients. *Journal of Thrombosis and Haemostasis*. 2005;3(4):692–4.
8. Schulman S, Angerås U, Bergqvist D, Eriksson B, Lassen MR, Fisher W. Definition of major bleeding in clinical investigations of antihemostatic medicinal products in surgical patients. *Journal of Thrombosis and Haemostasis*. 2010;8(1):202–4.
9. Ferreira FL. Serial Evaluation of the SOFA Score to Predict Outcome in Critically Ill Patients. *JAMA*. 10 ottobre 2001;286(14):1754.
10. Vincent JL, Moreno R, Takala J, Willatts S, De Mendonça A, Bruining H, et al. The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure. On behalf of the Working Group on Sepsis-Related Problems of the European Society of Intensive Care Medicine. *Intensive Care Med*. luglio 1996;22(7):707–10.
11. Borg GAV. Psychophysical bases of perceived exertion. *Medicine & Science in Sports & Exercise* 1982. 1982;14(5):377–81.

CTPA COMPARISON BETWEEN GROUPS

Number of total TC, with or without contrast, in the different centers, divided by treatment arm



Symptoms for requiring Angio TC examination (multiple symptoms may have warranted for requiring examination)



CHARACTERISTICS OF SUBJECT WITH PULMONARY EMBOLISM

The following table summarizes the clinical characteristics, treatments and symptoms of subjects that underwent pulmonary embolism.

		001	002	003	004	005	006
ARM	Treatment	40 mg od	40 mg od	40 mg od	40 mg od	40 mg od	40 mg od
ANAGRAPHS	Sex	M	M	M	M	M	M
	Ethnicity	CAUCASIC	CAUCASIC	CAUCASIC	CAUCASIC	ARAB	CAUCASIC
SYMPTOMS	Fever	Yes	No	Yes	No	Yes	No
	Mialgia	No	No	Yes	Yes	No	No
	Asthenia	No	No	Yes	No	No	No
	Headache	No	No	No	No	No	No
	Asthma	No	No	Yes	No	No	No
	Cough	No	No	Yes	Yes	Yes	Yes
	Dyspnea	No	No	Yes	Yes	No	Yes
	Nausea/ Vomiting	No	No	No	No	No	No
	Diarrhea	No	No	No	No	No	No
	Other Symptoms	No	No	No	No	Epigastralgia	Chest Pain, Cardiopalmus
EXAMS	Pulmonary Examination			VM reduction	normal	nds	respiratory insufficiency, mild intercostal algia
	TC Evaluation	ground-glass bilateral multiple thickening		negative for pulmonary embolism	bilateral ground- glass and right basal thickening with right pleuric effusion	Polmonite bilaterale AngioTC positiva per EP	Apical centrilobular emphysema, global increase in lung parenchyma density from confluent subsolid densities prevalent in both upper lobes, Corads 5, parenchymal commitment estimated about 30- 40%
BASELINE	Antiaggregants	Acetylsalilic Acid 100mg bid		Acetylsalilic Acid 100mg bid			
	Anticoagulants						
	Beta-Blockers	Nebivolol 1,25 mg sid		Atenolol 7,5 mg bid			

	Calcium Antagonists						
	Ace Inhibitors	Ramipril 10 mg sid		Elanapril 10 mg bid			
	Sartans						Olmesartan Medoxomil 10 mg od
	Sacubitril / Valsartan						
	Diuretics						Hydrochlorothiazide
	MRAs					Dexamethasone 6 mg od	
	Ivabradin						
	Antiarrhythmics						
	Statins			Rosuvastatin 40 mg od			
	Oral Hypoglicemics					Xelevia 100 mg od	
	Insulin					Yes	
	Immuno Suppressants			Dexamethasone 6 mg od			
	Antivirals			Remdesevir 100 mg od			
	Broncodilators						Acridinium Bromide
IN-HOSPITAL THERAPY	Antibiotics						
	Steroids						Dexamethasone 1.5 ml od
	Antithrombotics						
	Gastric Protector						Omeprazol 20 mg od
	Antiarrhythmics						
	Inotropes						
	Vasopressors						
	Other Drugs						Omelsartan Medoxomil 10 mg od
FU	Follow Up Status	Alive	Lost to FU	Alive	Alive	Alive	Alive

ENROLLMENTS INFORMATIONS

