

Supplementary Table 1 Demographic characteristics of the matched comparison group

	Concussion (n = 336)		Matched Comparison Group (n = 336)		p
	M	F	M	F	
n	203 (60.4%)	133 (39.6%)	199 (61.3%)	130 (38.7%)	.813
Average Age in Months	120 (7.50)	121 (7.42)	119 (7.38)	120 (7.59)	.167
Pubertal Status					
Prepubescence ^a	150 (73.9%)	34 (25.6%)	146 (70.9%)	35 (26.9%)	.816
Pubescence ^b	53 (26.1%)	99 (74.4%)	60 (29.1%)	95 (73.1%)	
Family Income^c					
< 50K	47 (23.2%)	30 (22.6%)	49 (23.8%)	24 (18.5%)	.888
50-99K	65 (32.0%)	33 (24.8%)	58 (28.2%)	38 (29.2%)	
> 100K	91 (44.8%)	70 (52.6%)	99 (48.1%)	68 (52.3%)	
Race					
Non-Hispanic White	124 (61.1%)	97 (72.9)	118 (57.3%)	84 (64.6%)	.203
Non-Hispanic Black	17 (8.4%)	10 (7.5%)	32 (15.5%)	10 (7.7%)	
Hispanic	35 (17.2%)	16 (12.0%)	38 (18.4%)	23 (17.7%)	
Asian ^d	4 (2.0%)	0 (0.0%)	4 (1.9%)	1 (0.8%)	
Other/Multi-Racial	23 (11.3%)	10 (7.5%)	14 (6.8%)	12 (9.2%)	

P values were calculated using Independent t-test for continuous variables and chi-square test for categorical variables.

^aPrepubescence includes children at pubertal stages 1 and 2, as defined by the Physical Health Measure completed by Parents.

^bPubescence includes children at pubertal stages 3-5, as defined by the Physical Health Measure completed by Parents.

^cFamily Income refers to total combined family income.

^dAsian race refers to Asian Indian, Chinese, Filipino, Japanese, Korean, Vietnamese, other Asian.

M, males; F, females.

Supplementary Table 2 Demographic and injury characteristics for participants scanned on Siemens Prisma scanners

	Concussion Scanned on Siemens (n = 223)		Siemens Comparison Group (n = 4498)		p
	M	F	M	F	
n	128 (57.4%)	95 (42.6%)	2381 (52.9%)	2117 (47.1%)	.192
Average Age in Months	121 (7.35)	120 (7.56)	119 (7.41)	119 (7.32)	.010
Pubertal Status					
Prepubescence ^a	94 (73.4%)	23 (24.2%)	1612 (67.7%)	632 (29.9%)	.452
Pubescence ^b	34 (26.6%)	72 (75.8%)	769 (32.3%)	1485 (70.1%)	
Family Income^c					
< 50K	32 (25.0%)	21 (22.1%)	715 (30.0%)	630 (29.8%)	.118
50-99K	43 (33.6%)	24 (25.3%)	648 (27.2%)	672 (31.7%)	
> 100K	53 (41.4%)	50 (52.6%)	1018 (42.8%)	815 (38.5%)	
Race					
Non-Hispanic White	82 (64.1%)	74 (77.9%)	1336 (56.1%)	1154 (54.5%)	<.001
Non-Hispanic Black	14 (10.9%)	8 (8.4%)	368 (15.5%)	321 (15.2%)	
Hispanic	20 (15.6%)	9 (9.5%)	431 (18.1%)	417 (19.7%)	
Asian ^d	3 (2.3%)	0 (0%)	34 (1.4%)	36 (1.7%)	
Other/Multi-Racial	9 (7.0%)	4 (4.2%)	212 (8.9%)	189 (8.9%)	

P values were calculated using Independent t-test for continuous variables and chi-square test for categorical variables.

^aPrepubescence includes children at pubertal stages 1 and 2, and pubescence includes children at pubertal stages 3-5, as defined by the Physical Health Measure completed by Parents.

^cFamily Income refers to total combined family income.

^dAsian race refers to Asian Indian, Chinese, Filipino, Japanese, Korean, Vietnamese, other Asian.

M, males; F, females.

Supplementary Table 3 Group comparisons of mean neurite density between children with concussion and matched comparison group and Siemens Prisma group

		Interaction				Main Effect				R ²
		Group by Sex		Age by Sex		Female Sex		Age		
		β	<i>p</i>	β	<i>p</i>	β	<i>p</i>	β	<i>p</i>	
Matched Comparison	SWM	-0.009	.039	-	-	0.007	.041	0.0007	<.001	0.498
	DWM	<i>-0.007</i>	<i>.070</i>	-	-	0.005	.121	0.0007	<.001	0.470
Siemens Prisma	SWM	-0.010	.007	-	-	0.007	.008	0.0005	<.001	0.182
	DWM	-0.006	.046	0.0002	.040	0.003	.133	0.0004	<.001	0.089

P < .05 is bolded, *p* < .1 is italicized with comparison group as the reference group.

Main effects of sex and age are reported for all models.

R² represents conditional R² which takes both the fixed and random effects into account.


SWM, superficial white matter; DWM, deep white matter.

Supplementary Table 4 Association between neurite density in the concussion group and injury variables


	Superficial White Matter						Deep White Matter					
	All		Female		Male		All		Female		Male	
	β	p	β	p	β	p	β	p	β	p	β	p
Injury Mechanism												
Fight	0.005	.637	0.004	.760	0.016	.302	0.006	.637	0.012	.338	0.009	.577
MVA	-0.0005	.945	-0.013	.299	0.008	.479	-0.018	.039	-0.015	.172	<i>-0.019</i>	<i>.097</i>
Play/Sports	0.003	.890	-0.004	.449	0.010	.049	0.0001	.979	-0.004	.396	0.003	.596
Other	-0.002	.806	-0.005	.726	0.003	.832	0.004	.806	-0.003	.809	0.008	.603
Number of Injuries	0.0002	.969	0.003	.647	-0.011	.235	-0.002	.969	-0.002	.773	-0.008	.364
w/ Loss of Consciousness <30 minutes	0.004	.580	<i>0.010</i>	<i>.066</i>	0.002	.710	-0.002	.667	0.008	.109	-0.007	.276
w/ Post-Traumatic Amnesia <24 hours	0.006	.477	-0.001	.904	<i>0.013</i>	<i>.064</i>	0.003	.505	-0.0004	.953	0.006	.437
Age at First Injury	-0.00002	.912	-0.0001	.668	0.0002	.544	-0.00003	.912	-0.00002	.939	0.0001	.687
Time Since First Injury	-0.00001	.946	-0.0002	.366	0.0003	.402	0.00002	.946	-0.0001	.612	0.0002	.459

$P < .05$ is bolded, $p < .1$ is italicized.

Effects are reported for concussion group analyses and sex stratified analyses.



Supplementary Figure 1 Mean ND in children with concussion compared to matched comparison group. (A) Deep white matter. (B) Superficial white matter.



Supplementary Figure 2 Mean ND from Siemens scanners in children with concussion compared to comparison group. (A) Deep white matter. (B) Superficial white matter.