

## Supplementary Material




Figure S1: Flowchart of participation and exclusions in each cohort yielding the study population.

**Table S1: Overview of information and various measures used in study**

Denmark DBDS	Estonia EstBB-C19	Iceland C19- Resilience	Norway MAP-19	Norway MoBa	Sweden Omtanke2020	UK CovidLife
<b>Ethical approvals</b>						
Zealand and Central Denmark Regional Committees on Health Research Ethics (SJ-740 and 1-10-72-95-13) and the Data Protection Agency (P-2019-99)	Estonian Committee on Bioethics and Human Research (1.1–12/1277)	National Bioethics Committee (NBC no. 20–073, 21–071) as well as the National Data Protection Authority	Regional Committee for Medical Research Ethics, reference number: 125510	Regional Committees for Medical and Health Research Ethics (127708/1414 0/20138)	Ethical approval no. 2020–01785	East of Scotland Research Ethics Service (EoSRES)
<b>COVID-19 related outcomes</b>						
<b>COVID-19 diagnosis</b>	Self-report of confirmed PCR-test	Self-report of confirmed PCR-test	Self-report of confirmed PCR-test	Self-report of confirmed PCR-test	Self-report of confirmed PCR-test	Self-report of confirmed PCR-test or positive antibody test with date of testing provided
<b>COVID-19 illness severity</b>	Never bedridden/Bedridden 1 day or more	No fever/ Fever 1-6 days/Fever 7 days or more	Never bedridden/Bedridden 1-6 days/Bedridden 7 days or more	Not measured	Never bedridden/Bedridden 1-6 days/Bedridden 7 days or more	Never bedridden/Bedridden 1-6 days/Bedridden 7 days or more (only asked at long surveys (baseline and 6-months FU))
<b>COVID-19 hospitalization</b>	Not measured	Non-hospitalized/Hospitalized (non-ICU)/Hospitalized (ICU)	Non-hospitalized/Hospitalized (non-ICU)/Hospitalized (ICU)	Not measured	Non-hospitalized/Hospitalized	Non-hospitalized/Hospitalized (non-ICU)/Hospitalized (ICU)
<b>Mental health instruments</b>						
<b>Depression</b>	PHQ-9	EST-Q2	PHQ-9	PHQ-9	PHQ-9	PHQ-9
<b>Anxiety</b>	ASS	EST-Q2	GAD-7	GAD-7	GAD-7	GAD-7
<b>COVID-19 related distress</b>	PC-PTSD-5 (modified)	Not measured	PC-PTSD-5 (modified)	PCL-5	PC-PTSD-5 (modified)	PC-PTSD-5 (modified)
<b>Sleep quality</b>	Not measured	EST-Q2	PSQI	Not measured	PSQI	One item of PSQI
						Binary variable of better/same sleep quality versus worse sleep quality
<b>Covariates</b>						
<b>Age</b>	x	x	x	x	x	x
<b>Gender</b>	x	x	x	x	x	x
<b>Education</b>	x	x	x	x	x	—
<b>Relationship status</b>	x	—	x	x	—	x
<b>Smoking</b>	x	x	x	x	x	x
<b>BMI</b>	x	x	x	x	x	x
<b>Pre-existing psychiatric conditions</b>	*	*	x	x	x	x
<b>Number of chronic medical conditions</b>	x	x	x	—	x	x
<b>Response period</b>	x	x	x	x	x	x

Note: Data from surveys are marked with x; register data is marked with a \*. “x” and “\*\*” denotes covariates were included in model while “—” indicates covariate was not included

Table S2: **Background characteristics of study participants that received a COVID-19 diagnosis at any time point and others in the seven COVIDMENT cohorts separately, combined, and combined with all missing values removed**

	Denmark		Estonia		Iceland		Norway				Sweden		United Kingdom		Overall		Overall	
	(DBDS)		(EstBB C-19)		(C19-Resilience)		(MAP-19)		(MoBa)		(Omtanke)		(CovidLife)				without missing values	
	(N=30 917)		(N=9989)		(N=21 615)		(N = 10 061)		(N=132 451)		(N=23 698)		(N=18 518)		(N= 247 249)		(N=247 249)	
	COVID-19 diagnosis	Others	COVID-19 diagnosis	Others	COVID-19 diagnosis	Others	COVID-19 diagnosis	Others	COVID-19 diagnosis	Others	COVID-19 diagnosis	Others	COVID-19 diagnosis	Others	COVID-19 diagnosis	Others	COVID-19 diagnosis	Others
	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)
Total	1111 (3.5)	29 806 (96.4)	2121 (21.2)	7868 (78.8)	1144 (5.3)	20 471 (94.7)	110 (1.1)	9951 (98.9)	1995 (1.5)	130 456 (98.5)	3175 (13.4)	20 523 (86.6)	323 (1.7)	18 195 (98.2)	9979 (4)	237270 (96)	9979 (4)	237270 (96)
Gender																		
Male	513 (46.2)	14 341 (48.1)	740 (34.9)	2224 (28.3)	424 (37.1)	6125 (29.9)	17 (15.5)	2166 (21.8)	815 (40.9)	56 098 (43.0)	607 (19.1)	3796 (18.5)	86 (26.6)	5928 (32.6)	3202 (32.1)	90678 (38.2)	3202 (32.1)	90678 (38.2)
Female	598 (53.8)	15 465 (51.9)	1381 (65.1)	5664 (71.7)	718 (62.8)	14 309 (69.9)	92 (83.6)	7760 (78.0)	1180 (59.1)	74 358 (57.0)	2568 (80.9)	16 727 (81.5)	235 (72.8)	12 138 (66.7)	6772 (67.9)	146401 (61.7)	6772 (67.9)	146401 (61.7)
Other	—	—	—	—	2 (0.2)	37 (0.2)	1 (0.9)	25 (0.3)	—	—	**	**	—	—	3 (0)	62 (0)	3 (0)	62 (0)
Missing	—	—	—	—	—	—	—	—	—	—	—	—	2 (0.6)	129 (0.7)	2 (0)	129 (0.1)	2 (0)	129 (0.1)
Age																		
Mean age (SD)	46.6 (14.8)	54.6 (16.6)	44.5 (13.4)	44.7 (13.3)	48.2 (14.7)	54.8 (14.1)	33.3 (12.2)	35.7 (13.5)	47.1 (5.41)	46.7 (5.39)	47.1 (13.8)	50.4 (16.1)	52.4 (13.5)	56.5 (14.3)	46.6	48.9	46.6	48.9
18-29 years	200 (18.0)	3053 (10.2)	273 (12.9)	970 (12.3)	168 (14.7)	1216 (5.9)	54 (49.1)	4349 (43.7)	—	—	419 (13.2)	2532 (12.3)	23 (7.1)	964 (5.3)	1137 (11.4)	13084 (5.5)	1137 (11.4)	13084 (5.5)
30-39 years	194 (17.5)	3554 (11.9)	573 (27.0)	2143 (27.2)	154 (13.5)	1974 (9.6)	23 (20.9)	2254 (22.7)	154 (7.7)	11 039 (8.5)	565 (17.8)	3542 (17.3)	38 (11.8)	1760 (9.7)	1701 (17)	26266 (11.1)	1701 (17.1)	26266 (11.1)
40-49 years	261 (23.5)	4911 (16.5)	563 (26.5)	1963 (24.9)	259 (22.6)	3620 (17.7)	21 (19.1)	1567 (15.7)	1232 (61.8)	81 932 (62.8)	677 (21.3)	3596 (17.5)	59 (18.3)	2508 (13.8)	3072 (30.8)	100097 (42.2)	3072 (30.8)	100097 (42.2)
50-59 years	244 (22.0)	6152 (20.5)	396 (18.7)	1607 (20.4)	284 (24.8)	5280 (25.8)	6 (5.5)	1088 (10.9)	575 (28.8)	35 620 (27.3)	908 (28.6)	4300 (21.0)	99 (30.7)	3955 (21.7)	2512 (25.2)	58002 (24.4)	2512 (25.2)	58002 (24.4)
60-69 years	117 (10.5)	4386 (14.7)	221 (10.4)	856 (10.9)	206 (18.0)	5335 (26.1)	5 (4.5)	557 (5.6)	33 (1.7)	1768 (1.4)	462 (14.6)	3505 (17.1)	74 (22.9)	5454 (30.0)	1118 (11.2)	21861 (9.2)	1118 (11.2)	21861 (9.2)
70 years +	95 (8.6)	7750 (26.0)	95 (4.5)	329 (4.2)	73 (6.4)	3046 (14.9)	1 (0.9)	136 (1.4)	1 (0.1)	97 (0.1)	144 (4.5)	3048 (14.9)	26 (8.0)	3348 (18.4)	435 (4.4)	17754 (7.5)	435 (4.4)	17754 (7.5)
Missing	—	—	—	—	—	—	—	—	—	—	—	—	4 (1.2)	206 (1.1)	4 (0)	206 (0.1)	4 (0)	206 (0.1)
Education																		
Compulsory	30 (2.7)	1501 (5.0)	44 (2.1)	140 (1.8)	144 (12.6)	2916 (14.2)	3 (2.7)	519 (5.2)	38 (1.9)	3498 (2.7)	*	*	23 (7.1)	1515 (8.3)	286 (2.9)	10469 (4.4)	290 (4.4)	10469 (5.2)
Upper secondary, vocational or other	60 (5.4)	1144 (3.8)	884 (41.7)	2825 (35.9)	361 (31.6)	6359 (31.1)	15 (13.6)	1771 (17.8)	606 (30.4)	42 530 (32.6)	*	*	103 (31.9)	6050 (33.3)	2029 (20.3)	60679 (25.6)	2029 (30.9)	60679 (30)
Bachelor's/ diploma university degree	698 (62.8)	20 142 (67.6)	582 (27.4)	2182 (27.7)	390 (34.1)	6473 (31.6)	92 (83.6)*1	7661 (77.0)	661 (33.1)	42 765 (32.8)	*	*	84 (26.0)	4120 (22.6)	2507 (25.1)	83343 (35.1)	2507 (38.2)	83343 (41.2)



April-June 2020	125 (11.3)	52 (0.2)	138 (6.5)	1113 (14.1)	372 (32.5)	19 807 (96.8)	14 (12.7)*3	9951 (100)	346 (17.3)	130 456 (100)	45 (1.4)	317 (1.5)	70 (21.7)	18 195 (100)	1110 (11.1)	179891 (75.8)	1110 (11.1)	179891 (75.8)
July-September 2020	—	—	155 (7.3)	1133 (14.4)	9 (0.8)	232 (1.1)	3 (2.7)	-	292 (14.6)	-	1322 (41.6)	9262 (45.1)	20 (6.2)	—	1801 (18)	10627 (4.5)	1801 (18)	10627 (4.5)
October-December 2020	947 (85.2)	29 049 (97.5)	166 (7.8)	1328 (16.9)	139 (12.2)	396 (1.9)	26 (23.6)	-	682 (34.2)	-	1368 (43.1)	8819 (43.0)	—	—	3328 (33.4)	39592 (16.7)	3328 (33.4)	39592 (16.7)
January-March 2021	39 (3.5)	705 (2.4)	905 (42.7)	2499 (31.8)	574 (50.2)	32 (0.2)	28 (25.5)	-	86 (4.3)	-	319 (10.0)	1524 (7.4)	233 (72.1)	—	2184 (21.9)	4760 (2)	2184 (21.9)	4760 (2)
April-June 2021	—	—	757 (35.7)	1795 (22.8)	50 (4.4)	4 (0.0)	39 (35.5)	-	589 (29.5)	-	121 (3.8)	601 (2.9)	—	—	1556 (15.6)	2400 (1)	1556 (15.6)	2400 (1)
<b>Illness severity</b>																		
Never bedridden	560 (50.4)	—	1061 (50.0)	—	428 (37.4)	—	-	-	636 (31.9)	—	475 (15.0)	—	—	—	3160 (31.7)	—	3160 (43.7)	—
Bedridden 1-6 days	551 (49.6)	—	773 (36.4)	—	455 (39.8)	—	-	-	213 (10.7)	—	461 (14.5)	—	—	—	2453 (24.6)	—	2453 (33.9)	—
Bedridden 7 days or more	—	—	287 (13.5)	—	261 (22.8)	—	-	-	685 (34.3)	—	380 (12.0)	—	—	—	1613 (16.2)	—	1613 (22.3)	—
Missing	—	—	—	—	—	—	-	-	461 (23.1)	—	1859 (58.6)	—	—	—	2753 (27.6)	—	—	—
Non-Hospitalized	—	—	2073 (97.7)	—	1056 (92.3)	—	-	-	1495 (74.9)	—	3090 (97.3)	—	286 (88.5)	—	8000 (80.2)	—	8000 (96.4)	—
Hospitalized	—	—	48 (2.3)	—	88 (7.7)	—	-	-	39 (2.0)	—	85 (0.3)	—	37 (11.4)	—	297 (3)	—	297 (3.6)	—
Missing	—	—	—	—	—	—	-	-	461 (23.1)	—	—	—	—	—	1682 (16.9)	—	—	—
<b>Time since diagnosis</b>																		
0-2 months	594 (53.5)	—	1055 (49.7)	—	444 (38.8)	—	17 (15.5)	-	568 (28.5)	—	215 (6.8)	—	215 (66.6)	—	3108 (31.1)	—	3108 (31.1)	—
2-6 months	373 (33.6)	—	954 (45.0)	—	541 (47.3)	—	43 (39.1)	-	951 (47.7)	—	682 (21.5)	—	98 (30.3)	—	3642 (36.5)	—	3642 (36.5)	—
6-16 months	144 (13.0)	—	112 (5.3)	—	159 (13.9)	—	50 (45.5)	-	476 (23.9)	—	2278 (71.7)	—	10 (3.1)	—	3229 (32.4)	—	3229 (32.4)	—
Missing	—	—	—	—	—	—	-	-	—	—	—	—	—	—	0 (0)	—	—	—

\* Information obtained from registers

\*\* Response option not offered to participants

\*\*\*Data only collected in long follow-ups, hence missing data independent of hospitalisation. Participants who got diagnosed at the other eight follow-ups have not answered this question

#### MAP-19

\*1 This number include higher education in Norway, encompassing both Bachelor, Master and PhD.

\*2 We ask about a current psychological disorder

\*3 do you want the number of COVID-19 infected at that timepoint or whether they filled out the questionnaires? Currently it is those who filled out the questionnaires at the different timepoints

**Table S3: Overview of total number of COVID-19 cases per country and proportion of population diagnosed with COVID-19<sup>1</sup>**

Cohort	Location	End of study	Total cases at end of study <sup>a</sup>	Population of country	Proportion of population with confirmed COVID-19 diagnosis
DBDS	Denmark	August 2021	347 028	5 813 302	6.0%
EstBB-C19	Estonia	August 2021	141 956	1 325 188	10.7%
C19-Resilience	Iceland	August 2021	10 789	343 360	3.1%
MAP-19	Norway	August 2021	160 174	5 465 629	2.9%
MoBa	Norway	May 2021	125 116	5 465 629	2.3%
Omtanke2020	Sweden	June 2021	1 089 990	10 160 159	10.7%
CovidLife	United Kingdom	February 2021	4 188 827	68 207 114	6.1%

<sup>a</sup>Number of cases on the last day of the month

Table S4: Prevalence and prevalence ratios of mental health indicators among individuals with and without a diagnosis of COVID-19, overall and in subgroup analyses by illness severity and time from diagnosis for each cohort. Combined prevalence for each mental health indicator included as well.

	DBDS (DK)															
	N=30 917															
	Overall		Diagnosed 0-2 months ago				Diagnosed 2-6 months ago				Diagnosed 6-16 months ago					
	n	n <sub>threshold</sub> (%)	PR (age+gender adj)	PR (multivar adj) <sup>a</sup>	n	n <sub>threshold</sub> (%)	PR (age+gender adj)	PR (multivar adj) <sup>a</sup>	n	n <sub>threshold</sub> (%)	PR (age+gender adj)	PR (multivar adj) <sup>a</sup>	n	n <sub>threshold</sub> (%)	PR (age+gender adj)	PR (multivar adj) <sup>a</sup>
<b>Depression (PHQ-9 / EST-Q2)</b>																
No confirmed diagnosis	29802	2232 (7.5)	Ref.	Ref.	-	-	-	-	-	-	-	-	-	-	-	-
Ever diagnosed with COVID-19	1111	136 (12.2)	1.33 (1.13-1.56)	1.45 (1.23-1.71)	594	80 (13.5)	1.49 (1.22-1.83)	1.64 (1.33-2.03)	373	41 (11.0)	1.20 (0.90-1.59)	1.28 (0.95-1.73)	144	15 (10.4)	1.06 (0.66-1.70)	1.05 (0.65-1.70)
Never bedridden	560	42 (7.5)	0.81 (0.61-1.09)	0.90 (0.67-1.22)	311	27 (8.7)	0.97 (0.68-1.38)	1.09 (0.76-1.56)	189	10 (5.3)	0.57 (0.31-1.04)	0.63 (0.34-1.17)	60	5 (8.3)	0.79 (0.34-1.83)	0.82 (0.34-1.96)
Bedridden 1-6 days	551	94 (17.1)	1.85 (1.55-2.24)	1.97 (1.63-2.40)	283	53 (18.7)	2.05 (1.61-2.62)	2.22 (1.72-2.87)	184	31 (16.8)	1.85 (1.35-2.52)	1.93 (1.38-2.67)	84	10 (11.9)	1.26 (0.71-2.24)	1.22 (0.69-2.16)
Bedridden 7 days or more	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hospitalized	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Hospitalized	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<b>Anxiety symptoms (GAD-7 / ASS / EST-Q2)</b>																
No confirmed diagnosis	29806	909 (3.0)	Ref.	Ref.	-	-	-	-	-	-	-	-	-	-	-	-
Ever diagnosed with COVID-19	1111	39 (3.5)	0.95 (0.96-1.30)	0.91 (0.66-1.26)	594	23 (3.9)	1.04 (0.70-1.57)	1.02 (0.68-1.54)	373	12 (3.2)	0.89 (0.51-1.55)	0.82 (0.46-1.47)	144	<5	-	-
Never bedridden	560	12 (2.1)	0.57 (0.32-1.00)	0.55 (0.31-0.98)	311	8 (2.6)	0.69 (0.34-1.36)	0.68 (0.34-1.36)	189	<5	—	—	60	<5	-	-
Bedridden 1-6 days	551	27 (4.9)	1.35 (0.93-1.96)	1.29 (0.88-1.88)	283	15 (5.3)	1.45 (0.88-2.38)	1.40 (0.85-2.30)	184	9 (4.9)	1.40 (0.73-2.65)	1.27 (0.66-2.46)	84	<5	-	-



	EstBB C-19 (EE)															
	N=9989															
			Overall			Diagnosed 0-2 months ago				Diagnosed 2-6 months ago				Diagnosed 6-16 months ago		
	n	n <sub>threshold</sub> (%)	PR (age+gender adj)	PR (multivar adj) <sup>a</sup>	n	n <sub>threshold</sub> (%)	PR (age+gender adj)	PR (multivar adj) <sup>a</sup>	n	n <sub>threshold</sub> (%)	PR (age+gender adj)	PR (multivar adj) <sup>a</sup>	n	n <sub>threshold</sub> (%)	PR (age+gender adj)	PR (multivar adj) <sup>a</sup>
<b>Depression (PHQ-9 / EST-Q2)</b>																
No confirmed diagnosis	7868	1640 (20.8)	Ref.	Ref.	-	-	-	-	-	-	-	-	-	-	-	-
Ever diagnosed with COVID-19	2121	402 (19.0)	0.92 (0.84-1.02)	0.91 (0.83-1.00)	1055	211 (20.0)	0.96 (0.85-1.09)	0.92 (0.82-1.04)	954	171 (17.9)	0.87 (0.76-1.01)	0.89 (0.77-1.03)	112	20 (17.9)	0.92 (0.63-1.35)	0.90 (0.61-1.33)
Never bedridden	1061	174 (16.4)	0.80 (0.70-0.92)	0.80 (0.70-0.92)	513	88 (17.2)	0.83 (0.68-1.00)	0.82 (0.69-0.99)	484	75 (15.5)	0.76 (0.62-0.94)	0.76 (0.62-0.94)	64	11 (17.2)	0.94 (0.57-1.55)	0.88 (0.53-1.46)
Bedridden 1-6 days	773	163 (21.1)	0.99 (0.86-1.14)	0.96 (0.84-1.11)	403	91 (22.6)	1.05 (0.88-1.25)	0.96 (0.81-1.15)	339	66 (19.5)	0.93 (0.74-1.15)	0.96 (0.77-1.20)	31	6 (22.9)	0.92 (0.45-1.88)	0.90 (0.44-1.85)
Bedridden 7 days or more	287	65 (22.6)	1.20 (0.97-1.49)	1.19 (0.97-1.48)	139	32 (23.0)	1.27 (0.95-1.70)	1.19 (0.89-1.60)	131	30 (22.9)	1.17 (0.85-1.61)	1.22 (0.89-1.66)	17	3 (17.6)	0.86 (0.33-2.22)	0.98 (0.34-2.80)
Non-Hospitalized	2073	396 (19.1)	0.93 (0.84-1.02)	0.92 (0.83-1.01)	1040	209 (20.1)	0.97 (0.85-1.09)	0.92 (0.82-1.04)	931	168 (18.0)	0.88 (0.76-1.01)	0.90 (0.78-1.04)	102	19 (18.6)	0.95 (0.64-1.40)	0.92 (0.61-1.36)
Hospitalized	35	5 (14.3)	0.82 (0.38-1.78)	0.80 (0.36-1.79)	12	2 (16.7)	0.99 (0.28-3.46)	1.08 (0.28-4.08)	16	2 (12.5)	0.73 (0.21-2.56)	0.62 (0.18-2.07)	7	1 (16.7)	0.72 (0.14-3.65)	0.87 (0.14-5.32)
<b>Anxiety symptoms (GAD-7 / ASS / EST-Q2)</b>																
No confirmed diagnosis	7868	1540 (19.6)	Ref.	Ref.	-	-	-	-	-	-	-	-	-	-	-	-
Ever diagnosed with COVID-19	2121	379 (17.9)	0.93 (0.85-1.03)	0.97 (0.87-1.07)	1055	193 (18.3)	0.95 (0.84-1.09)	0.96 (0.84-1.09)	954	164 (17.2)	0.90 (0.78-1.03)	0.96 (0.83-1.11)	112	22 (19.6)	1.08 (0.75-1.57)	1.05 (0.72-1.53)
Never bedridden	1061	174 (16.4)	0.86 (0.75-0.99)	0.89 (0.78-1.02)	513	89 (17.3)	0.90 (0.75-1.09)	0.94 (0.78-1.12)	484	72 (14.9)	0.78 (0.63-0.97)	0.81 (0.66-1.01)	64	13 (20.3)	1.18 (0.74-1.90)	1.10 (0.67-1.80)
Bedridden 1-6 days	773	144 (18.6)	0.94 (0.81-1.09)	0.97 (0.83-1.12)	403	83 (20.6)	1.03 (0.85-1.25)	1.00 (0.83-1.21)	339	57 (16.8)	0.86 (0.68-1.08)	0.95 (0.75-1.20)	31	4 (12.9)	0.66 (0.27-1.63)	0.63 (0.26-1.56)

Bedridden 7 days or more	287	61 (21.3)	1.20 (0.96-1.50)	1.27 (1.01-1.59)	139	21 (15.1)	0.89 (0.61-1.31)	0.92 (0.62-1.37)	131	35 (26.7)	1.44 (1.08-1.92)	1.56 (1.18-2.08)	17	5 (29.4)	1.54 (0.77-3.09)	1.80 (0.95-3.38)
Non-Hospitalized	2073	372 (17.9)	0.93 (0.85-1.03)	0.97 (0.87-1.07)	1040	192 (18.5)	0.96 (0.84-1.09)	0.97 (0.85-1.10)	931	160 (17.2)	0.89 (0.77-1.03)	0.96 (0.83-1.11)	102	20 (19.6)	1.07 (0.73-1.58)	1.03 (0.69-1.53)
			0.52 (0.18-1.49)	0.53 (0.18-1.54)	12	1 (8.3)	0.51 (0.08-3.36)	0.58 (0.08-3.99)	16	1 (6.2)	0.39 (0.06-2.59)	0.35 (0.06-2.20)	7	1 (14.3)	0.77 (0.15-3.94)	0.86 (0.15-5.07)
<b>COVID-19 related distress symptoms (PC-PTSD-5 / PCL-5)</b>																
No confirmed diagnosis																
Ever diagnosed with COVID-19	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Never bedridden	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Bedridden 1-6 days	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Bedridden 7 days or more	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Non-Hospitalized	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Hospitalized	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
<b>Sleep quality (PSQI, EST-Q2, Single-item measures)</b>																
No confirmed diagnosis																
Ever diagnosed with COVID-19	7868	2406 (30.6)	Ref.	Ref.	-	-	-	-	-	-	-	-	-	-	-	
	2121	642 (30.3)	1.01 (0.94-1.08)	1.03 (0.96-1.11)	1055	304 (28.8)	0.97 (0.88-1.08)	0.98 (0.89-1.09)	954	295 (30.9)	1.02 (0.92-1.12)	1.07 (0.97-1.18)	112	43 (38.4)	1.26 (0.99-1.60)	1.19 (0.94-1.52)
Never bedridden	1061	319 (30.1)	1.00 (0.91-1.10)	1.02 (0.93-1.12)	513	142 (27.7)	0.94 (0.81-1.08)	0.96 (0.84-1.10)	484	151 (31.2)	1.03 (0.90-1.17)	1.05 (0.92-1.20)	64	26 (40.6)	1.30 (0.96-1.76)	1.24 (0.91-1.70)
	773	223 (28.8)	0.98 (0.87-1.09)	0.99 (0.88-1.11)	403	117 (29.0)	0.99 (0.85-1.16)	0.97 (0.83-1.14)	339	94 (27.7)	0.93 (0.78-1.10)	0.99 (0.83-1.18)	31	12 (38.7)	1.31 (0.85-2.03)	1.17 (0.75-1.82)
Bedridden 7 days or more	287	100 (34.8)	1.13 (0.96-1.32)	1.19 (1.01-1.39)	139	45 (32.4)	1.06 (0.83-1.35)	1.10 (0.86-1.41)	131	50 (38.2)	1.21 (0.97-1.49)	1.30 (1.06-1.60)	17	5 (29.4)	1.01 (0.48-2.11)	1.04 (0.51-2.10)

Non-Hospitalized	2073	628 (30.3)	1.01 (0.94-1.09)	1.04 (0.96-1.11)	1040	299 (28.7)	0.97 (0.88-1.08)	0.98 (0.89-1.08)	931	288 (30.9)	1.02 (0.92-1.13)	1.07 (0.97-1.19)		41 (40.2)	1.32 (1.04-1.67)	1.24 (0.97-1.59)
	Hospitalized	35	10 (28.6)	0.89 (0.53-1.48)	0.91 (0.56-1.47)	12	4 (33.3)	1.02 (0.48-2.19)	1.18 (0.55-2.51)	16	4 (25.0)	0.76 (0.33-1.74)	0.73 (0.33-1.60)	7	2 (28.6)	0.98 (0.31-3.02)



No confirmed diagnosis	20 471	5578 (27.2)	Ref.	Ref.	-	-	-	-	-	-	-	-	-	-	-	
Ever diagnosed with COVID-19	1144	393 (34.4)	1.15 (1.05-1.25)	1.09 (0.97-1.24)	444	144 (32.4)	1.08 (0.95-1.24)	1.08 (0.94-1.24)	541	206 (38.1)	1.24 (1.11-1.38)	1.04 (0.83-1.30)	159	43 (27.0)	0.97 (0.75-1.25)	0.74 (0.51-1.06)
Never bedridden	428	127 (29.7)	0.95 (0.82-1.11)	0.94 (0.79-1.11)	159	43 (27.0)	0.87 (0.67-1.12)	0.92 (0.71-1.18)	206	63 (30.6)	0.96 (0.78-1.19)	0.83 (0.62-1.11)	63	21 (33.3)	1.14 (0.81-1.61)	0.90 (0.58-1.38)
	455	154 (33.8)	1.11 (0.98-1.27)	1.02 (0.87-1.19)	165	51 (30.9)	1.03 (0.82-1.29)	0.96 (0.76-1.20)	236	92 (39.0)	1.26 (1.07-1.47)	1.03 (0.79-1.33)	54	11 (20.4)	0.70 (0.42-1.17)	0.56 (0.32-0.99)
Bedridden 1-6 days	261	112 (42.9)	1.55 (1.35-1.69)	1.44 (1.22-1.69)	120	50 (41.7)	1.49 (1.21-1.83)	1.49 (1.22-1.83)	99	51 (51.5)	1.81 (1.49-2.21)	1.45 (1.10-1.91)	42	11 (26.2)	1.07 (0.64-1.77)	0.72 (0.40-1.28)
Non-Hospitalized	1056	356 (33.7)	1.10 (1.01-1.21)	1.06 (0.94-1.20)	407	130 (31.9)	1.05 (0.91-1.21)	1.06 (0.91-1.23)	505	187 (37.0)	1.19 (1.06-1.33)	1.00 (0.79-1.25)	144	39 (27.1)	0.94 (0.72-1.23)	0.73 (0.51-1.05)
	61	29 (47.5)	1.93 (1.51-2.48)	1.66 (1.27-2.17)	22	10 (45.5)	1.74 (1.18-2.55)	1.47 (1.00-2.17)	30	16 (53.3)	2.16 (1.55-3.00)	1.74 (1.17-2.60)	9	3 (33.3)	1.63 (0.63-4.18)	1.10 (0.40-2.98)





(GAD-7 / ASS / EST-Q2)																
No confirmed diagnosis	98 195	2724 (2.8)	Ref.	Ref.	-	-	-	-	-	-	-	-	-	-	-	-
	295	10 (3.4)	1.26 (0.68-2.31)	1.26 (0.69-2.32)	102	2 (2.0)	0.72 (0.18-2.87)	0.59 (0.15-2.28)	193	8 (4.1)	1.54 (0.78-3.03)	1.77 (0.89-3.50)	-	-	-	-
Ever diagnosed with COVID-19	79	3 (3.8)	1.43 (0.47-4.34)	1.64 (0.58-4.63)	10	1 (10.0)	4.22 (0.66-26.99)	4.02 (1.02-15.78)	69	2 (2.9)	1.07 (0.27-4.21)	1.27 (0.34-4.72)	-	-	-	-
	19	0 (0)	-	-	2	0 (0)	-	-	17	0 (0)	-	-	-	-	-	-
	122	6 (4.9)	1.81 (0.83-3.94)	2.07 (0.93-4.6)	15	0 (0)	-	-	107	6 (5.6)	2.08 (0.96-4.52)	2.39 (1.08-5.32)	-	-	-	-
	212	9 (4.2)	1.57 (0.83-2.98)	1.78 (0.95-3.37)	23	1 (4.3)	1.57 (0.23-10.77)	1.67 (0.30-9.36)	189	8 (4.2)	1.57 (0.80-3.09)	1.8 (0.91-3.57)	-	-	-	-
Hospitalized	8	0 (0.0)	-	-	4	0 (0)	-	-	4	0 (0)	-	-	-	-	-	-
COVID-19 related distress symptoms (PC-PTSD-5 / PCL-5)																
No confirmed diagnosis	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Ever diagnosed with COVID-19	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Never bedridden	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Bedridden 1-6 days	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Bedridden 7 days or more	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hospitalized	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Hospitalized	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Sleep quality (PSQI, EST-Q2, Single-item measures)																
No confirmed diagnosis	77 044	16 062 (20.8)	Ref.	Ref.	-	-	-	-	-	-	-	-	-	-	-	-
	1061	289 (27.2)	1.32 (1.19-1.46)	1.32 (1.20-1.46)	205	49 (23.9)	1.15 (0.90-1.47)	1.15 (0.90-1.47)	568	162 (28.5)	1.38 (1.21-1.57)	1.36 (1.20-1.55)	288	78 (27.1)	1.32 (1.09-1.60)	1.35 (1.12-1.63)

Never bedridden	398	80 (20.1)	0.98 (0.81-1.20)	1.01 (0.83-1.23)	62	12 (19.4)	0.94 (0.57-1.56)	0.94 (0.57-1.55)	243	52 (21.4)	1.04 (0.82-1.33)	1.07 (0.84-1.36)	93	16 (17.2)	0.86 (0.55-1.35)	0.91 (0.58-1.43)
Bedridden 1-6 days	124	31 (25.0)	1.24 (0.92-1.68)	1.24 (0.92-1.67)	46	10 (21.7)	1.07 (0.62-1.85)	1.09 (0.64-1.87)	61	18 (29.5)	1.46 (0.99-2.14)	1.43 (0.98-2.08)	17	3 (17.6)	0.91 (0.32-2.53)	0.95 (0.36-2.48)
Bedridden 7 days or more	410	145 (35.4)	1.68 (1.47-1.91)	1.64 (1.44-1.87)	61	18 (29.5)	1.38 (0.94-2.04)	1.35 (0.91-2.00)	215	80 (37.2)	1.75 (1.47-2.08)	1.69 (1.42-2.01)	134	47 (35.1)	1.69 (1.34-2.12)	1.70 (1.35-2.13)
Non-Hospitalized	912	249 (27.3)	1.32 (1.18-1.47)	1.32 (1.19-1.47)	168	39 (23.2)	1.12 (0.85-1.47)	1.11 (0.85-1.46)	510	149 (29.2)	1.40 (1.22-1.61)	1.40 (1.22-1.60)	234	61 (26.1)	1.27 (1.03-1.58)	1.32 (1.06-1.63)
Hospitalized	20	7 (35.0)	1.85 (1.03-3.30)	1.71 (0.95-3.07)	1	1 (100.0)	5.98 (5.83-6.14)	6.33 (6.01-6.67)	9	1 (11.1)	0.60 (0.10-3.63)	0.58 (0.10-3.36)	10	5 (50.0)	2.54 (1.38-4.66)	2.25 (1.17-4.3)

	Omtanke (SE)															
	N=23 698															
	Overall				Diagnosed 0-2 months ago				Diagnosed 2-6 months ago				Diagnosed 6-16 months ago			
	n	n <sub>threshold</sub> (%)	PR (age+gender adj)	PR (multivar adj) <sup>a</sup>	n	n <sub>threshold</sub> (%)	PR (age+gender adj)	PR (multivar adj) <sup>a</sup>	n	n <sub>threshold</sub> (%)	PR (age+gender adj)	PR (multivar adj) <sup>a</sup>	n	n <sub>threshold</sub> (%)	PR (age+gender adj)	PR (multivar adj) <sup>a</sup>
<b>Depression (PHQ-9 / EST- Q2)</b>																
No confirmed diagnosis	20010	3702 (18,5)	Ref.	Ref.	-	-	-	-	-	-	-	-	-	-	-	
Ever diagnosed with COVID-19	3060	706 (23.1)	1.16 (1.09 - 1.25)	1.18 (1.10 - 1.27)	209	48 (23.0)	1.42 (1.07 - 1.89)	1.42 (1.00 - 2.02)	933	219 (23.5)	1.10 (0.97 - 1.24)	1.09 (0.97 - 1.23)	1918	439 (22.9)	1.17 (1.07 - 1.28)	1.26 (1.15 - 1.38)
Never bedridden	451	77 (17,1)	0.80 (0.65 - 0.98)	0.90 (0.74 - 1.09)	10	2 (20.0)	1.17 (0.32 - 4.28)	0.87 (0.24 - 3.13)	207	34 (16.4)	0.72 (0.53 - 0.98)	0.77 (0.56 - 1.04)	234	44 (17.5)	0.86 (0.648 - 1.13)	1.04 (0.79 - 1.35)
Bedridden 1-6 days	433	93 (21.5)	1.03 (0.87 - 1.23)	1.05 (0.88 - 1.26)	15	2 (13.3)	0.96 (0.26 - 3.31)	0.95 (0.270 - 3.35)	184	47 (25.5)	1.17 (0.92 - 1.50)	1.21 (0.95 - 1.53)	234	41 (18.8)	0.91 (0.70 - 1.17)	0.94 (0.72 - 1.23)
Bedridden 7 days or more	367	142 (38.7)	2.12 (1.87 - 2.40)	1.69 (1.47 - 1.93)	9	3 (33.3)	2.05 (0.83 - 5.09)	1.51 (0.62 - 3.70)	97	38 (39.2)	2.09 (1.64 - 2.66)	1.60 (1.22 - 2.09)	261	101 (38.7)	2.09 (1.80 - 2.43)	1.75 (1.49 - 2.05)
Non-Hospitalized	2978	671 (22.5)	1.12 (1.05 - 1.21)	1.16 (1.08 - 1.24)	208	48 (23.1)	1.43 (1.08 - 1.89)	1.43 (1.01 - 2.02)	918	214 (23.3)	1.08 (0.95 - 1.22)	1.09 (0.96 - 1.22)	1852	409 (22.1)	1.11 (1.01 - 1.22)	1.23 (1.12 - 1.35)

Hospitalized	63	28 (44.4)	2.67 (2.08 - 3.43)	1.67 (1.24 - 2.25)	1	0 (0)	-	-	12	3 (25.0)	2.75 (1.24 - 6.10)	1.44 (0.56 - 3.71)	50	25 (50.0)	2.59 (1.99 - 3.36)	1.76 (1.29 - 2.40)
<b>Anxiety symptoms (GAD-7 / ASS / EST-Q2)</b>																
No confirmed diagnosis	20 327	2755 (13.6)	Ref.	Ref.	-	-	-	-	-	-	-	-	-	-	-	
Ever diagnosed with COVID-19	3154	429 (13.6)	0.93 (0.85 - 1.02)	0.95 (0.87 - 1.05)	212	24 (11.3)	0.91 (0.61 - 1.36)	1.04 (0.66 - 1.65)	958	125 (13.0)	0.85 (0.72 - 1.01)	0.85 (0.71 - 1.00)	1984	280 (14.1)	0.94 (0.84 - 1.06)	1.03 (0.92 - 1.16)
Never bedridden	474	47 (9.9)	0.64 (0.49 - 0.88)	0.71 (0.55 - 0.92)	12	2 (16.7)	1.28 (0.33 - 5.00)	1.02 (0.31 - 3.34)	211	20 (9.5)	0.61 (0.40 - 0.92)	0.64 (0.43 - 0.96)	251	25 (10.0)	0.63 (0.44 - 0.91)	0.74 (0.52 - 1.06)
Bedridden 1-6 days	459	69 (15.0)	1.01 (0.82 - 1.25)	1.03 (0.84 - 1.28)	15	2 (13.3)	1.53 (0.42 - 5.60)	1.51 (0.37 - 6.14)	193	33 (17.1)	1.15 (0.85 - 1.55)	1.16 (0.86 - 1.56)	251	34 (13.5)	0.87 (0.64 - 1.18)	0.91 (0.67 - 1.24)
Bedridden 7 days or more	377	86 (22.8)	1.78 (1.48 - 2.13)	1.46 (1.21 - 1.77)	9	1 (11.1)	1.01 (0.16 - 6.18)	0.76 (0.17 - 3.46)	99	23 (23.2)	1.81 (1.31 - 2.49)	1.43 (1.00 - 2.06)	269	62 (23.0)	1.72 (1.38 - 2.14)	1.49 (1.19 - 1.86)
Non-Hospitalized	3070	413 (13.5)	0.91 (0.83 - 1.00)	0.95 (0.86 - 1.04)	211	24 (11.4)	0.91 (0.61 - 1.36)	1.04 (0.66 - 1.65)	943	124 (13.1)	0.85 (0.72 - 1.01)	0.85 (0.72 - 1.01)	2	265 (13.8)	0.91 (0.81 - 1.03)	1.02 (0.91 - 1.15)
Hospitalized	65	15 (23.1)	1.75 (1.16 - 2.63)	1.13 (0.74 - 1.75)	1	0 (0)	-	-	12	1 (8.3)	0.91 (0.15 - 5.49)	0.51 (0.07 - 3.87)	52	14 (26.9)	1.74 (1.14 - 2.65)	1.24 (0.80 - 1.91)
<b>COVID-19 related distress symptoms (PC-PTSD-5 / PCL-5)</b>																
No confirmed diagnosis	20 134	10 702 (53.2)	Ref.	Ref.	-	-	-	-	-	-	-	-	-	-	-	
Ever diagnosed with COVID-19	2800	1546 (55.2)	1.03 (0.99 - 1.07)	1.03 (1.00 - 1.07)	205	118 (57.6)	1.08 (0.95 - 1.23)	1.15 (0.97 - 1.36)	883	483 (54.7)	1.03 (0.97 - 1.10)	1.03 (0.97 - 1.10)	1712	945 (55.2)	1.02 (0.98 - 1.07)	1.04 (1.00 - 1.09)
Never bedridden	456	216 (47.4)	0.89 (0.81 - 0.98)	0.92 (0.84 - 1.02)	12	7 (58.3)	1.08 (0.68 - 1.72)	1.07 (0.67 - 1.72)	208	98 (47.1)	0.89 (0.77 - 1.03)	0.91 (0.79 - 1.04)	236	111 (47.0)	0.88 (0.77 - 1.01)	0.94 (0.82 - 1.07)
Bedridden 1-6 days	432	229 (53.0)	0.98 (0.90 - 1.07)	0.99 (0.91 - 1.08)	16	8 (50.0)	0.99 (0.61 - 1.61)	0.99 (0.60 - 1.63)	183	98 (53.6)	1.00 (0.88 - 1.15)	1.00 (0.87 - 1.14)	233	123 (52.8)	0.97 (0.86 - 1.10)	0.99 (0.87 - 1.12)
Bedridden 7 days or more	366	241 (65.8)	1.22 (1.13 - 1.31)	1.15 (1.07 - 1.24)	9	5 (55.6)	1.06 (0.60 - 1.88)	0.94 (0.53 - 1.69)	98	61 (62.2)	1.19 (1.03 - 1.39)	1.12 (0.96 - 1.31)	259	175 (67.6)	1.23 (1.13 - 1.34)	1.18 (1.08 - 1.28)

Non-Hospitalized	2718	1494 (55.0)	1.02 (0.99 - 1.06)	1.03 (1.00 - 1.07)	203	117 (57.6)	1.08 (0.94 - 1.23)	1.15 (0.97 - 1.36)	868	474 (54.6)	1.03 (0.96 - 1.09)	1.03 (0.97 - 1.10)	1647	903 (54.8)	1.01 (0.97 - 1.06)	1.04 (0.99 - 1.09)
	63	38 (60.3)	1.25 (1.06 - 1.47)	1.11 (0.94 - 1.32)	2	1 (50.0)	1.28 (0.37 - 4.38)	1.03 (0.38 - 2.78)	12	9 (75.0)	1.38 (0.89 - 2.15)	1.17 (0.73 - 1.90)	49	31 (63.3)	1.23 (1.03 - 1.47)	1.11 (0.93 - 1.34)
<b>Sleep quality (PSQI, EST-Q2, Single-item measures)</b>																
No confirmed diagnosis	20 508	4859 (23.7)	Ref.	Ref.	-	-	-	-	-	-	-	-	-	-	-	
Ever diagnosed with COVID-19	2891	779 (26.9)	1.10 (1.03 - 1.18)	1.11 (1.04 - 1.19)	206	56 (27.2)	1.11 (0.86 - 1.42)	1.08 (0.80 - 1.47)	909	248 (27.3)	1.14 (1.02 - 1.28)	1.14 (1.01 - 1.27)	1776	475 (26.7)	1.07 (0.99 - 1.17)	1.11 (1.02 - 1.21)
Never bedridden	474	98 (20.7)	0.83 (0.69 - 0.99)	0.89 (0.74 - 1.06)	12	1 (8.3)	0.33 (0.05 - 2.11)	0.32 (0.06 - 1.84)	211	42 (19.9)	0.82 (0.63 - 1.08)	0.84 (0.64 - 1.11)	251	55 (21.9)	0.86 (0.68 - 1.09)	0.97 (0.76 - 1.22)
Bedridden 1-6 days	461	130 (28.2)	1.14 (0.98 - 1.32)	1.15 (0.99 - 1.33)	16	4 (25.0)	1.06 (0.45 - 2.49)	1.04 (0.42 - 2.54)	193	61 (31.6)	1.32 (1.07 - 1.62)	1.31 (1.06 - 1.62)	252	65 (25.8)	1.01 (0.82 - 1.25)	1.04 (0.84 - 1.29)
Bedridden 7 days or more	380	142 (37.4)	1.56 (1.37 - 1.78)	1.40 (1.22 - 1.59)	9	4 (44.4)	1.83 (0.89 - 3.77)	1.52 (0.81 - 2.87)	100	36 (36.0)	1.57 (1.20 - 2.05)	1.40 (1.07 - 1.83)	271	102 (37.6)	1.52 (1.30 - 1.78)	1.39 (1.19 - 1.62)
Non-Hospitalized	2806	740 (26.4)	1.07 (1.00 - 1.15)	1.09 (1.02 - 1.16)	204	55 (27.0)	1.09 (0.85 - 1.41)	1.07 (0.78 - 1.45)	894	239 (26.7)	1.11 (0.99 - 1.25)	1.11 (0.99 - 1.25)	1708	446 (26.1)	1.04 (0.95 - 1.13)	1.09 (1.00 - 1.19)
Hospitalized	66	32 (48.5)	2.05 (1.62 - 2.60)	1.62 (1.27 - 2.08)	2	1 (50.0)	2.67 (0.66 - 10.74)	1.89 (0.66 - 5.46)	12	7 (58.0)	3.02 (1.96 - 4.65)	2.38 (1.68 - 3.36)	52	24 (46.2)	1.80 (1.36 - 2.39)	1.47 (1.09 - 2.00)




Bedridden 1-6 days	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Bedridden 7 days or more	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hospitalized	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Hospitalized	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<b>Sleep quality (PSQI, EST-Q2, Single-item measures)</b>																
No confirmed diagnosis	17 374	5155 (29.7)	Ref.	Ref.	-	-	-	-	-	-	-	-	-	-	-	-
Ever diagnosed with COVID-19	293	108 (36.9)	1.12 (0.96-1.32)	1.03 (0.74-1.44)	190	67 (35.3)	1.08 (0.88-1.32)	1.04 (0.75-1.45)	93	36 (38.7)	1.20 (0.92-1.57)	1.23 (0.94-1.62)	10	5 (50)	1.33 (0.61-2.87)	1.39 (0.62-3.13)
Never bedridden	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Bedridden 1-6 days	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Bedridden 7 days or more	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hospitalized	260	92 (35.4)	1.08 (0.90-1.28)	0.94 (0.64-1.37)	162	52 (32.1)	0.96 (0.75-1.22)	0.84 (0.55-1.28)	88	35 (39.8)	1.26 (0.97-1.64)	1.31 (1.00-1.71)	10	5 (50)	1.33 (0.61-2.87)	1.39 (0.62-3.13)
Hospitalized	31	14 (45.2)	1.43 (0.95-2.16)	1.28 (0.82-2.01)	26	13 (50)	1.67 (1.14-2.45)	1.55 (1.03-2.35)	5	1 (20)	0.00 (0.00-0.00)	0.00 (0.00-0.00)	-	-	-	-


		Overall							
		N = 247 519							
		Overall		Diagnosed 0-2 months ago		Diagnosed 2-6 months ago		Diagnosed 6-16 months ago	
		n	n <sub>threshold (%)</sub>	n	n <sub>threshold (%)</sub>	n	n <sub>threshold (%)</sub>	n	n <sub>threshold (%)</sub>
<b>Depression (PHQ-9 / EST-Q2)</b>									
No confirmed diagnosis		197243	22241 (11.3)	-					
Ever diagnosed with COVID-19		8150	1643 (20.2)	2547	523 (20.5)	3192	611 (19.1)	2393	507 (21.2)
Never bedridden		2585	353 (13.7)	1001	136 (13.6)	1163	155 (13.3)	421	65 (15.4)
Bedridden 1-6 days		2230	452 (20.3)	866	189 (21.8)	961	197 (20.5)	403	63 (15.6)
Bedridden 7 days or more		1030	299 (29)	272	77 (28.3)	438	112 (25.6)	320	110 (34.4)

Non-Hospitalized	6593	1373 (20.8)	1840	400 (21.7)	2645	529 (20)	2108	444 (21.1)
Hospitalized	200	69 (34.5)	64	27 (42.2)	70	13 (18.6)	66	29 (43.9)
<b>Anxiety symptoms (GAD-7 / ASS / EST-Q2)</b>								
No confirmed diagnosis	204151	15140 (7.4)						
Ever diagnosed with COVID-19	8249	1076 (13)	2631	362 (13.8)	3159	393 (12.4)	2459	317 (12.9)
Never bedridden	2602	267 (10.3)	1005	118 (11.7)	1159	107 (9.2)	438	38 (8.7)
Bedridden 1-6 days	2257	300 (13.3)	868	122 (14.1)	969	135 (13.9)	420	40 (9.5)
Bedridden 7 days or more	1047	194 (18.5)	283	48 (17)	436	77 (17.7)	328	69 (21)
Non-Hospitalized	6688	971 (14.5)	1856	313 (16.9)	2660	366 (13.8)	257,916	292 (113.2)
Hospitalized	203	39 (19.2)	68	18 (26.5)	67	6 (9)	68	15 (22.1)
<b>COVID-19 related distress symptoms (PC-PTSD-5 / PCL-5)</b>								
No confirmed diagnosis	74902	23859 (31.9)						
Ever diagnosed with COVID-19	5165	2116 (41)	1260	401 (31.8)	1840	713 (38.8)	2065	1002 (48.5)
Never bedridden	1444	416 (28.8)	482	113 (23.4)	603	175 (29)	359	128 (35.7)
Bedridden 1-6 days	1438	505 (35.1)	464	140 (30.2)	603	214 (35.5)	371	151 (40.7)
Bedridden 7 days or more	627	316 (50.4)	129	47 (36.4)	197	90 (45.7)	301	179 (59.5)
Non-Hospitalized	3774	1693 (44.9)	610	213 (34.9)	1373	568 (41.4)	1791	912 (50.9)
Hospitalized	124	55 (44.4)	24	10 (41.7)	42	16 (38.1)	58	32 (55.2)
<b>Sleep quality (PSQI, BIS, EST-Q2, Single-item measures)</b>								
No confirmed diagnosis	143265	34060 (23.8)						
Ever diagnosed with COVID-19	7510	2211 (29.4)	2100	620 (29.5)	3065	947 (30.9)	2345	644 (27.5)
Never bedridden	2361	624 (26.4)	746	198 (26.5)	1144	308 (26.9)	471	118 (25.1)
Bedridden 1-6 days	1813	538 (29.7)	630	182 (28.9)	829	265 (32)	354	91 (25.7)
Bedridden 7 days or more	1338	499 (37.3)	329	117 (35.6)	545	217 (39.8)	464	165 (35.6)
Non-Hospitalized	7107	2065 (29.1)	1981	575 (29)	2928	898 (30.7)	2198	592 (26.9)
Hospitalized	213	92 (43.2)	63	29 (46)	72	29 (40.3)	78	34 (43.6)


<sup>a</sup> Adjusted for gender, age, education, relationship status, smoking, BMI, previous psychiatric diagnosis, number of comorbidities and response period



**Figure S2: Change in mental health indicators before and after diagnosis of COVID-19, sub-analysis within cohorts with repeated measures (Median time between follow up for cohorts: C19-Resilience: 7.3 months, Omtanke2020: 1 month, CovidLife: 10 months).**



**Figure S3: Mental health indicators among individuals with a diagnosis of COVID-19 compared with individuals without COVID-19 by hospitalization status.**



**Figure S4: Change in mental health indicators from first to last measure after diagnosis of COVID-19, sub-analysis of patients with COVID-19 with repeated measures (Median time between follow up for cohorts: C19-Resilience: 12.7 months, MoBa: 10.1 months, Omtanke2020: 6 months, CovidLife: 10 months).**

**Table S5: Overall prevalence ratios of mental health indicators during the first 16 months after diagnosis of COVID-19 by illness severity**

Mental health indicator	Time since diagnosis	Time spent bedridden	PR (95% confidence interval)
<i>Depressive symptoms</i>	0–2 months	Never bedridden	0·84 (0·69-1·03)
		1–6 days	1·32 (0·85-2·04)
		> 7 days	1·66 (1·11-2·47)
	2–6 months	Never bedridden	0·76 (0·66-0·89)
		1–6 days	1·21 (0·89-1·63)
		> 7 days	1·53 (1·26-1·86)
	6–16 months	Never bedridden	0·96 (0·77-1·19)
		1–6 days	0·97 (0·78-1·21)
		> 7 days	1·60 (1·17-2·18)
<i>Anxiety symptoms</i>	0–2 months	Never bedridden	0·91 (0·78-1·07)
		1–6 days	1·01 (0·86-1·18)
		> 7 days	1·47 (0·80-2·72)
	2–6 months	Never bedridden	0·71 (0·55-0·92)
		1–6 days	1·03 (0·87-1·22)
		> 7 days	1·46 (1·20-1·79)
	6–16 months	Never bedridden	0·87 (0·60-1·26)
		1–6 days	0·82 (0·58-1·17)
		> 7 days	1·47 (1·19-1·81)

		Never bedridden	1·01 (0·86-1·19)
<i>COVID-19 related distress symptoms</i>	0–2 months	1–6 days	1·22 (1·06-1·39)
		> 7 days	1·78 (1·29-2·46)
		Never bedridden	0·82 (0·47-1·42)
	2–6 months	1–6 days	1·05 (0·86-1·27)
		> 7 days	1·17 (0·93-1·47)
		Never bedridden	0·88 (0·73-1·06)
	6–16 months	1–6 days	0·91 (0·72-1·15)
		> 7 days	1·01 (0·52-1·96)
		Never bedridden	0·94 (0·84-1·06)
<i>Poor sleep quality</i>	0–2 months	1–6 days	0·98 (0·86-1·11)
		> 7 days	1·35 (1·11-1·64)
		Never bedridden	0·98 (0·87-1·11)
	2–6 months	1–6 days	1·14 (0·96-1·35)
		> 7 days	1·48 (1·29-1·69)
		Never bedridden	1·02 (0·87-1·20)
	6–16 months	1–6 days	0·95 (0·71-1·27)
		> 7 days	1·26 (0·90-1·77)

## **References**

1. Ritchie H, Mathieu E, Rodés-Guirao L, et al. Coronavirus Pandemic (COVID-19). *Published online at OurWorldinData.org* 2020; (Retrieved from: '<https://ourworldindata.org/coronavirus>' [Online Resource]).