

# Breakthrough Infections with SARS-CoV-2 in Vaccinated Students After a Festive Event – Identification of Risk Factors

## Appendix

### Methods

#### Description of the Outbreak Setting

The venue was a private setting, with a main room connected to a larger garage where some party activities, which required large floor space, took place. The garage was less frequented. The main venue's area was  $\approx 85 \text{ m}^2$  ( $915 \text{ ft}^2$ ) with approximately  $250 \text{ m}^3$  ( $8,829 \text{ ft}^3$ ) of airspace. To avoid noise pollution in the surrounding neighborhood, windows were closed from 10 pm onwards. Assuming an equispaced distribution of all guests in the main room, the arithmetic mean of the area per person was  $0.85 \text{ m}^2$  ( $9.1 \text{ ft}^2$ ) corresponding to a personal radius of  $0.52 \text{ m}$  ( $1.7 \text{ ft}$ ).

#### Case Survey

A team of students from the Paracelsus Medical University in Nuremberg conducted telephone interviews with participants of the festive event that had given written consent. Data were collected in pseudonymized form, with individuals arbitrarily assigned unique numbers as identifiers. Data were compiled to an Excel file after plausibility check together with staff of the Institute of Clinical Hygiene, Medical Microbiology and Infectiology, Klinikum Nürnberg. Data collected pertained to the following variables:

- Sex (male/female)
- SARS-CoV-2-test before the event
- COVID-symptoms on the day of the event
- COVID-19 vaccination status (including date of vaccination and type of vaccine)
- Previously diagnosed SARS-CoV-2 infection

- Risk factors (see below)
- Arrival and departure at the event
- Contact to assumed index person (<1.5 m distance and >10 minutes or shorter/more distant conversation)
- Individual contacts to other guests (<1.5 m distance and >10 minutes)
- Shared use of drinking vessels
- SARS-CoV-2 infection after the event (date and kind of test)
- Symptoms up to 14 days after the event
- Symptoms 1 month after the event
- Need for hospitalization
- Use and alert of Coronavirus warning app

### **Definition of Risk Factors**

We defined 12 risk factors for severe progression of COVID-19 according to the „Epidemiologischer Steckbrief zu SARS-CoV-2 und COVID-19“ [https://www.rki.de/DE/Content/InfAZ/N/Neuartiges\\_Coronavirus/Steckbrief.html](https://www.rki.de/DE/Content/InfAZ/N/Neuartiges_Coronavirus/Steckbrief.html) (German only), as follows:  $\geq 70$  years of age; male sex; smoker; (severe) obesity (body-mass-index  $> 30$  kg/m<sup>2</sup>); cardiovascular diseases; chronic lung diseases; chronic kidney and liver diseases; psychiatric disorders; diabetes mellitus; cancer; immune suppression, trisomy 21.

### **Definition of Symptoms**

We defined a case with antigen point-of-care test or PCR-confirmed SARS-CoV-2 infection as symptomatic if persons reported one or more of the following symptoms 1-8 days after the event: cough, fever ( $\geq 38.0^\circ$ ), common cold (blocked/runny nose), headache, sore throat, anosmia (loss of smell), or ageusia (loss of taste), shortness of breath, gastrointestinal disorders, melalgia, chest pain, tachycardia. A follow-up was conducted with those who had tested positive after the event, one month after to inquire for long-term effects.

### **Statistics and Software**

Data were checked for consistency. Crosstabulation tables were analyzed using Pearson's Chi-Square test, Fisher's Exact test and Kruskal-Wallis test for singly ordered tables. Mann-Whitney U test and bootstrap-t tests were used to test continuously distributed variables. Logistic

regression analysis was applied to test and illustrate the effect of various risk factor on the risk of infection. All reported tests were two-sided, and p-values < 0.05 were considered statistically significant. All statistical analyses in this report were performed by use of STATISTICA 13 (Hill, T. & Lewicki, P. Statistics: Methods and Applications. StatSoft, Tulsa, OK) and PASW 24 (IBM SPSS Statistics for Windows, Version 21.0., Armonk, NY).

### **Sequencing and strain assignment**

Sequencing of SARS-CoV-2 isolates was performed on the Illumina platform. SARS-CoV-2 lineages were assigned using PANGO lineages (<https://cov-lineages.org>) (ref DOI:10.1093/ve/veab064). Characteristic mutations were furthermore identified using outbreak.info (<https://outbreak.info>). Multiple sequence alignments were performed using QIAGEN CLC Genomics Workbench 21.0.5 (<https://digitalinsights.qiagen.com>).

### **Ethics**

This outbreak investigation was approved by the Institutional Review Board of the Paracelsus Medical University in Nuremberg (IRB-2021-037). Participation in the interrogation by telephone was voluntary. Written consent was obtained by all participants. Participants were informed that data protection measures were taken according to the General Data Protection Regulation of the European Union.

### **Appendix URL 1:**

Interactive representation of interactions between participants at the event (as interrogated and in part supported by pictures taken at the event). Each guest is represented by a node and all interactions longer than 10 minutes and a distance less than 1.5 m away are represented as edges between the nodes. The color of the nodes represents test results after the event, red indicating infection, green: no infection detected, grey: unknown. The blue node represents the putative index. To create the visualization, each node started at a random position. The spacing on the diagram and the dimensions of the axes are arbitrary.

URL 1: “all\_nodes.html“

**Appendix URL 2:**

Interactive representation of interactions between individuals tested positive after the event.

Coloring scheme as in URL1.

URL 2: “only\_pos“

**Appendix Table 1. Vaccination status of participants**

Number of participants	Previous SARS-CoV-2 infection	1 <sup>st</sup> vaccination*	2 <sup>nd</sup> vaccination*	3 <sup>rd</sup> vaccination*
1	yes	JNJ	-	-
1	yes	BNT	-	-
1	yes	BNT	BNT	-
1	yes	BNT	BNT	BNT
1	no	-	-	-
1	no	JNJ	-	-
10	no	AZD	AZD	-
26	no	BNT	BNT	-
3	no	MOD	MOD	-
44	no	AZD	BNT	-
1	no	AZD	MOD	-
1	no	AZD	BNT	BNT
4	no	BNT	BNT	BNT

Table footnotes:

\*AZD, AstraZeneca; BNT, BioNTech/Pfizer; JNJ, Johnson&Johnson; MOD, Moderna

**Appendix Table 2. Selected data of infected individuals**

Individual	Vaccination status#	Duration since last vaccine dose (d)	Number of total contacts	Duration at the event (min)	Kind of contact to index person*	Day of first positive test after event	AgPO CT	PCR (cycle threshold)	Number of symptoms
1.2 (put. index)	2xBNT	178	12	270	n/a	+1	pos	pos (24)	7
1.6	2xBNT	136	12	315	++	+4	pos	pos (29)	3

1.8	2xBNT	192	11	255	++	+5	neg	pos (32)	7
1.9	2xBNT	268	18	330	++	+2	pos	neg (36)	3
1.10	2xBNT	112	18	450	++	+5	-	pos (19)	1
1.11	2xBNT	154	33	360	++	+6	neg	pos (32)	7
1.12	1xJNJ	157	19	330	++	+6	neg	pos (32)	0
1.13	2xMOD	143	16	360	++	+5	neg	pos (27)	2
2.4	AZD+ 2xBNT	99	7	330	-	+4	pos	pos (20)	6
2.5	AZD+BNT	141	19	285	+	+6	neg	pos (30)	8
2.10	none	-	4	360	++	+6	neg	pos (24)	4
2.16	AZD+BNT	136	23	300	++	+4	pos	pos (24)	4
2.17	AZD+BNT	142	50	300	++	+4	pos	pos (20)	6
2.24	2xAZD	156	52	360	++	+3	pos	pos (18)	5
2.27	AZD+BNT	141	19	360	++	+5	neg	pos (unknow n)	6
3.2	AZD+BNT	136	26	345	-	+5	-	pos (17)	7
3.11	2xAZD	165	20	330	-	+6	pos	pos (21)	3
3.22	AZD+BNT	157	38	510	-	+7	neg	pos (30)	0
3.26	AZD+BNT	127	16	210	-	+5	neg	pos (25)	5
3.27	AZD+BNT	141	11	240	-	+6	neg	pos (34)	6
3.28	AZD+BNT	141	8	240	-	+6	pos	pos (24)	2
3.30	AZD+BNT	141	33	435	-	+5	pos	-	5
3.33	AZD+BNT	141	28	390	-	+6	neg	pos (unknow n)	7
3.34	AZD+BNT	148	25	450	-	+6	neg	pos (32)	5


#AZD, AstraZeneca; BNT, BioNTech/Pfizer; JNJ, Johnson&Johnson

\* n/a: not applicable; ++: closer than 1.5m (4.9 ft) and for longer than 10 min; +: conversation shorter and/or more distant; -: no direct contact


**Appendix Table 3. Risk of infection relative to conversation with index person**

	No infection	Infection	Totals
<i>No conversation with index person</i>	48	10	58
<i>Row %</i>	82.8%	17.2%	
<i>Conversation with index person</i>	23	13	36
<i>Row %</i>	63.9%	36.1% <sup>1</sup>	
<i>Totals</i>	71	23	94

<sup>1</sup> Risk of infection is 2.1 times higher (relative risk) upon conversation with index, Fisher's Exact test (two-sided, p = 0.0497)



**Appendix Figure 1. Histogram of severity of COVID-19 sickness with infected participants, according to the number of symptoms**



Appendix Figure 2. Calculated risk of infection relative to the duration of stay at the event