

1 Sharp Increases in Drug Overdose Deaths Among High-School-Age Adolescents 2 During the US COVID-19 Epidemic and Illicit Fentanyl Crisis

3 Joseph Friedman, MPH¹, Morgan Godvin, BA², Chelsea Shover³, PhD, Joseph P. Gone, PhD⁴, Helena Hansen, MD,
4 PhD¹, David Schriger, MD, MPH⁵

5 ¹*Center for Social Medicine and Humanities, University of California, Los Angeles, California*

6 ²*Local Public Safety Coordinating Committee, Multnomah County, Oregon*

7 ³*Division of General Internal Medicine and Health Services Research, University of California, Los Angeles, California*

8 ⁴*Department of Anthropology, Department of Global Health and Social Medicine, and Native American Program, Harvard University, Cambridge,
9 Massachusetts*

10 ⁵*Department of Emergency Medicine, University of California, Los Angeles*

11

12 Abstract

13 Although overdose deaths in the US have increased exponentially for the past four decades, these shifts
14 have historically affected adults, while pediatric overdose rates remained stable. However, this may be
15 changing, given that the illicit drug supply has become increasingly hazardous in recent years, as illicitly-
16 manufactured-fentanyls (IMFs) and other synthetic opioid and benzodiazepine analogues are
17 increasingly sold as heroin and counterfeit prescription pills. We calculated drug overdose deaths per
18 100,000 population by 5-year age groups for the 2010-2021 period. For high-school-aged adolescents
19 (age 14-18), we stratified rates by race/ethnicity, census region, associated substance, and ICD-10 cause-
20 of-death intent categories. Adolescent overdose mortality saw a sharp increase between 2019 and 2020,
21 from 2.35 per 100,000 to 4.58 per 100,000, representing a 94.3% increase, the largest percent increase
22 of any 5-year age group. American Indian or Alaska Native (AIAN) adolescents, Latinx adolescents, and
23 adolescents in the West census region were disproportionately affected, overdose death rates 2.15,
24 1.31, and 1.68 times the national average in 2021, respectively. Trends were driven by fatalities
25 involving IMFs, which nearly tripled from 2019 to 2020, and represented 76.6% of adolescent overdose
26 deaths in 2021. Sharp increases in adolescent drug overdose deaths, despite flat or declining drug use
27 rates, and no increase in deaths from alcohol or most drugs, reinforce that rising fatalities are likely
28 driven by an increasingly toxic, IMF-contaminated drug supply. Rising racial disparities in overdose
29 require a prevention approach that ameliorates deep-seated social and economic inequalities as well as
30 poor access to mental and physical healthcare and social services for AIAN and Latinx adolescents. Our
31 results should also be understood in the context of rising rates of adolescent mental illness during the
32 COVID-19 pandemic. These findings highlight the urgent need for accurate, harm-reduction-oriented
33 education for early adolescents about the risks of an evolving drug supply, as well as greater access to
34 naloxone and services that check drugs for the presence of IMFs.

35 **Introduction**

36 The United States is in the midst of an unprecedented drug overdose crisis, recently crossing the
37 milestone of 100,000 deaths in a 12-month period¹. Although overdose deaths in the US have increased
38 exponentially for the past four decades, these shifts have historically affected adults, while pediatric
39 overdose rates remained stable². The illicit drug supply has become increasingly hazardous in recent
40 years, as illicitly-manufactured-fentanyls (IMFs) and other synthetic opioid and benzodiazepine
41 analogues are increasingly sold as heroin and counterfeit prescription pills³. Although the number of
42 adolescents using drugs reportedly declined in 2021⁴, media accounts of overdose deaths among high-
43 school-age adolescents have become more frequent, often implicating counterfeit pills containing IMFs.

44 **Methods**

45 We calculated drug overdose deaths per 100,000 population by 5-year age groups for the 2010-2021
46 period using data from the National Center for Health Statistics (see appendix). Records from 2020-2021
47 were provisional and may underestimate final mortality. Data from 2021 corresponded to January-May
48 and were scaled proportionally to represent annual death counts and rates. For high-school-aged
49 adolescents (age 14-18), we stratified rates by race/ethnicity, census region, associated substance, and
50 ICD-10 cause-of-death intent categories. Analyses were conducted using R version 4.0.3. This work was
51 deemed exempt from review by the University of California, Los Angeles, as it deals with anonymized
52 and publicly available records.

53 **Results**

54 Between 2010 and 2018, adolescents aged 14-18 had stable overdose mortality, substantially below the
55 high and rising rates seen among the general population (supplemental Table). However, adolescents
56 saw a sharp increase between 2019 and 2020, from 2.35 per 100,000 to 4.58 per 100,000, representing
57 a 94.3% increase (Figure 1A). This was much larger than the percent increase observed among the all-
58 age population in 2020, of 30.6% (Figure 1B). 14–18-year-old-individuals saw the highest percent
59 increase in 2020 of any 5-year age group for whom trends could be assessed (Figure 1C). Overdose
60 death rates among adolescents rose further in the first 5 months of 2021 to 5.50 per 100,000.

61

62 Stratified by race/ethnicity, American Indian or Alaska Native (AIAN) adolescents saw disproportionate
63 overdose mortality, especially after 2017. In provisional data from January-May 2021 overdose mortality
64 in this group was 2.15-fold higher than the national average (Figure 1D). Latinx adolescents were also
65 disproportionately affected, with a rate 1.31 times greater than the national average in 2021. Although

66 adolescents in all census regions saw sharp increases in 2020-2021, those in the West were
67 disproportionately affected, with an overdose death rate 1.68 times the national average. (Figure 1A).

68 Increases in adolescent overdose deaths can be attributed to rising fatalities involving IMFs and other
69 synthetic opioids, which nearly tripled from 2019 to 2020, and represented 76.6% of all deaths by 2021
70 (Figure 2A). Throughout available data, most adolescent overdose deaths were categorized as
71 unintentional, with a minority categorized as suicides, representing 84.5% and 13.4% of deaths in 2021,
72 respectively (Figure 2B).

73 **Discussion**

74 For decades, overdose mortality among high-school-aged adolescents remained stable and substantially
75 lower than rates among the general population. However, beginning in 2020 they saw the largest annual
76 percent change of any age group. The predominance of adolescent overdose deaths on the West coast
77 may be related to the region's high prevalence of counterfeit oxycodone and benzodiazepine pills
78 containing high quantities of IMFs³. Sharp increases in adolescent drug overdose deaths, despite flat or
79 declining drug use rates, and no increase in deaths from alcohol or most drugs, reinforce that rising
80 fatalities are likely driven by an increasingly toxic, IMF-contaminated drug supply.

81 We noted sharp racial disparities for AIAN adolescents, of larger magnitude than those seen among
82 adults in this population⁵. Disproportionately high trends among Latinx adolescents stand in contrast to
83 relatively lower rates among Latinx adults⁵. Rising racial disparities in overdose require a prevention
84 approach that ameliorates deep-seated social and economic inequalities as well as poor access to
85 mental and physical healthcare and social services for AIAN and Latinx adolescents.

86 Our results should be understood in the context of rising rates of adolescent mental illness during the
87 COVID-19 pandemic⁶. Although detected suicides remain a small proportion of adolescent overdose
88 deaths, sharply increasing youth suicidal ideation may be a key contextual factor. These findings
89 highlight the urgent need for accurate, harm-reduction-oriented education for early adolescents about
90 the risks of an evolving drug supply, as well as greater access to naloxone and services that check drugs
91 for the presence of IMFs.

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/).




Figure 1. Adolescent Overdose Deaths, 2010-2021

Drug overdose deaths among high-school age adolescents (age 14-18) are shown per 100,000 individuals of the same age by race/ethnicity (A) and US Census Region (B). Year-to-year percent increases are shown for adolescents aged 14-18 and compared to the all-population figures (C). The percent change in drug overdose deaths between 2019 and 2020 is shown for 5-year age groups from ages 14-18 through ages 79-83. A vertical dashed line separates trends before and during the COVID-19 pandemic (A-C). *2021 refers to January-May 2021, and rates have been annualized.




Figure 2. Substances-Involved and Intent of Adolescent Overdose Deaths, 2010-2021

Overdose deaths among high-school-aged adolescents (age 14-18) are shown per 100,000 individuals of the same age by substance used (A) and ICD-10 drug-related cause of death intent (B). A vertical dashed line separates trends before and during the COVID-19 pandemic (A). *2021 refers to January-May 2021, and rates and counts have been annualized.

**Alcohol-related overdose deaths are shown here for comparative purposes but are not included in drug overdose rates shown elsewhere in the analysis, including Figure 1.

93 **References**

- 94 1. Products - Vital Statistics Rapid Release - Provisional Drug Overdose Data. Published December 8,
95 2020. Accessed December 25, 2020. <https://www.cdc.gov/nchs/nvss/vsrr/drug-overdose-data.htm>
- 96 2. Jalal H, Buchanich JM, Roberts MS, Balmert LC, Zhang K, Burke DS. Changing dynamics of the drug
97 overdose epidemic in the United States from 1979 through 2016. *Science*.
98 2018;361(6408):eaau1184. doi:10.1126/science.aau1184
- 99 3. Ciccarone D. The rise of illicit fentanyl, stimulants and the fourth wave of the opioid overdose
100 crisis. *Current Opinion in Psychiatry*. 2021;34(4):344-350. doi:10.1097/YCO.0000000000000717
- 101 4. *Percentage of Adolescents Reporting Drug Use Decreased Significantly in 2021 as the COVID-19*.
102 National Institute on Drug Abuse; 2021:4.
103 https://www.drugabuse.gov/sites/default/files/NIDA_MTF_2021_Cleared_FINAL.pdf
- 104 5. Friedman J, Hansen H. *Black and Native Overdose Mortality Overtook That of White Individuals*
105 *During the COVID-19 Pandemic.*; 2021:2021.11.02.21265668. doi:10.1101/2021.11.02.21265668
- 106 6. *Protecting Youth Mental Health: The U.S. Surgeon General's Advisory*. Surgeon General; 2021.
107 Accessed December 19, 2021. <https://www.hhs.gov/sites/default/files/surgeon-general-youth-mental-health-advisory.pdf>

109

110 Supplemental Methods

111 Data Preparation and Definitions

- 112 1. Final drug overdose mortality and population counts were obtained for the 2010-2019 period
113 from the CDC Wonder Final Multiple Cause of Death Platform (<https://wonder.cdc.gov/mcd-icd10.html>)
114
- 115 2. Provisional drug overdose counts for January 2020 – May 2021 were obtained from the CDC
116 Wonder Provisional Multiple Cause of Death Platform ([https://wonder.cdc.gov/mcd-icd10-
117 provisional.html](https://wonder.cdc.gov/mcd-icd10-provisional.html)).
- 118 3. Drug overdose mortality rates per 100,000 individuals were calculated by 5-year age groups
119 from ages 14-18 through ages 79-83. Individuals under age 14 were excluded due to insufficient
120 overdose deaths to calculate a stable rate. Individuals over 83 were excluded due to missing
121 population numbers in the Wonder platform.
- 122 4. Additionally, for individuals aged 14-18, overdose deaths per 100,000 individuals were
123 calculated stratified by race/ethnicity, census region, substance-involved, and ICD-10 cause-of-
124 death intent categories.
- 125 5. Drug overdoses were defined by the following underlying cause of death ICD-10 codes:

X40 (Accidental poisoning by and exposure to nonopioid analgesics, antipyretics and antirheumatics)
X41 (Accidental poisoning by and exposure to antiepileptic, sedative-hypnotic, antiparkinsonism and psychotropic drugs, not elsewhere classified)
X42 (Accidental poisoning by and exposure to narcotics and psychodysleptics [hallucinogens], not elsewhere classified)
X43 (Accidental poisoning by and exposure to other drugs acting on the autonomic nervous system)
X44 (Accidental poisoning by and exposure to other and unspecified drugs, medicaments and biological substances)
X60 (Intentional self-poisoning by and exposure to nonopioid analgesics, antipyretics and antirheumatics)
X61 (Intentional self-poisoning by and exposure to antiepileptic, sedative-hypnotic, antiparkinsonism and psychotropic drugs, not elsewhere classified)
X62 (Intentional self-poisoning by and exposure to narcotics and psychodysleptics [hallucinogens], not elsewhere classified)
X63 (Intentional self-poisoning by and exposure to other drugs acting on the autonomic nervous system)
X64 (Intentional self-poisoning by and exposure to other and unspecified drugs, medicaments and biological substances)
X85 (Assault by drugs, medicaments and biological substances)
Y10 (Poisoning by and exposure to nonopioid analgesics, antipyretics and antirheumatics, undetermined intent)
Y11 (Poisoning by and exposure to antiepileptic, sedative-hypnotic, antiparkinsonism and psychotropic drugs, not elsewhere classified, undetermined intent)
Y12 (Poisoning by and exposure to narcotics and psychodysleptics [hallucinogens], not elsewhere classified, undetermined intent)
Y13 (Poisoning by and exposure to other drugs acting on the autonomic nervous system, undetermined intent)
Y14 (Poisoning by and exposure to other and unspecified drugs, medicaments and biological substances, undetermined intent)

- 126
- 127 6. Age-specific populations served as the denominator and overdose death counts served as the
128 numerator for the calculation of rates. Population data corresponding to the provisional period
129 (2020-2021) were not available, so values from 2019 were used for these years, assuming no
130 population change. Population change has been very small year-to-year, however this
131 assumption could lead to slightly elevated rates.
- 132 7. ‘Latinx’ individuals were defined as any persons for which ethnicity was given as Hispanic or
133 Latino, regardless of race. All other race/ethnicity groups were defined as individuals given to be
134 of each race, who had their ethnicity listed as “non-Hispanic.”
- 135 8. With the exception of alcohol overdose deaths, substance-specific drug overdose deaths were
136 defined by the following multiple cause of death ICD-10 codes, given that the underlying cause
137 of death was assigned to one of the aforementioned underlying cause of death codes:

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

Substance	Definition	Notes
Alcohol	Underlying cause of death in the "Alcohol poisonings (overdose) (X45, X65, Y15)" drug/alcohol induced cause set.	Not included in total drug overdose rates. Shown here for comparative purposes.
Benzodiazepines	MCD code "T42.4 (Benzodiazepines)"	
Cocaine	MCD Code "T40.5 (Cocaine)"	
Heroin	MCD Code "T40.1 (Heroin)"	
Illicit Fentanyl and Synthetics	MCD Code "T40.4 (Other synthetic narcotics)"	This category is known to represent almost entirely illicitly-manufactured fentanyl and other synthetic opioids. A very small percentage of this category now includes prescription fentanyl.
Methamphetamine	MCD Code "T43.6 (Psychostimulants with abuse potential)"	This category is known to represent almost entirely illicit methamphetamine, and is traditionally referred to as such. See: https://jamanetwork.com/journals/jamapsychiatry/fullarticle/2774859
Prescription Opioids	MCD Code "T40.2 (Other opioids)"	This category refers almost entirely to prescription opioids, such as oxycodone, not including methadone or prescription fentanyl.

138 9. Intent was defined by ICD-10 drug/alcohol induced cause of death codes.

139 10. Data were visualized using R version 4.0.3

140 Methodological Considerations

- 141 1. Records from 2020-2021 are provisional and may underestimate final mortality rates.
- 142 2. Data from 2021 represent only January-May and could overestimate or underestimate
- 143 subsequently observed trends representing the entire year.
- 144 3. Race/ethnicity can be incorrectly assigned in some death investigations. This is a well-
- 145 documented issue for American Indian and Alaska Native individuals in particular, and should
- 146 inspire caution in the interpretation of results.
- 147 4. These findings are descriptive in nature and cannot indicate causally if increases in overdose
- 148 mortality were directly linked to the COVID-19 pandemic or any other factor.
- 149 5. For drug overdoses, as with nearly all causes of death, mortality rates remain much higher for
- 150 adults than for pediatric populations. The key finding of this work include that adolescents aged
- 151 14-18 now represent the fastest growing segment of the drug overdose crisis in relative, year-
- 152 to-year percent increase space. This is particularly notable after many years of flat trends.
- 153 However, these findings should not discount the importance of drug overdose among adults,
- 154 whose overall rates still remain higher. Rather, as has been seen for numerous demographic
- 155 groups (e.g., Black communities in urban areas) early exponential increases may precede
- 156 subsequent large increases in level. Therefore, there may be an opportunity now for early
- 157 intervention among a segment of the population with rapidly growing overdose mortality. Many
- 158 of these considerations can be seen in the primary results when shown numerically, as below:

Year	Adolescents (14-18)				All Population			
	Deaths	Death Rate per 100,000	Percent Change	Absolute Change	Deaths	Death Rate per 100,000	Percent Change	Absolute Change
2010	518	2.4	5.4	0.1	38329	12.4	2.9	0.4
2011	497	2.3	-2.4	-0.1	41340	13.3	6.9	0.9
2012	429	2.0	-12.9	-0.3	41502	13.2	-0.4	0.0
2013	399	1.9	-6.4	-0.1	43982	13.9	5.2	0.7
2014	382	1.8	-4.5	-0.1	47055	14.8	6.1	0.8
2015	445	2.1	16.2	0.3	52404	16.3	10.5	1.5
2016	491	2.3	10.6	0.2	63632	19.7	20.8	3.4
2017	449	2.1	-8.7	-0.2	70237	21.6	9.5	1.9
2018	450	2.1	0.2	0.0	67367	20.6	-4.5	-1.0
2019	492	2.4	9.9	0.2	70630	21.5	4.5	0.9
2020	956	4.6	94.3	2.2	92232	28.1	30.6	6.6
2021	1147	5.5	20.0	0.9	101460	30.9	10.0	2.8

