

**Supplemental Figure 1. Original cohort characteristics prior to exclusion, N = 352**

	WHO ordinal scores <6 N =287	WHO ordinal scores 6-9 N =65	
	N (%)		p- values
Symptoms			
Cough	147 (51.2)	39 (60.0)	-
Dyspnea	150 (52.3)	45 (69.2)	
Rhinorrhea	13 (4.5)	5 (7.7)	
Anosmia	16 (5.6)	3 (4.6)	
Dysgeusia or hypogeusia	27 (9.4%)	9 (13.8)	
Sore throat	15 (5.2)	4 (6.2)	
Fever	116 (40.4)	36 (55.4)	
Hemoptysis	4 (1.4)	0 (0)	
Nausea and/or vomiting	66 (23)	10 (15.4)	
Diarrhea	54 (18.8)	20 (30.8)	
Myalgia and/or arthritis	47 (16.4)	11 (16.9)	
Rash	1 (0.3)	0 (0)	
Altered mental status	16 (5.6)	5 (7.7)	
Asymptomatic	54 (18.8)	4 (6.2)	
Comorbidities			
Cardiovascular	172 (59.9)	41 (63.1)	-
Diabetes mellitus	87 (30.3)	24 (36.9)	
Lung disease	56 (19.5)	8 (12.3)	
Renal	34 (11.8)	6 (9.2)	
Liver	23 (8)	10 (15.4)	
HIV	2 (0.7)	2 (3.1)	
Malignancy on chemotherapy	4 (1.4)	1 (1.5)	
Solid organ transplant	6 (2.1)	1 (1.5)	
Dementia	16 (5.6)	3 (4.6)	
Others	46 (16.0)	6 (9.2)	
No known PMH	35 (12.2)	8 (12.3)	
Campuses			
Angleton	22 (7.7)	4 (6.2)	-
League City	44 (15.3)	17 (26.2)	

Clear Lake	51 (17.8)	9 (13.8)	
Galveston	165 (57.5)	32 (49.2)	
Non UTMB transfers	4 (1.4)	2 (3.1)	
Missing values	1 (0.3)	1 (1.5)	
Demographics			
Sex			
Male	165 (57.5)	44 (67.7)	0.17
Female	122 (14.5)	21 (32.3)	
Code status			
Comfort care	8 (2.8)	10 (15.4)	-
DNI	18 (6.3)	6 (9.2)	
Full code	261 (90.9)	47 (72.3)	
Missing values	0 (0)	2 (3.1)	

Clinical variables **	Mean (+ SD)		p- values
BMI	31.4 (7.43)	33.9 (8.86)	0.0408
O2 saturation, %	95.6 (6.58)	93.0 (5.62)	0.00133
NLR	5.69 (5.10)	9.92 (11.2)	0.00505
CRP	4.46 (7.62)	11.5 (10.2)	< 0.001
D-dimer	1.99 (7.86)	3.62 (16.6)	0.449
qSOFA	0.286 (0.475)	0.569 (0.558)	<0.001
LDH	402 (418)	783 (559)	<0.001
	N (%)		p- values
Categorical LDH			
Normal value	197 (68.6)	19 (29.2)	<0.001
1-2 x ULN	77 (26.38)	30 (46.2)	
> 2 x ULN	13 (4.5)	14 (21.5)	

\*\* 2 subjects in ordinal score 6-9 group had consistently missed values on clinical variables

Supplemental Figure 2. Initial regression model of cohort, N=329

<b>Analysis of Maximum Likelihood Estimates</b>					
<b>Parameter</b>	<b>DF</b>	<b>Estimate</b>	<b>Standard</b>	<b>Wald</b>	<b>Pr &gt; Chi Sq</b>
			<b>Error</b>	<b>Chi-Square</b>	
<b>Intercept</b>	1	3.18E+55	1.97092E+47	1.3761	0.2408
<b>Sex</b>	1	0.918237	6.234510078	0.0022	0.9628
<b>Age</b>	1	0.985309	1.066838925	0.0521	0.8195
<b>Sex*Age</b>	1	0.991982	1.028910011	0.0797	0.7777
<b>Admission Date</b>	1	0.994326	1.004952222	1.3234	0.25
<b>BMI</b>	1	0.945066	1.119743902	0.2499	0.6171
<b>Age*BMI</b>	1	1.002062	1.001881768	1.2095	0.2714
<b>Admission SpO2</b>	1	0.930345	1.052638638	1.9798	0.1594
<b>Duration Sx</b>	1	0.947432	1.106276642	0.2857	0.593
<b>Admit NLR</b>	1	1.178686	1.064281581	6.9766	0.0083
<b>Admit Procalcitonin</b>	1	0.835939	1.566587991	0.1594	0.6897
<b>Admit D-Dimer</b>	1	0.938943	1.046341715	1.9368	0.164
<b>Admit LDH</b>	1	2.308505	1.372179017	6.993	0.0082
<b>Admit CRP</b>	1	1.807058	1.546354152	1.8428	0.1746
<b>qSOFA</b>	1	2.932338	1.429178781	9.0749	0.0026

\*Note that complete case exclusion not applied for this table

Supplemental Figure 3. DNI Sensitivity Analysis

<b>Analysis of Maximum Likelihood Estimates</b>					
<b>Parameter</b>	<b>DF</b>	<b>Estimate</b>	<b>Standard</b>	<b>Wald</b>	<b>Pr &gt; Chi Sq</b>
			<b>Error</b>	<b>Chi-Square</b>	
<b>Intercept</b>	1	77.2154835	175.3002	0.7078	0.4002
<b>Age</b>	1	0.97384804	1.051061	0.2828	0.5949
<b>BMI</b>	1	0.93304672	1.104839	0.4836	0.4868
<b>Age*BMI</b>	1	1.00215231	1.001561	1.8954	0.1686
<b>Admit SpO2</b>	1	0.92154838	1.043312	3.7208	0.0537
<b>Admit NLR</b>	1	1.07422557	1.026238	7.66	0.0056
<b>Admit D-Dimer</b>	1	0.98049278	1.01349	2.1498	0.1426
<b>Admit LDH</b>	1	3.46668914	1.288785	24.006	<.0001
<b>qSOFA</b>	1	2.37049786	1.370533	7.5001	0.0062

Supplemental Figure 4. Stratified BMI Analysis

Analysis of Maximum Likelihood Estimates					
Parameter	DF	Estimate	Standard	Wald	Pr > Chi Sq
			Error	Chi-Square	
Intercept	1	0.000554	6.30599	16.5798	<.0001
Age	1	1.032931	1.019284	2.8751	0.09
Admit NLR	1	1.190294	1.09319	3.8177	0.0507
Admit LDH	1	5.452641	1.594085	13.2276	0.0003

Lower BMI Analysis (BMI <30)

Analysis of Maximum Likelihood Estimates					
Parameter	DF	Estimate	Standard	Wald	Pr > Chi Sq
			Error	Chi-Square	
Intercept	1	7.73534434	204.95696	0.1477	0.7007
Sex	1	0.32184024	1.6594729	5.0094	0.0252
Age	1	1.05812661	1.0196914	8.401	0.0038
BMI	1	1.06822672	1.0338607	3.9238	0.0476
Admit SpO2	1	0.90429468	1.054746	3.5653	0.059
Admit NLR	1	1.16602438	1.0778842	4.1962	0.0405
Admit D-Dimer	1	0.65422758	1.215311	4.736	0.0295
Admit LDH	1	2.40873087	1.4663847	5.2744	0.0216
Admit CRP	1	2.86967849	1.708473	3.8735	0.0491

Higher BMI Analysis (BMI => 30)

Supplemental Figure 5. Ordinal Logistic Regression

Analysis of Maximum Likelihood Estimates						
Parameter		DF	Estimate	Standar	Wald	Pr > Chi Sq
				d	Chi-Square	
				Error		
Intercept	2	1	62.1903598	150.475	0.6786	0.4101
Intercept	1	1	247.769778	151.973	1.2041	0.2725
Age		1	0.95218113	1.04907	1.0425	0.3072
BMI		1	0.91805314	1.0968	0.8549	0.3552
Age*BMI		1	1.00259336	1.00148	3.0732	0.0796
Admit SpO2		1	0.91310902	1.04185	4.9188	0.0266
Admit NLR		1	1.10494991	1.03915	6.7523	0.0094
Admit LDH		1	2.49502812	1.2766	14.0199	0.0002
Admit qSOFA		1	2.08318922	1.36944	5.4482	0.0196