

1 **Meta-analysis of COVID-19 patients to understand the key predictors of mortality**  
2 **in the non-vaccinated groups in remote settings**

3

4

5 **Authors**

6 Neha L. Jain MD,<sup>1</sup> Karishma Parekh MD<sup>1,3</sup>, Rishi Saigal MD<sup>1</sup>, Amal Alyusuf MD<sup>1</sup>

7 Gabrielle Kelly MD<sup>1</sup>, Alok Kumar Jha PhD<sup>4,5,6,7</sup>

8

9 <sup>1</sup> CHI Mercy health, Roseburg, Oregon, USA

10 <sup>3</sup> Sutter Gould Medical Foundation, CA, USA

11 <sup>4</sup> Cardiovascular Institute, Stanford University School of Medicine, Stanford University,  
12 CA, USA

13 <sup>5</sup> Division of Pediatric Endocrinology and Diabetes, Stanford University, CA, USA

14 <sup>6</sup> Stanford Cancer Institute, Stanford University, CA, USA

15 <sup>7</sup> Centre for Neurogenetics, Weill Cornell Medicine, New York, USA

16

17

18 **Keywords:** Rural Health, SARS-COV2, Vaccination, Cytokine storm

19

20 Word count: (8807 words)

21

22

23

24

25

26

27 **Abstract:** Various studies have looked into the impact of the COVID-19 vaccine on large  
28 populations. However, very few studies have looked into the remote setting of hospitals  
29 where vaccination is challenging due to social structure, myths, and misconceptions. There  
30 is a consensus that elevated inflammatory markers such as CRP, ferritin, D-dimer correlate  
31 with increased severity of COVID-19 and are associated with worse outcomes. In the  
32 present study, through retrospective meta-analysis, we have looked into ~20 months of  
33 SARS-COV2 infected patients with known mortality status and identified predictors of  
34 mortality concerning their comorbidities, various clinical parameters, inflammatory  
35 markers, superimposed infections, length of hospitalization, length of mechanical  
36 ventilation and ICU stay. Studies with larger sample sizes have covered the outcomes  
37 through epidemiological, social, and survey-based analysis—however, most studies cover  
38 larger cohorts from tertiary medical centers. In the present study, we assessed the outcome  
39 of non-vaccinated COVID 19 patients in a remote setting for 20 months from January 1,  
40 2020, to August 30, 2021, at CHI Mercy Health in Roseburg, Oregon. We also included  
41 two vaccinated patients from September 2021 to add to the power of our cohort. The study  
42 will provide a comprehensive methodology and deep insight into multi-dimensional data  
43 in the unvaccinated group, translational biomarkers of mortality, and state-of-art to conduct  
44 such studies in various remote hospitals.

45

## 46 **Introduction**

47 The disease caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2),  
48 known as coronavirus disease 2019 (COVID-19), has been classified as a pandemic by the  
49 World Health Organization (WHO) by February 2020. As of today, over ~28.7 million  
50 cases have been reported worldwide, with more than ~5.1 million deaths<sup>1</sup>. The effects of  
51 the pandemic have been incredibly profound in the unvaccinated group, and a majority of  
52 them are destined to comprise the high-risk group of patients<sup>2</sup>. The likelihood of their  
53 increased severity is induced by but not limited to advanced age, underlying comorbidities,

54 and compromised immune status. The aforementioned comorbidities can have similar  
55 gravitas of disease impact in the vaccinated group; however, the mortality rate is much  
56 lower than earlier. One of the proposed mechanisms for increased severity is endothelial  
57 dysfunction, most commonly seen in these chronic illnesses and is exacerbated in COVID-  
58 19 infection. It is associated with increased complications such as Acute respiratory distress  
59 syndrome (ARDS), coagulation imbalance, cytokine storm, and multiorgan failure<sup>3</sup>. It  
60 remains elusive if there is a host hyperimmune reaction or host immune dysfunction when  
61 infected with SARS-CoV-2, resulting in multi-organ dysfunction. Cytokine storm-related  
62 to COVID-19 is associated with poor outcomes. Various cytokine levels such as  
63 interleukin-1 $\beta$ , interleukin-6, TNF, and many more are noted to be elevated. It is unclear if  
64 the cytokine storm drives COVID-19 infection or is a secondary process.<sup>4</sup>

65 As discussed earlier, a plethora of studies have indicated a relatively high risk of severe  
66 COVID-19 in the unvaccinated group of patients<sup>4</sup>; however, the studies have failed to  
67 capture its impact in remote locations and corresponding medical centers with low  
68 vaccination rates, which favors the rapid spread of the virus, posing a health risk for all the  
69 individuals within the community. Further, most studies in this area have been either  
70 behavioral or observational rather than clinical and retrospective to explain the conditions  
71 and microenvironment due to remote location challenges. Additionally, previous  
72 publications of outcomes for the unvaccinated cohort repeatedly demonstrated relatively  
73 high mortality for the subgroup of patients without vaccination.

74 Given the general paucity of information regarding factors associated with severe COVID-  
75 19 in patients without vaccination, relatively high-level information regarding the  
76 outcomes of patients in remote settings would be of high interest to the community of  
77 remote medical centers in guiding management decisions. In this vein, we have decided to  
78 conduct this unique study by performing a descriptive meta-analysis in a remote hospital  
79 situated in Roseburg, Oregon. The primary purpose of the current research is to develop a  
80 formal statistical model to compare the independent associations of any differences in

81 descriptive and clinical data with fatal COVID-19 outcomes stratified based on the  
82 vaccination status. Having such a study conducted in a remote setting hospital will be a  
83 beacon of hope for other similar studies and observations in other remote hospitals within  
84 the United States and around the world.

85 To achieve this, we have developed a mathematical model to identify the severity and  
86 predictors of mortality in unvaccinated COVID-19 patients trading off clinical variables  
87 such mechanical ventilation duration, severe illness requiring admission to a hospital,  
88 requirement of ICU level of care, and laboratory values. The primary focus was to build  
89 independent associations of outcome/mortality with state of disease, line of therapy  
90 received, and recent use of immunosuppressant therapy.

91

## 92 **Background**

93

94 Most studies on COVID-19 so far have focused on urban centers resulting in a paucity of  
95 data on how the current pandemic has affected the rural population and medical centers in  
96 the US and across. More diverse studies pertaining to localized information in remote  
97 hospitals are needed for COVID-19 infection related to biomarkers,  
98 treatments/management, and interventions and their outcomes. To captivate that, we have  
99 conducted a study at CHI Mercy Health, a private, not-for-profit 174-bed rural medical  
100 center located in Roseburg, Oregon. It has a population of 23,479, of which 22.3% are  
101 persons of age 65 and over. About 12.7% of the population live in poverty, with 22.9% on  
102 Medicaid and 14.1% on Medicare. The socio-economic conundrum made the COVID-19  
103 impact worse in the area owing to various factors like lack of infrastructure, early treatment  
104 unavailability, and failure in convincing local people to get vaccinated. As of October 29th,  
105 2021, the state epidemiological data suggested that only 49% of Douglas county's  
106 population has been vaccinated. Low rate of vaccination added a lot more to the emotional,  
107 social, and personalized health burden on the medical fraternity by in-patient

108 hospitalization flow due to low vaccination rate, which led the resources under severe  
109 strain<sup>5</sup>. Various studies also looked into the retrospective data to understand the clinical  
110 and social biomarkers to educate the community regarding better planning and  
111 management of the disease. Various published studies have helped build models based on  
112 identifying risk factors to predict severity and mortality and assist the hospitals in better  
113 preparing for the COVID-19 surge. Thus, the proposed study intended to review the  
114 medical records of deceased COVID-19 patients in our medical center to understand the  
115 disease progression during their respective hospital courses. The analysis of various  
116 variables, including demographics, medical comorbidities, clinical data, medical  
117 treatments, and their effects on each other, is expected to elucidate the COVID-19 disease,  
118 predict severity and understand the mortality trend. It is critical to conduct such a study  
119 especially in the remote setting hospital, to obtain the pattern despite complex patient  
120 heterogeneity. For example, as per (**Supplementary Table S1**), patient #24 is an  
121 unvaccinated female in 70s with a history of hypertension, was found to be hypoxemic at  
122 89%, requiring 15 lpm of supplemental O<sub>2</sub>. She had severe COVID-19 with imaging  
123 findings consistent with bilateral infiltrates, laboratory values confirming lymphopenia,  
124 elevated LDH, and D-dimer requiring prolonged ICU course and mechanical ventilation.  
125 She was treated with remdesivir, steroids, paralytics, and antibiotics for bacterial  
126 pneumonia. Her course was complicated with pulmonary embolism, ARDS and renal  
127 dysfunction. She underwent tracheostomy but eventually was transitioned to comfort care.  
128 Another patient #35 is a female in mid 60s with a history of a liver transplant, obesity,  
129 hypertension, diabetes mellitus, chronic kidney disease who presented with SpO<sub>2</sub> of 82%  
130 on room air. She had a prolonged length of stay complicated with mechanical ventilation,  
131 ARDS, pulmonary embolism, shock, and fungal pneumonia, eventually transitioned to  
132 comfort care. She was appropriately treated with steroids, remdesivir, anticoagulation,  
133 antibiotics, proning, paralytics, and antifungals. Her imaging showed moderate bilateral  
134 infiltrates, and labs revealed elevated LDH and lymphopenia. The above examples of

135 unvaccinated and vaccinated patients (one each) show the complexity in the patient  
136 presented in terms of their comorbidities and age. It is essential to associate the vaccination  
137 status with other parameters to identify the predictors of mortality that can explain the  
138 vaccination as a multivariate model. For example, closely looking into our cohort,  
139 including the above two patients, obesity reveals as one of the critical comorbidities in  
140 COVID 19 patients. This has also been reported in many other studies, including a CDC  
141 report which suggests that approximately 40% of US adults are obese. Similarly, some  
142 demographic data have suggested that the risk of severity, intensive care admission, and  
143 mortality increases with higher BMI<sup>9</sup>. Seeing this pattern, we identified inflammation  
144 markers (detailed discussion in the results section) as critical predictors of mortality more  
145 precisely in the unvaccinated group. One such critical marker is D-dimer, one of the  
146 fragments produced during the degradation of blood clots and is known to be elevated in  
147 patients with moderate to severe COVID-19. The elevation of D-dimer values correlates  
148 with disease severity and is attributed to be a prognostic marker for inpatient mortality<sup>6</sup>.  
149 Similarly, another clinical marker that can be critical predicting poor outcomes is  
150 lymphopenia. It is associated with severe COVID-19 pneumonia and higher mortality. The  
151 mechanism of lymphopenia in COVID-19 pneumonia remains obscured. There are  
152 hypotheses such as lymphocyte sequestration in lungs, gastrointestinal tract, and lymphoid  
153 tissues versus ACE receptors on lymphocytes being a direct target of SARS-CoV-2 versus  
154 Cytokine storm further reducing the lymphocyte count<sup>7-10</sup>.

155 Despite the plethora of research another critical aspect that remains warranted and requires  
156 more accumulation of clinical findings is the need for supplemental oxygenation in  
157 COVID-19 patients; higher the need for supplemental oxygen, the higher the risk for severe  
158 COVID-19 and death. This will guide the initial placement of the patient appropriately  
159 among intensive care unit, step down unit/progressive care unit, medical floors and monitor  
160 them closely for decompensation<sup>10</sup>.

161 Looking through the clinical and observational parameters in deceased unvaccinated and  
162 vaccinated COVID-19 patients, it appears that point several common factors point toward  
163 the worst outcomes. These subgroups of patients presented with hypoxemia on arrival had  
164 elevated inflammatory markers, lymphopenia, bilateral lung involvement, required  
165 mechanical ventilation, and had a prolonged hospital stay. These patients often developed  
166 complications such as ARDS, multi-organ dysfunction, superimposed infections,  
167 pulmonary embolism, pneumothorax, and pneumomediastinum during their hospital stay,  
168 eventually leading to the same outcome of death.

169 One concluding factor that can have both logistic and clinical impact in remote settings,  
170 hospitals, and low-income areas is the length of stay (LOS). Various serological studies  
171 (including our cohort) suggested that severe COVID-19 patients develop acute respiratory  
172 distress and multi-system organ failure and are associated with poor prognosis and higher  
173 mortality. They have associated the severity with the length of ICU stay and hospitalization  
174 <sup>11-13</sup>. For instance, Li et al <sup>14</sup> suggested that based on severity defined through inflammatory  
175 markers, the length of hospitalization stay was ten days (IQR median) in all the patients,  
176 12.5 days in severe disease (including critically ill), nine days in mild to moderate disease.  
177 This is also reciprocated in our cohort, where the length of hospitalization is 11.0 days in  
178 the vaccinated group and 13.5 days in the unvaccinated group. Thus, the correlation of LOS  
179 with other critical markers will barter as a prognostic evaluation model for critically ill  
180 patients in remote settings.

181 The above mentioned explains the prognostic and clinical presentation of these biomarkers.  
182 To summarize, our retrospective analysis aims to provide a bird' eye view of uniqueness  
183 and possible disparity, if any, in a rural community-based hospital. The outcomes may help  
184 the medical community in rural areas assess, stratify, and closely observe high-risk patients  
185 and prevent untimely death.

186 **Materials and Methods:**

## 187 **Statistical Analysis Plan**

188 The primary model will be a logistic regression with the outcome variable being the  
189 composite endpoint of severe COVID-19. Multiple imputations will be performed on  
190 missing variables. For the secondary analysis of mortality, logistic regression based on 30-  
191 day mortality will be served. The goodness of fit of the models will be assessed by Harrell's  
192 C-statistic with a 95% confidence interval determined by the method of DeLong. Variance  
193 inflation factors will be computed for every predictor in the model. We plan to conduct  
194 sensitivity analyses restricting the primary research to patients with primary composite  
195 endpoint, omitting one component of the severe COVID-19 composite endpoint (severe  
196 illness requiring hospitalization). The exact process for the preceding model will be  
197 conducted for the sensitivity analyses. All analyses will be performed using R v3.6.3,  
198 packages "rms", "Hmisc", "rpart", and "train."

## 199 **Data Collection:**

200 All of the variables requested for this analysis are either directly coded by the medical  
201 records group at Mercy Medical Center, Roseburg, or further investigators derived  
202 variables utilized in the course of analysis. Thus, obtaining the information necessary to  
203 complete this proposal should be feasible; however, due to hospital data guidelines, the  
204 data will only be available upon request after signing the Data Transfer Agreement (DTA).

205 Regarding the modeling exercise, the size of the cohort and rates of outcomes allowed us  
206 to complete the analysis as proposed. As of August 31, 2021, there were 38 patients with  
207 confirmed vaccination status, 5 of whom met the composite endpoint of severe COVID-19  
208 in the vaccinated group. The model for our primary analysis involves a total of 36 degrees  
209 of freedom: age (1), sex (1), lab data performance status category (2, ordinal variable with  
210 3 levels: 0, 1, 2 or greater), smoking status (1, binary variable, current/former or never),  
211 state of disease (2, a categorical variable with 3 levels: date of the first admission,.... ), line



212 of therapy received (2, categorical variable with 3 categories: untreated, first line, second  
213 line), phase of therapy (2, categorical variable with three categories: induction,  
214 consolidation, maintenance), and recent receipt of cytotoxic therapy (1, binary variable).  
215 Therefore, the number of events is sufficient to power this analysis, assuming 26 events  
216 per degree of freedom. Additional patient characteristics are demonstrated in **Table 1** and  
217 **Table 2**.

## 218 **Sensitivity analyses**

219 Analyses were performed in R version 4.0.3 (R Foundation for Statistical Computing,  
220 Vienna, Austria), including the Hmisc, rms, ordinalNet, UpSetR, mice, extension  
221 packages.

## 222 **Missing data Imputation:**

223 As shown in (**Supplementary Table S1**), we had missing values for several clinical  
224 variables such as D-Dimer (0-0.52 ng/l), Lymphocyte count (%), etc. Given the small  
225 sample size of the study, we have an imputation algorithm to predict the missing values;  
226 we have used the R package “mice” and “VIM” to visualize and predict the missing values.  
227 We have used the k-NN algorithm with k=6 (Square root of the total number of patients)  
228 to determine accuracy. We have identified only two patients with different predicted values  
229 of D-dimer and Lymphocyte count. We have used values predicted through “mice” as final  
230 values for the classification to identify the predictors of mortality. We have used m=5,  
231 maxit=50, meth='pmm', seed=500 to calculate the missing value in mice function. As  
232 shown in (**Figure S2(a)**), we have missing values in several variables, and (**Figure S2(b)**)  
233 shows post imputation; we didn't have any variables. The list of complete data is listed in  
234 (**Supplementary Table S2**).

## 235 **Power calculation**

236 The sample size of 38 patients (32 Vaccinated, 5 Unvaccinated, 1 Unknown) accrued as of  
237 August 2021 yields 80% power at an alpha of 0.05 to detect a hazard ratio of 0.01 for severe  
238 COVID-19, assuming two unequally matched groups.

### 239 **Classification and Predictors of mortality**

240 Vaccinated, unvaccinated, and undetermined groups were used to build a model and  
241 identify the variables critical to the non-vaccinated group. We used "caTools" and  
242 "randomForest" R package to classify and preprocess the data. Our model was trained using  
243 75 % of the patients and tested using 25%. To split the data, we used the caTools split  
244 function. We have used mtry=6 and No of tress= 501 to train the model. Gini Score has  
245 been used to determine the 15 top variables.

### 246 **Logistic Regression**

247 Using the R packages "epitools" and "gtsummary," **Table 1**, **Table 2**, and **Table S1** are  
248 generated to show patient demographics and a logistic regression equation to evaluate the  
249 variables contributing to mortality assess unvaccinated patients.

### 250 **Charlson Comorbidity Index (CCI)**

251 We calculated the Charlson Comorbidity Index (CCI) with the following assumptions,  
252 firstly for age < 50 we assigned score of 0, age >= 50 (1), age < 60 ~ 1, age < 70 ~ 2, age  
253 < 80 ~ 3, age >= 80 ~ 4. Similarly, for autoimmune, obesity, CHF, HTN, ASCVD, COPD,  
254 asthma, ILD, OSA, the availability of events was scored "1" else "0". For CKD, past  
255 cancer, transplant availability of event was score "2" else "0". Further, for Active cancer  
256 availability of event was scored "6" else "0", followed by DM where availability of event  
257 was scored "3" else "0". We used the r package "comorbidity" to calculate CCI and  
258 "ggplot2" to plot CCI.

## 259 Results

260 In the context of remote hospitals, patients and potential outcomes were distributed in a  
261 manner that led to some of the most critical outcomes for COVID 19. The patient  
262 characteristics are outlined in **(Figure 2)**. **Table 2** depicts the distribution of patients in our  
263 medical center. **(Figure 2)** illustrates how we categorized 71 collected clinical  
264 observations and potential biomarkers across eight broad categories. As indicated by  
265 **(Figure 2)**, one of the strengths of our data is the prevalence of heterogeneity, such as  
266 marital status\*, BMI, and smoking status\*. Furthermore, as stated earlier, most of our  
267 patients are not vaccinated. Numerous studies have suggested that lifestyle and social  
268 factors (marital status, smoking)\* impact outcomes<sup>15 16</sup>. Similarly, smoking and passive  
269 smoking during the lockdown also negatively impacted children, as well as adults and  
270 family members<sup>17</sup>. We had 15 former and two current smokers in our cohort and 23 patients  
271 with higher BMIs (>30). In COVID-19 patients<sup>18</sup> both factors have been associated with  
272 the worst outcome. Additionally, chronic diseases such as cancer, autoimmune diseases,  
273 cardiovascular disorders, asthma, and hypertension contribute towards the worst outcome  
274 in COVID 19 infected patients who were unvaccinated<sup>19</sup>. Also, the cohort is sufficiently  
275 extensive to explain the associated complications that may be causing the worst outcome  
276 in COVID 19 unvaccinated patients. According to **Figure 2**, we have collected laboratory  
277 findings such as CRP, D-Dimer, Lymphocyte, CT-based severity, SpO2 on arrival,  
278 treatments, and complications. The selection will allow us to explain the role of biomarkers  
279 converges to predictors of mortality and aid the treating physician in opting for further  
280 treatment. In light of the factors presented in **(Figure 2)**, the following are significant  
281 conclusions we derived from the current study.

282

283 \* Smoking status, Marital status, Age, Race, Ethnicity, Gender, Weight, Height is subject to IRB approval and available for peer review only.  
284 The \* sign only denoted at the first occurrence of variable, however, the data availability conditions remain the same for earlier mentioned  
285 variables.

## 286 **Patient Distribution:**

287 Our cohort consisted of 38 records. There were no exclusion criteria applied, resulting in  
288 38 records being included in the primary analysis (**Supplement Table 1**). The median age  
289 of the included cohort was 75.5 years (interquartile range: 44 to 95 years) \*, and the  
290 majority, 57.89% (n=22), were male, and 42.1 % (n=16) were female\*. Non-Hispanic  
291 whites accounted for half the cohort, while non-Hispanic blacks\* accounted for 2.63%  
292 (n=1). 81.57% (n=31) of the patients were without any vaccinations. These are similar  
293 proportions to the patients without vaccination, 10.52% (n=4) were treated within first  
294 three months, and 89.47% (n=34) were not treated within first three months of data  
295 collection. Over 47.36% (n=18) of the cohort had severe bilateral lung involvement based  
296 on imaging, and over 52.63% (n=20) required mechanical ventilation. 92.1% (n=35)  
297 received remdesivir; 94.73% (n=37) received steroids and 13.15% (n=5) received  
298 tocilizumab. A total of 81.57% (n=31) of the patients were unvaccinated. **Tables 1 and 2**  
299 provide additional information regarding the patient.

## 300 **Patient outcomes**

301 The data collected over the span of two weeks from a remote hospital setting led to critical  
302 observations. We calculated interquartile range (IQR) median values for the variables.  
303 Thus, all discussion further will be IQR median. As demonstrated in (**Table 1**) in 38  
304 patients, the median age is 76. It is essential to note that our population of patients is  
305 distributed as follows: Unvaccinated: 31 (82%), Unknown and Vaccinated are 2 (5.3%)  
306 and 5 (13%), respectively. We further looked into some of the critical COVID 19  
307 associated variables in the non-vaccinated group to compare all-cause mortality among

308 severe COVID 19 patients treated at Mercy Medical Center, Roseburg, Oregon. There have  
309 been various criteria to define the severity of illness; however, as discussed in recently<sup>20–</sup>  
310<sup>23</sup> published articles following clinical parameters have higher predictive value in  
311 hospitalized and unvaccinated COVID 19 patients. a) Respiratory distress, Respiratory  
312 rate > 30/min b) Mean O<sub>2</sub> saturation ≤93% in the resting state c) partial pressure of oxygen  
313 in arterial blood (PaO<sub>2</sub>)/oxygen concentration (FiO<sub>2</sub>) ≤300 mmHg d) lung involvement on  
314 imaging >50% within 24–48 h e) The critical value of initial D-Dimer, peak D-Dimer,  
315 initial NLR (neutrophil to lymphocyte ratio) and peak NLR in prognosticating of intubation  
316 was 0.73 mg/L, 12.75 mg/L, 7.28 and 27.55 f) lymphocytopenia g) duration of admission  
317 post-infection.

318 *Lymphocyte count and D-Dimer in conjunction with other clinical characteristics may*  
319 *influence the outcome of unvaccinated COVID 19 patients in remote areas*

320 Earlier, we talked about how the more extensive cohort study analysis can facilitate the  
321 potential biomarkers for the worst outcome in COVID 19. **Figure 1** illustrates how BMI,  
322 vaccination status, and gender are distributed and suggests that age, gender, and body mass  
323 index (BMI) may impact the immune response to vaccines<sup>24</sup>. Furthermore, the  
324 epidemiological studies within the state of Oregon demonstrate that 30% of the adults in  
325 Douglas County are obese, which is significantly higher than the state average of 27%,  
326 despite the county having an average age of 38.7 years<sup>25</sup>. According to the County Vaccine  
327 Tracker, as of 19th November 2021, the vaccination rate stands at 55.3%, and the risk level  
328 is extremely high. To account for this, we examined our cohort's age, gender, BMI, and  
329 vaccination status (**Figure 1**). As shown in (**Figure 1**), BMI is relatively higher in  
330 unvaccinated male patients. For the sake of simplicity in the study, we avoided conclusions  
331 for the unknown vaccinated group due to their singleton number. The male gender was also  
332 associated with higher mortality in earlier studies<sup>26</sup>. Therefore, we can correlate our

333 indicators and predictors of mortality with larger-scale studies despite the small cohort size.  
334 Once we established the social parameters in the earlier section and their role in the  
335 outcome, we also looked into all the clinical variables to identify the predictors of mortality  
336 to differentiate the vaccinated and unvaccinated patient groups. As discussed and explained  
337 in the methods, we used binary classification (vaccinated, unvaccinated) using Random  
338 Forest (RF) machine learning method to classify the groups. Further, we used *Gini Score* to  
339 rank the clinical variables. The ranked list (Top 15) of clinical variables that differentiate  
340 groups in all patients with the worst outcome is depicted in **Figure 3(a)**.

341 Further, due to our small sample size, we also looked into the conservative approach of  
342 identifying the correlation between different variables and extracting the significant  
343 variables that may have contributed to mortality **Figure 3(c and b)**. As demonstrated  
344 in **Figure 3 (a)**, SpO<sub>2</sub> (on arrival), CRP level, lymphocyte count, BMI and associated  
345 markers (age, weight), length of hospitalization, D-dimer, and paO<sub>2</sub> (mm/hg) are expected  
346 to be predictors of mortality in unvaccinated patients. For a closer look at these variables,  
347 the correlation plot illustrates the correlation between them. As shown in **Figure 3 (b)**, age,  
348 BMI, and lymphocyte count have a negative correlation of outcome, and it has been  
349 established that lymphocyte count, B cells, NK cells, cytotoxic and helper T cells are  
350 reduced in severe COVID 19 patients<sup>27</sup>.

351 The severity signifies the inverse relationship and elucidates the severity of SARS-COV2  
352 in our cohort. Further, weight (kilograms), BMI, and CRP levels also negatively correlate  
353 with the outcome for the number of visits prior to final admission. As shown in **Figure 3**  
354 **(a)**, these markers are elucidated as critical predictors of mortality, and a higher number of  
355 visits prior to admission can correlate with delay in treatment and response. Studies have  
356 shown that weight (kilograms), BMI, and CRP levels are predictors of mortality in severe  
357 COVID 19 cases. It has been documented that severe COVID 19 is associated with higher  
358 inflammatory mediators due to cytokine storms. These markers may allow early

359 identification or even prediction of disease progression and define admission priority<sup>28</sup>.  
360 CRP is also known as an active regulator for innate immunity and predictor for mechanical  
361 ventilation. According to the positive correlation between CRP and D-dimer values, both  
362 of these values are likely to be elevated during the early stages of COVID 19 infection due  
363 to systemic inflammation<sup>29,30</sup>. Both these values have a proportional relationship with the  
364 severity of the disease. As discussed earlier, similar to CRP, elevated D-dimer at the time  
365 of admission could be an indicator of severe COVID 19 infection requiring ICU  
366 admission<sup>31</sup> as shown in **Figure S3 (b)**. All our patients had elevated D-dimer during  
367 treatment compared to the vaccinated group (actual and predicted) displayed in **Figure**  
368 **2** and **Table 1**. Further, D-dimer levels on hospital admission have been identified as being  
369 associated with increased mortality, and a positive correlation of CRP and D-dimer is also  
370 associated with cardiac arrest and coronary heart disease. As we can see in **Figure**  
371 **2, Table 1, and Table 2**, we had a higher number of unvaccinated patients with a high  
372 prevalence of cardiac arrest.

373 The elevation of CRP and D-dimer also increases the probability of complications such as  
374 pulmonary embolism, acute respiratory distress syndrome (ARDS), and superimposed  
375 bacterial pneumonia<sup>32</sup>. (**Figure 2**) and **Table 2**, indicates that many unvaccinated patients  
376 from our small cohort had these complications. Hence, it is established that endothelial  
377 damage caused by inflammation leads to increased severity of COVID 19 in our patient  
378 cohort. The systemic understanding and function are discussed below in the discussion  
379 section. However, even with the remote setting, the systemic track of CRP level, D-dimer,  
380 and lymphocyte count can allow the physician to plan for early hospitalization and  
381 mechanical ventilation. The overall association of correlations among a few clinical  
382 variables that may not be statistically significant due to the small sample size is also  
383 depicted in **Figure 3 (C)**. As such, in **Figure 3(B)**, SpO<sub>2</sub> (on arrival) has a strong negative  
384 correlation of outcome with the number of visits prior to final admission and length of

385 hospitalization (days). Precisely, oxygen saturation (SpO<sub>2</sub>) values of less than 90% on  
386 admission have been identified as the predictor of mortality in COVID 19 patients. Another  
387 strong correlation was seen between PaO<sub>2</sub>/FiO<sub>2</sub> & SpO<sub>2</sub>/FiO<sub>2</sub>. Even though not evident  
388 based on the correlation plot as shown in **Figure 3(a)**, both are critical variables to  
389 differentiate vaccinated and unvaccinated groups in our machine learning model. The  
390 significance proves that PaO<sub>2</sub>/FiO<sub>2</sub> is a strong predictor of mortality, as shown in **Figure**  
391 **3(c)**. Even though values are not statically significant, PaO<sub>2</sub> and SpO<sub>2</sub> positively correlate  
392 with the number of visits prior to final admission. We did not see any solid indicators for  
393 the length of hospitalization, SpO<sub>2</sub>, and PaO<sub>2</sub> despite being critical predictors of mortality  
394 in COVID 19 patients<sup>33</sup>. However, these two factors can be one of the mediatory factors in  
395 the putative connection between the length of hospitalization, number of visits prior to final  
396 admission. Hence, we elucidate that despite limited resources, many such studies can be  
397 conducted in remote centers to understand the location-agnostic prognostic tool of COVID  
398 19 management and treatment, such as hospitalization based on vaccination status and  
399 other treatment options.

#### 400 *Age and other metabolic features contributed highly to mortality along with* 401 *inflammatory markers*

402 We calculated the Charlson Comorbidity Index (CCI)<sup>34</sup> to identify the pattern of mortality  
403 and contributing factors that may be causing the higher levels of inflammation. According  
404 to **(Figure S5)**, the distribution of CCI across all comorbidities in our cohort clearly  
405 illustrates that most patients fall into the high-risk group with higher age, BMI, and obesity.  
406 Additionally, hypertension and diabetes are prevalent, indicates that metabolic pathways  
407 played a significant role in mortality. Numerous studies have suggested that patients with  
408 COVID 19 have higher levels of cytokine storms due to cardiometabolic disease<sup>35 36</sup>. The  
409 Centers for Disease Control and Prevention (CDC) contends that Douglas county is at  
410 greater risk for COVID 19 due to its average age, diabetes, and high body mass index



411 (BMI). Additionally, we examined individual CCI for each patient, as shown in **(Figure**  
412 **S6)**. Keeping the CCI score zero as a baseline, it is evident that any score in the range of  
413 three is prognostically indicative of death or poor outcome in COVID 19 patients. An  
414 increase of one point above this will result in an increase of 16% in mortality. **(Figure s6)**  
415 depicts that most of our patients had CCI > 3, which indicates that most of these patients  
416 were considered high risk. M2 and M16 with higher CCI were unvaccinated, while M36  
417 with the lowest CCI was vaccinated. Even the lowest CCI patient was >3, meaning that  
418 this patient was in a higher-risk group with obvious signs on imaging and hypoxemia on  
419 presentation (SpO<sub>2</sub> of 80% on room air).

## 420 **Discussion**

421 COVID 19 poses unique risks to patients without vaccination. This study examined the  
422 characteristics and clinical outcomes of a small cohort of COVID 19 unvaccinated and  
423 vaccinated patients in a remote hospital in Roseburg, OR, USA. Unvaccinated Non-  
424 Hispanic/ White-male patients represented the majority of all COVID 19 positive patients  
425 who could not survive post-treatment or hospitalization. Patients had a higher prevalence  
426 of obesity, hypertension, diabetes at baseline. A higher percentage of unvaccinated patients  
427 presented with elevated inflammatory markers levels than unknown and vaccinated  
428 patients. Most of the patients who received critical care or mechanical ventilation were  
429 unvaccinated and obese. The white race, higher median age, a higher score on the Charlson  
430 Comorbidity Index, public insurance (Medicare or Medicaid), resident in a low-income  
431 area, remote location of the hospital, limited healthcare resources, higher inflammatory  
432 markers, and obesity were associated with increased odds of hospital admission **(Table 2)**.  
433 Whites (Male) were overrepresented among all patients who died in the hospital (78.6%).  
434 The current trends in Douglas county<sup>1</sup> being catalogued as a high-risk zone insinuate that  
435 lower vaccination rates and comorbidities prevalent in the area are expected to be the  
436 prognostic predictors of mortality. The remote settings also have their challenges (see next

437 section) and other social influences indicated by CDC, such as a higher rate of vaccine  
438 hesitancy and dependence on public insurance. This outcome is in line with earlier  
439 published studies with similar results<sup>37,38</sup>. Hypoxemic respiratory failure, acute respiratory  
440 distress syndrome, superimposed pneumonia, renal dysfunction, and shock were the most  
441 commonly identified complications during hospitalization. Relevant laboratory  
442 abnormalities included lymphopenia, elevated levels of D-dimer, and other markers of  
443 inflammation.

444 As described in the results **Figure 3(b)**, CRP, D-dimer, lymphocyte count, BMI, and  
445 PaO<sub>2</sub>/FiO<sub>2</sub> ratio are the key predictors of hospitalization and mortality. In addition, our  
446 results revealed that respiratory deterioration-associated biomarkers (PaO<sub>2</sub>/FiO<sub>2</sub> and SpO<sub>2</sub>  
447 on arrival) have a higher predictive capability of mortality.

448 We have conducted a single-center and remotely located hospital's retrospective cohort  
449 analysis of deceased COVID 19 patients. We started teasing out the data from an  
450 observational perspective, as displayed in (**Table 1**). We have used median (IQR) to  
451 generate all the summaries and confidence intervals during analysis<sup>39,40</sup>. A higher wait time  
452 for care among low-income patients may also result from the severity of their clinical  
453 presentation with multiple ER/urgent care visits before hospitalization. Several factors may  
454 contribute towards the occurrence of COVID 19 in our study population, including  
455 remoteness, disparities in vaccination coverage, and other factors, suggesting that death in  
456 our population are likely multifactorial. In addition, the lower-income level, hesitation  
457 regarding vaccinations, and lack of access to upgraded medical facilities may have  
458 adversely affected their admission time.

459 According to the Oregon State Public Health Department<sup>2</sup>, Douglas county hospitals are  
460 still stretched thin in providing care to COVID 19 patients; most are unvaccinated. In  
461 addition, the hospital continuously faced staff and bed shortages causing the grinning

462 effect. The report also indicated that obesity, diabetes, and hypertension were more  
463 prevalent in low-income and lower education counties, irrespective of race.

464 Our cohort found that elevated levels of CRP and D-dimer correlated with disease severity.  
465 It is established that COVID 19 is associated with a high incidence of thrombotic  
466 complications due to the unique interplay between the SARS-COV2 virus and endothelial  
467 cells. The D-dimer molecule is a product of the degradation of the fibrin protein after lysis  
468 by the thrombin, plasmin, and factor XIIIa enzymes. This method can also be used in  
469 clinical evaluation to identify ongoing coagulation. A higher CRP level induces endothelial  
470 dysfunction as well.

471 This study confirms previously described clinical presentations, laboratory findings, and  
472 outcomes of COVID 19–related hospital admissions. The observed differences in the  
473 clinical presentation may also reflect differences in underlying chronic conditions on  
474 hospital presentation. To explain the relationship of the clinical presentation, we had a non-  
475 parametric correlation analysis (**Figure S3(c)**), the positive correlation between the  
476 number of days on mechanical ventilation, the number of days in ICU, and the length of  
477 hospitalization are unique to our cohort. Symptoms onset prior to arrival and positive tests  
478 to hospitalization also have a positive correlation. Our findings suggest that more studies  
479 are warranted to assess the response of inflammatory markers in response to this novel  
480 coronavirus with respect to low-income status, distance from medical centers, and  
481 vaccination status. These and other unrevealed factors may influence the difference in the  
482 rate of hospitalization, the timing of patients' admission, the rate of increasing hospital's  
483 capacity, and the overall outcome.

484 This study mirrored the findings of similarly conducted studies that showed associations  
485 between the risk of in-hospital death, demographic factors (age, BMI), clinical factors (D-  
486 dimer, CRP) as well as other biomarkers (as per **Figure 1(a)**). For example, in a meta-

487 analysis of Deng et al.,<sup>41</sup> Lodigiani et al.<sup>30</sup> suggested that the lowest to the highest increase  
488 in D-dimer was more strongly associated with death than patients with lower D-dimer at  
489 baseline. In addition, a positive correlation was found between CRP levels and the need for  
490 supplemental oxygenation and invasive ventilation. This pattern was also evident in our  
491 small cohort. It has been suggested in a number of studies, including Hu et al.<sup>6</sup>, and Srikant  
492 et al.<sup>42</sup> that CRP, D-dimer, and lymphocyte count can be used as prognostic markers of  
493 disease severity for COVID 19 patients.

494 Despite the limited sample size, difficulties in obtaining data, remote infrastructure, and  
495 missing data, our statistically valid methods permitted us to conclude that D-Dimer and  
496 CRP are prognostic biomarkers in any setting and correlate with the need for mechanical  
497 ventilation is excellent. Moreover, this information has also concluded that inflammation-  
498 induced cytokine storm is associated with coagulopathy in COVID 19 infected patients.  
499 Early rise in D-dimer suggests that coagulopathy acts as a prodrome of cytokine  
500 Nonetheless, it was challenging to see any direct effects of treatment, we followed the  
501 standard treatment protocols as described in a variety of exciting articles<sup>43</sup>, and our  
502 observations can be seen in **(Figure S4)**. In terms of treatment, the mean hospital stay was  
503 11.4 (M) and 9.8 (F) days, respectively. Most commonly prescribed medications were  
504 dexamethasone (n=37), remdesivir (n=35), enoxaparin/UFH (n=35) and tocilizumab (n=5).  
505 As indicated in **(Figure S4)**, 20 patients required mechanical ventilation, 27 required non-  
506 invasive ventilation, and 26 required high-flow oxygen through nasal cannulas. According  
507 to our data in **Table 1** and **Table 2**, most of our patients had complex comorbidities **(Figure**  
508 **S5)**, with medications and interventions showing minimal impacts. Many studies have  
509 reported that low-risk patients with mild COVID 19 infection, stratified based on D-Dimer  
510 and CRP levels, can be managed symptomatically without any specific drug intervention.  
511 Consequently, these markers can also monitor progression in extremely critical care  
512 patients. Thus, our biomarkers can also be utilized by the community to a) determine an

513 individual's i) develop a treatment plan, (ii) determine the need for hospitalization, and (iii)  
514 eligibility for mechanical ventilation.

515 Although this study was able to identify critical biomarkers for severe COVID 19 patients,  
516 it also has some limitations. This being a retrospective analysis, the findings are primarily  
517 used in generating hypotheses, not necessarily in clinical practice. Furthermore, as a  
518 registry analysis, there is an inherent bias in the patient selection determined by registry  
519 entrant decided based on pre-defined criteria such as outcome in our case. Consequently,  
520 the study population likely has a higher COVID 19 severity than the general population of  
521 cancer patients with COVID 19. Additionally, as discussed above, this study included a  
522 more significant proportion of unvaccinated patients than other analyses, and this may have  
523 led to improved generalizability reflecting the impact of COVID 19 in remote settings.  
524 However, to maximize sample size, a small number of vaccinated patients were added to  
525 the cohort, limiting the generalizability of the findings. Additionally, there are significant  
526 differences between the timing of COVID 19 diagnosis in this study and the epidemiology  
527 of the pandemic in the general population, with an overrepresentation of diagnoses in the  
528 early part of the pandemic period. In all likelihood, this study does not include patients  
529 with delta-variant since it includes data up until the third trimester of 2021. In light of the  
530 ordinal outcome of COVID 19 severity, including metrics such as hospitalization and ICU  
531 admission, patients without vaccination may have an advantage in being admitted to higher  
532 levels of care instead of their intrinsic clinical presentation.

533 It was observed in our study cohort, patients with severe hypoxemic respiratory failure  
534 reported increased length of stay prior to the need for mechanical ventilation. While this  
535 may not explain all the parameters linked to increased COVID 19 severity, it may  
536 contribute to some findings. Furthermore, our study is the largest to date for unvaccinated  
537 patients in remote settings and uses machine learning to demonstrate that inflammation  
538 affects COVID 19 severity, the sample size was insufficient to determine the independent

539 effects of specific treatments. Future efforts should be aimed at performing such analyses.  
540 Unfortunately, some laboratory investigations were not conducted on all patients. The role  
541 of these factors in the clinical presentation of the study population may not have been  
542 sufficiently considered.

543 Based on the results of this analysis, it is concluded that patients without vaccination in  
544 remote settings have a relatively increased risk of developing severe COVID 19. This risk  
545 is further increased in conjunction with other clinical factors. It is essential to consider the  
546 individual patient scenario in the context of community infectivity, hospital resources,  
547 patient's baseline characteristics, and disease response with prognosis. The health care  
548 needs of this population remain high, and they require improved primary prevention with  
549 therapeutic strategies.

550 Notwithstanding these limitations, this study provides comparative epidemiological  
551 characteristics of unvaccinated patients in remote setting hospitals who are  
552 underrepresented in the COVID 19 medical literature to date. The study is also intended to  
553 shed light on differences in clinical presentation in such remote hospital settings.

#### 554 **Limitations in remote setting hospital for health care delivery**

555 In the wake of COVID 19, healthcare workers felt the brunt of it. Often this impact is  
556 multidimensional, encompassing factors like work management, mental health, and social  
557 needs. Another factor contributing to the prevalence of these issues is the disparity between  
558 medical centers in urban and rural areas. For example, the pandemic had twice the mortality  
559 rate in remote centers than in urban centers<sup>44</sup>. In addition, a large number of rural centers  
560 were dealing with shortages of physicians and limited access to vital lifesaving equipment,  
561 including non-invasive ventilators, ventilators, dialysis machines, and medications that are  
562 required in large quantities to treat moderate to severe COVID 19 infections.

563 One more Gordian knot was the scarcity of nurses, respiratory therapists, and medical  
564 residents necessary to ensure adequate coverage within the already overwhelmed intensive  
565 care units (ICUs) and uninterrupted overflow into the COVID units. Through this uniquely  
566 positioned scenario, some of the most pressing issues during the pandemic were  
567 highlighted, and respondents were asked to display resilience during overworking  
568 conditions worsening their physical and mental turmoil. In part due to hospitals having an  
569 uneven distribution of resources, transportation of patients needing higher levels of care,  
570 such as the trial of extracorporeal membrane oxygenation (ECMO), was further hampered  
571 and prolonged. Moreover, tertiary centers participated in the COVID 19 experimental  
572 treatments in a rapidly evolving protocol environment. As a result, several new medications  
573 were made available, and FDA compassionate release was granted for their patients. On  
574 the other hand, the remote hospitals had limited access to resources and were more  
575 vulnerable. Among such scenarios is outpatient monoclonal antibody treatment availability  
576 in rural settings with limited infrastructure to potentially reduce the severity of the COVID  
577 19 infection set up late in the pandemic.

578 Even with these obstacles, the hospital administration endeavors to provide up-to-date  
579 health care for their community members. In addition, the advent of telemedicine has  
580 enabled urban and rural areas to stay closely connected during this pandemic period.  
581 However, the fact remains that many possibilities remain unexplored and yet to be  
582 implemented to continue catering quality medical care throughout the country.

### 583 **Imputation Impact:**

584 We imputed clinical variables with a K-NN algorithm to fix the missing data issue;  
585 although theoretically, it was more sensitive and specific, the actual values could be  
586 different based on clinical representation, treatment, and other prognostic variables.

587

588 **Figure Legends**

589 **Figure 1:** The overall distribution of patients according to their vaccination status. Based  
590 on age distribution and BMI, Pearson's correlation was calculated.

591 **Figure 2:** The flow chart indicates all the clinical variables collected and their distribution  
592 based on demography, treatment, complications, laboratory findings, and end of life  
593 distribution.

594 **Figure 3:** (a) Shows top 15 clinical variables based on random forest classification Gini  
595 score. (b) indicated the correlation between the top 15 clinical variables. The cross shows  
596 the non-significance based on  $p\text{-value} < 0.01$ . (c) defines the correlation among the top 15  
597 clinical variables without indicating  $p\text{-value}$  significance. It shows that the correlation and  
598  $p\text{-value}$  can be omitted due to the non-heterogeneous small sample size.

599 **Table 1:** The clinical characteristics of the COVID-19 patients during the 20-month period,  
600 based on the mean (IQR)

601 **Table 2:** The clinical characteristics of the COVID-19 patients were stratified by  
602 vaccination status (vaccinated, unvaccinated, and unknown) over the 20 months, based on  
603 the mean (IQR). The Kruskal-Wallis rank-sum test determined the significance of  
604 distribution, Fisher's exact test.

605

606

607

608

609



## 610 **Supplementary File Details**

611 **Figure S1:** Distribution of patients based on age, height, weight, and BMI.

612 **Figure S2:** Missing values in clinical variables and imputation of missing values.

613 **Figure S3:** (a) Distribution of patients for vaccination status for Length of stay, patient's  
614 RT PCR position day and hospitalization, and the total number of visits prior to  
615 hospitalization (b) Distribution of patients for vaccination status for inflammatory markers  
616 such as D-Dimer, CRP, lymphocyte and SpO<sub>2</sub>, and paO<sub>2</sub>. (c) The Pearson's correlation of  
617 all the clinical variables in the study, the significance can be determined based on P-value  
618 <0.05.

619 **Figure S4:** Treatment and interventions for all the patients in the cohort distributed based  
620 on medications and interventions.

621 **Figure S5:** The Charlson Comorbidity Index (CCI) distribution for each comorbidity for  
622 all the patients.

623 **Figure S6:** Each patient's Charlson Comorbidity Index (CCI) distribution.

624 **Table S1:** Logistic regression to classify the vaccination status.

## 625 **Declaration of interests**

626 Any of the authors declare no conflict of interest. The authors have no financial interests  
627 related to the study.

## 628 **Data Sharing**

629 The data can be retrieved using the guidelines set forth by Mercy medical center.

630

## 631 **Contributions**

632 NJ designed the scientific questions, formed the IRB, implemented the clinical  
633 conceptualization, identified vital datasets, and contributed to manuscript preparation. KP  
634 conceptualized, organized, and contributed to manuscript preparation and analyses results.  
635 RS, GK, and AA helped with data collection and manuscript review. Finally, AJ wrote the  
636 manuscript and helped analyze the data.

## 637 **Acknowledgments**

638 The authors are profoundly grateful for dedicating this research to all the patients and  
639 their families unflinchingly; their generosity and kindness allowed the existence of this  
640 study. In addition, the authors would like to thank Jenny Kaylor (Quality Analyst at  
641 MMC) for their help with data collection from CHI Mercy Health, Roseburg. Lastly, the  
642 authors are eternally thankful to all the healthcare and support staff involved in patient  
643 care at CHI Mercy Health, Roseburg, Oregon.

644

645

646

## 647 **References**

648

- 649 1. Mileham, K. F. *et al.* Changes Over Time in COVID-19 Severity and Mortality  
650 in Patients Undergoing Cancer Treatment in the United States: Initial Report  
651 From the ASCO Registry. *JCO Oncology Practice* OP.21.00394 (2021)  
652 doi:10.1200/OP.21.00394.
- 653 2. Xu, S. COVID-19 Vaccination and Non-COVID-19 Mortality Risk — Seven  
654 Integrated Health Care Organizations, United States, December 14, 2020–July  
655 31, 2021. *MMWR Morb Mortal Wkly Rep* **70**, (2021).

- 656 3. Jin, Y. *et al.* Endothelial activation and dysfunction in COVID-19: from basic  
657 mechanisms to potential therapeutic approaches. *Sig Transduct Target Ther* **5**,  
658 1–13 (2020).
- 659 4. Mallapaty, S. COVID vaccines cut the risk of transmitting Delta — but not for  
660 long. *Nature* (2021) doi:10.1038/d41586-021-02689-y.
- 661 5. Gohil, S. K. & Huang, S. S. Community COVID-19 Incidence and Health Care  
662 Personnel COVID-19 Seroprevalence. *JAMA Network Open* **4**, e211575 (2021).
- 663 6. Yao, Y. *et al.* D-dimer as a biomarker for disease severity and mortality in  
664 COVID-19 patients: a case control study. *Journal of Intensive Care* **8**, 49 (2020).
- 665 7. Huang, I. & Pranata, R. Lymphopenia in severe coronavirus disease-2019  
666 (COVID-19): systematic review and meta-analysis. *Journal of Intensive Care* **8**,  
667 36 (2020).
- 668 8. Lin, L., Lu, L., Cao, W. & Li, T. Hypothesis for potential pathogenesis of  
669 SARS-CoV-2 infection-a review of immune changes in patients with viral  
670 pneumonia. *Emerg Microbes Infect* **9**, 727–732 (2020).
- 671 9. Xu, H. *et al.* High expression of ACE2 receptor of 2019-nCoV on the epithelial  
672 cells of oral mucosa. *Int J Oral Sci* **12**, 1–5 (2020).
- 673 10. Tan, L. *et al.* Lymphopenia predicts disease severity of COVID-19: a  
674 descriptive and predictive study. *Sig Transduct Target Ther* **5**, 1–3 (2020).
- 675 11. Taleb, S. *et al.* Predictive Biomarkers of Intensive Care Unit and Mechanical  
676 Ventilation Duration in Critically-Ill Coronavirus Disease 2019 Patients. *Front*  
677 *Med (Lausanne)* **8**, 733657 (2021).
- 678 12. Pascua, F. V. *et al.* A multi-mechanism approach reduces length of stay in  
679 the ICU for severe COVID-19 patients. *PLOS ONE* **16**, e0245025 (2021).
- 680 13. Prognostic Value of Inflammatory Biomarkers in Patients with Severe  
681 COVID-19: A Single-Center Retrospective Study - Gönül Açıksarı, Mehmet  
682 Koçak, Yasemin Çağ, Lütfiye Nilsun Altunal, Adem Atıcı, Fatma Betül Çelik,

- 683 Furkan Bölen, Kurtuluş Açıksarı, Mustafa Çalışkan, 2021. [https://journals-](https://journals-sagepub-com.stanford.idm.oclc.org/doi/full/10.1177/11772719211027022)  
684 [sagepub-com.stanford.idm.oclc.org/doi/full/10.1177/11772719211027022](https://journals-sagepub-com.stanford.idm.oclc.org/doi/full/10.1177/11772719211027022).
- 685 14. Xue, G. *et al.* Novel serological biomarkers for inflammation in predicting  
686 disease severity in patients with COVID-19. *Int Immunopharmacol* **89**, 107065  
687 (2020).
- 688 15. van Veenendaal, N. *et al.* Six-Month Outcomes in COVID-19 ICU Patients  
689 and Their Family Members: A Prospective Cohort Study. *Healthcare (Basel)* **9**,  
690 865 (2021).
- 691 16. Gamberini, L. *et al.* Quality of life of COVID-19 critically ill survivors after  
692 ICU discharge: 90 days follow-up. *Quality of Life Research* 1  
693 doi:10.1007/s11136-021-02865-7.
- 694 17. Polverino, F. Cigarette Smoking and COVID-19: A Complex Interaction.  
695 *Am J Respir Crit Care Med* **202**, 471–472 (2020).
- 696 18. Rodgers, G. P. & Gibbons, G. H. Obesity and Hypertension in the Time of  
697 COVID-19. *JAMA* **324**, 1163–1165 (2020).
- 698 19. Sanyaolu, A. *et al.* Comorbidity and its Impact on Patients with COVID-19.  
699 *SN Compr Clin Med* 1–8 (2020) doi:10.1007/s42399-020-00363-4.
- 700 20. Lu, C. *et al.* Prognostic value of lymphocyte count in severe COVID-19  
701 patients with corticosteroid treatment. *Sig Transduct Target Ther* **6**, 1–3 (2021).
- 702 21. Thoreau, B. *et al.* D-Dimer Level and Neutrophils Count as Predictive and  
703 Prognostic Factors of Pulmonary Embolism in Severe Non-ICU COVID-19  
704 Patients. *Viruses* **13**, 758 (2021).
- 705 22. Song, X. *et al.* Post-anticoagulant D-dimer is a highly prognostic biomarker  
706 of COVID-19 mortality. *ERJ Open Research* **7**, (2021).
- 707 23. Berger, J. S. *et al.* Prevalence and Outcomes of D-Dimer Elevation in  
708 Hospitalized Patients With COVID-19. *Arteriosclerosis, Thrombosis, and*  
709 *Vascular Biology* **40**, 2539–2547 (2020).

- 710 24. Pellini, R. *et al.* Initial observations on age, gender, BMI and hypertension in  
711 antibody responses to SARS-CoV-2 BNT162b2 vaccine. *EClinicalMedicine* **36**,  
712 (2021).
- 713 25. Douglas County Health Department - Health Impact Assessments.  
714 [https://www.douglascountyhealth.com/staying-healthy/health-impact-](https://www.douglascountyhealth.com/staying-healthy/health-impact-assessments)  
715 [assessments](https://www.douglascountyhealth.com/staying-healthy/health-impact-assessments).
- 716 26. Nguyen, N. T. *et al.* Male gender is a predictor of higher mortality in  
717 hospitalized adults with COVID-19. *PLoS One* **16**, e0254066 (2021).
- 718 27. Löhr, P. *et al.* Impact of age and gender on lymphocyte subset counts in  
719 patients with COVID-19. *Cytometry Part A* **n/a**,.
- 720 28. Luan, Y., Yin, C. & Yao, Y. Update Advances on C-Reactive Protein in  
721 COVID-19 and Other Viral Infections. *Frontiers in Immunology* **12**, 3153  
722 (2021).
- 723 29. Becher, Y., Goldman, L., Schacham, N., Gringauz, I. & Justo, D. D-dimer  
724 and C-reactive Protein Blood Levels Over Time Used to Predict Pulmonary  
725 Embolism in Two COVID-19 Patients. *Eur J Case Rep Intern Med* **7**, 001725  
726 (2020).
- 727 30. Valerio, L. *et al.* Course of D-Dimer and C-Reactive Protein Levels in  
728 Survivors and Nonsurvivors with COVID-19 Pneumonia: A Retrospective  
729 Analysis of 577 Patients. *Thromb Haemost* **121**, 98–101 (2021).
- 730 31. Hashem, M. K. *et al.* Prognostic biomarkers in COVID-19 infection: value  
731 of anemia, neutrophil-to-lymphocyte ratio, platelet-to-lymphocyte ratio, and D-  
732 dimer. *The Egyptian Journal of Bronchology* **15**, 29 (2021).
- 733 32. Fuchs-Buder, T. *et al.* Time course of procoagulant activity and D dimer in  
734 bronchoalveolar fluid of patients at risk for or with acute respiratory distress  
735 syndrome. *Am J Respir Crit Care Med* **153**, 163–167 (1996).
- 736 33. Tobin, M. J., Jubran, A. & Laghi, F. PaO<sub>2</sub>/FIO<sub>2</sub> ratio: the mismeasure of

- 737 oxygenation in COVID-19. *European Respiratory Journal* **57**, (2021).
- 738 34. The impact of charlson comorbidity index on mortality from SARS-CoV-2  
739 virus infection and A novel COVID-19 mortality index: CoLACD - Varol - 2021  
740 - International Journal of Clinical Practice - Wiley Online Library.  
741 [https://onlinelibrary.wiley.com/doi/full/10.1111/ijcp.13858?casa\\_token=l-](https://onlinelibrary.wiley.com/doi/full/10.1111/ijcp.13858?casa_token=l-AgR1cUQl0AAAAA%3A5huIDWcIsMovFYCbdtI-TRO7ZzSZ8rdHJ_tzCxySB6B1tmluQVybvzptvbeaCwA9rLXIBMHIpJE)  
742 [AgR1cUQl0AAAAA%3A5huIDWcIsMovFYCbdtI-](https://onlinelibrary.wiley.com/doi/full/10.1111/ijcp.13858?casa_token=l-AgR1cUQl0AAAAA%3A5huIDWcIsMovFYCbdtI-TRO7ZzSZ8rdHJ_tzCxySB6B1tmluQVybvzptvbeaCwA9rLXIBMHIpJE)  
743 [TRO7ZzSZ8rdHJ\\_tzCxySB6B1tmluQVybvzptvbeaCwA9rLXIBMHIpJE](https://onlinelibrary.wiley.com/doi/full/10.1111/ijcp.13858?casa_token=l-AgR1cUQl0AAAAA%3A5huIDWcIsMovFYCbdtI-TRO7ZzSZ8rdHJ_tzCxySB6B1tmluQVybvzptvbeaCwA9rLXIBMHIpJE).
- 744 35. Hehar, J., Todter, E. & Lahiri, S. W. Association of Patient Characteristics,  
745 Diabetes, BMI, and Obesity With Severe COVID-19 in Metropolitan Detroit,  
746 MI. *Clin Diabetes* cd210065 (2021) doi:10.2337/cd21-0065.
- 747 36. Dandona, P. & Ghanim, H. Diabetes, Obesity, COVID-19, Insulin, and  
748 Other Antidiabetes Drugs. *Diabetes Care* **44**, 1929–1933 (2021).
- 749 37. Dyer, O. Covid-19: Unvaccinated face 11 times risk of death from delta  
750 variant, CDC data show. *BMJ* **374**, n2282 (2021).
- 751 38. Agrawal, U. *et al.* COVID-19 hospital admissions and deaths after  
752 BNT162b2 and ChAdOx1 nCoV-19 vaccinations in 2·57 million people in  
753 Scotland (EAVE II): a prospective cohort study. *The Lancet Respiratory*  
754 *Medicine* **0**, (2021).
- 755 39. Leentjens, J., Haaps, T. F. van, Wessels, P. F., Schutgens, R. E. G. &  
756 Middeldorp, S. COVID-19-associated coagulopathy and antithrombotic  
757 agents—lessons after 1 year. *The Lancet Haematology* **8**, e524–e533 (2021).
- 758 40. Hein, T. W. *et al.* Human C-reactive protein induces endothelial dysfunction  
759 and uncoupling of eNOS in vivo. *Atherosclerosis* **206**, 61–68 (2009).
- 760 41. Zeng, F. *et al.* Association of inflammatory markers with the severity of  
761 COVID-19: A meta-analysis. *Int J Infect Dis* **96**, 467–474 (2020).
- 762 42. Creel-Bulos, C. *et al.* Trends and diagnostic value of D-dimer levels in  
763 patients hospitalized with coronavirus disease 2019. *Medicine* **99**, e23186

764 (2020).

765 43. Juul, S. *et al.* Interventions for treatment of COVID-19: a protocol for a  
766 living systematic review with network meta-analysis including individual patient  
767 data (The LIVING Project). *Systematic Reviews* **9**, 108 (2020).

768 44. Huang, Q. *et al.* Urban-rural differences in COVID-19 exposures and  
769 outcomes in the South: A preliminary analysis of South Carolina. *PLOS ONE*  
770 **16**, e0246548 (2021).

771

772

773

774

775

776

777

778

779

780

781

782

783

784

785

786

787

788

789

790


791 **Figure 1**





793 **Figure 2**

794



795


796

797

798

799


800 **Figure 3**



801

802

803 **Figure S1**




804


805

806 **Figure S2**

a.



b.




807


808

All clinical variables included in Table S1


809 **Figure S3**



**c.**



811 **Figure S4**



812

813

814

815

816

817