

1 Supporting Information 1

2

3 Metagenomic analysis of antimicrobial resistance genes in wastewaters in Benin and Burkina Faso indicates a  
4 serious health risk from untreated hospital wastewaters in low-income countries

5

6 Melina A. Markkanen<sup>a#</sup>, Kaisa Haukka<sup>a, f</sup>, Katariina M. M. Pärnänen<sup>a</sup>, Victorien Tamegnon Dougnon<sup>b</sup>, Isidore  
7 Juste O. Bonkougou<sup>c</sup>, Zakaria Garba<sup>d</sup>, Halidou Tinto<sup>d</sup>, Anniina Sarekoski<sup>a</sup>, Antti Karkman<sup>a</sup>, Anu Kantele<sup>e, f\*</sup>,  
8 Marko P. J. Virta<sup>a#\*</sup>

9

10 <sup>a</sup> Department of Microbiology, University of Helsinki, Finland

11 <sup>b</sup> Research Unit in Applied Microbiology and Pharmacology of Natural Substances, Polytechnic School of  
12 Abomey-Calavi, University of Abomey-Calavi, Benin

13 <sup>c</sup> Department of Biochemistry and Microbiology, University Joseph KI-ZERBO, Ouagadougou, Burkina Faso

14 <sup>d</sup> Clinical Research Unit Nanoro, Institute for Research in Health Sciences, National Center for Scientific and  
15 Technological Research, Burkina Faso

16 <sup>e</sup> Meilahti Infectious Diseases and Vaccine Research Center, MeiVac, Department of Infectious Diseases,  
17 University of Helsinki and Helsinki University Hospital, Helsinki, Finland

18 <sup>f</sup> Human Microbiome Research Program, Faculty of Medicine, University of Helsinki, Helsinki, Finland

19

20 Corresponding authors:

21 # Marko P. J. Virta, [marko.virta@helsinki.fi](mailto:marko.virta@helsinki.fi)

22 # Melina A. Markkanen, [melina.markkanen@helsinki.fi](mailto:melina.markkanen@helsinki.fi)

23 \* Equal contribution

24

25 Table of Contents:

26 Figure S1. Sample collection in Benin and Burkina Faso.....2

27 Figure S2. Examples of sampling sites.....2

28 Figure S3. PCoA plots for resistomes, taxonomical compositions, and mobilomes.....4

29 Table S1. Significance levels for ordination plots.....5

30 Figure S4. Unique and shared ARGs between hospital wastewaters in Benin, Burkina Faso, and Finland....5

31 Figure S5. Relative abundance of clinically relevant species.....6

32 Figure S6. Multidrug-resistance fragments putatively encoded by class 1 integron gene cassettes.....6

33 Figure S7. Contigs harboring mobile colistin resistance gene *mcr-5.1* .....7

34 References .....8

35


36 Pages: 9

37 Figures: 7

38 Tables: 1

39

40




42

43 **Figure S1.** Sample collection in Benin and Burkina Faso. Sample collection sites are denoted with red dots on the  
 44 map. The climate in the coastal region of Benin is hot and humid. The average daytime temperature in Cotonou  
 45 varies between 28–32 °C and humidity between 77–84 %, with an annual rainfall of 1,308 mm. Burkina Faso is  
 46 characterized by a dry tropical climate. The average daytime temperature in Ouagadougou varies between 31–  
 47 39 °C, and the annual rainfall is 745 mm. During the sampling period in November/December, it typically rains  
 48 only a little in Cotonou and not at all in Ouagadougou.

49



50 A) B) C)  
 51 D) E) F)  
 52 G) H)  
 53 **Figure S2.** Examples of sampling sites. The samples collected in Benin included A) HWW from an open cesspit  
 54 next to the wall at the hospital B (sample BH29); B) moist soil between two septic tanks (samples BH31 and  
 55 BH34) at hospital B; C) the central septic tank collecting wastewater from different sections of hospital C (sample  
 56 BH44); D) water puddle surrounding the surgery room septic tank at hospital C (sample BH48); E) water from a  
 57 tank used for washing hands at hospital C (sample BH52); F) tap water in a distant village in Benin (sample  
 58 BSE93); and G) well water 100 m away from hospital A in Benin (sample BH11). H) In Burkina Faso, the HWW  
 59 septic tank is shown for hospital F (sample BFH1). Photos by K. Haukka, A. Sarekoski and A. Butcher.




60

61 **Figure S3.** PCoA plots describe the ordinations based on A) resistomes, B) taxonomical compositions, and C)  
 62 mobilomes in HWWs from Benin, Burkina Faso, and Finland. Countries are coded as colors, and the specific  
 63 hospitals are marked with a corresponding capitalized letter. Confidence ellipses represent 90 % confidence levels.  
 64

65 **Table S1.** Significance levels for ordination plots. The significance was measured using the adonis test for equal-  
 66 sized sample groups (n=8/country) of randomly selected samples. The triplicated tests were mainly in line with  
 67 each other, suggesting that sample selection did not significantly affect the results. The significance levels are  
 68 denoted as follows: ‘\*\*\*’ ≤ 0.001; ‘\*\*’ = 0.001-0.01; ‘\*’ = 0.01-0.05; ‘.’ = 0.05-0.1; ‘ ’ = 0.1-1.


ARG, Horn									
sample set #1			sample set #2			sample set #3			
$R^2$	$p$ value	sig. level	$R^2$	$p$ value	sig. level	$R^2$	$p$ value	sig. level	
Benin-Finland	0.29492	0.001	***	0.30982	0.002	**	0.33668	0.001	***
Benin-Burkina Faso	0.09321	0.121		0.17575	0.002	**	0.16236	0.007	**
Burkina Faso-Finland	0.24544	0.001	***	0.25117	0.001	***	0.25582	0.001	***
Taxonomical composition, Horn									
sample set #1			sample set #2			sample set #3			
$R^2$	$p$ value	sig. level	$R^2$	$p$ value	sig. level	$R^2$	$p$ value	sig. level	
Benin-Finland	0.17282	0.002	**	0.20603	0.001	***	0.18496	0.002	**
Benin-Burkina Faso	0.08476	0.159		0.14289	0.007	**	0.08541	0.007	
Burkina Faso-Finland	0.16481	0.001	***	0.14788	0.006	**	0.16983	0.002	**
MGEs, Horn									
sample set #1			sample set #2			sample set #3			
$R^2$	$p$ value	sig. level	$R^2$	$p$ value	sig. level	$R^2$	$p$ value	sig. level	
Benin-Finland	0.31312	0.002	**	0.34187	0.001	***	0.34196	0.001	***
Benin-Burkina Faso	0.06114	0.497		0.10201	0.098	.	0.0994	0.094	.
Burkina Faso-Finland	0.20934	0.001	***	0.22952	0.001	***	0.20678	0.002	**

69  
70  
71



72  
73  
74  
75


**Figure S4.** Unique and shared ARGs between hospital wastewaters in Benin, Burkina Faso, and Finland. A total of 2,331 ARGs were detected among 67 hospital wastewater samples in the three countries. The number of unique ARGs was substantially higher ( $n = 333$ ) in Burkina Faso compared to Benin ( $n = 59$ ) or Finland ( $n = 126$ ).



76

77 **Figure S5.** Relative abundance of clinically relevant species. The heatmap shows the relative abundances of  
 78 species listed in the ESKAPEEc group (*Enterococcus faecium*, *Staphylococcus aureus*, *Klebsiella pneumoniae*,  
 79 *Acinetobacter baumannii*, *Pseudomonas aeruginosa*, *Enterobacter* spp., and *Escherichia coli*) defined by de  
 80 Angelis and colleagues (1). In addition, *Acinetobacter* species with known intrinsic resistance to carbapenems  
 81 according to the Beta-lactamase Database (2) are shown. Of these, species with a total relative abundance higher  
 82 than 0.05 in the samples are shown.

83



84

85

86 **Figure S6.** Examples of assembly graphs showing multidrug-resistance fragments putatively encoded by class 1  
 87 integron gene cassettes in samples A) BH02, B) BH48, and C) BFH19.


88

89

90




91



92




93



94



95




96

97



98




99

100



101



102 **Figure S7.** Comparative analysis of contigs that harbor mobile colistin resistance gene *mcr-5.1* in four different  
 103 metagenomic samples. Visualizations of the contigs were created using anvi'o (v7) metagenomics workflow for  
 104 single profiles (3). Blue arrows indicate putative encoding genes. The gene names were determined based on  
 105 BLAST search results. The recruited reads are shown as a dark grey histogram with an overlay line (pink) denoting  
 106 the GC content. Single nucleotide variation is shown on top of the panel with colorful vertical lines. Inverse repeat  
 107 sequences of Tn3-like transposase sequences (38 bp) are shown with dark pink (IRL: 5'  
 108 GGGGTCGTCTCAGAAAACGGAAAAATCGTACGCTAAG 3'(4)) and green triangles (IRR: 5'  
 109 CTTAGCGTACGATTTTTTCCGTTTTCTAGACGACCCC 3' (4)). The genetic environments are shown for  
 110 samples A) BH02 B) BH48, C) BFH26 D) BFH41, and E) FH9.

111

112 References

- 113 1. de Angelis G, Fiori B, Menchinelli G, Tiziana D'inzeo &, Flora &, Liotti M, Morandotti GA,  
 114 Sanguinetti M, Posteraro B, Spanu T. 2007. Incidence and antimicrobial resistance trends in  
 115 bloodstream infections caused by ESKAPE and Escherichia coli at a large teaching hospital in Rome, a  
 116 9-year analysis (2007-2015). *European Journal of Clinical Microbiology & Infectious Diseases*  
 117 37:1627–1636.
- 118 2. Naas T, Oueslati S, Bonnin RA, Laura Dabos M, Zavala A, Dortet L, Retailleau P, Iorga BI, emy  
 119 Bonnin RA. 2017. Beta-lactamase database (BLDB)-structure and function. *Journal of Enzyme*  
 120 *Inhibition and Medicinal Chemistry* 32:917–919.
- 121 3. Murat Eren A, Kiefl E, Shaiber A, Veseli I, Miller SE, Schechter MS, Fink I, Pan JN, Yousef M,  
 122 Fogarty EC, Trigodet F, Watson AR, Esen ÖC, Moore RM, Clayssen Q, Lee MD, Kivenson V, Graham  
 123 ED, Merrill BD, Karkman A, Blankenberg D, Eppley JM, Sjödin A, Scott JJ, Vázquez-Campos X,  
 124 McKay LJ, McDaniel EA, R Stevens SL, Anderson RE, Fuessel J, Fernandez-Guerra A, Maignien L,  
 125 Delmont TO, Willis AD. 2021. Community-led, integrated, reproducible multi-omics with anvi'o.  
 126 *Nature Microbiology* 6:3–6.
- 127 4. Borowiak M, Fischer J, Hammerl JA, Hendriksen RS, Szabo I, Malorny B. 2017. Identification of a  
 128 novel transposon-associated phosphoethanolamine transferase gene, *mcr-5*, conferring colistin  
 129 resistance in d-tartrate fermenting *Salmonella enterica* subsp. *enterica* serovar Paratyphi B. *Journal of*  
 130 *Antimicrobial Chemotherapy* 72:3317–3324.

131