## Working from home due to the COVID-19 pandemic abolished the sleep disturbance vulnerability of late chronotypes relieving their predisposition to depression


Federico Salfi<sup>1</sup>, Aurora D'Atri<sup>1</sup>, Giulia Amicucci<sup>1,2</sup>, Lorenzo Viselli<sup>1</sup>, Maurizio Gorgoni<sup>2</sup>, Serena Scarpelli<sup>2</sup>, Valentina Alfonsi<sup>2</sup>, and Michele Ferrara<sup>1\*</sup>

<sup>1</sup>Department of Biotechnological and Applied Clinical Sciences, University of L'Aquila, L'Aquila, Italy <sup>2</sup>Department of Psychology, Sapienza University of Rome, Rome, Italy

> \*Corresponding author Prof. Michele Ferrara, Ph.D. Department of Biotechnological and Applied Clinical Sciences University of L'Aquila Via Vetoio (Coppito 2) 67100 Coppito (AQ) Italy <u>michele.ferrara@univaq.it</u>

## **Supplemental materials**

Two additional moderation models were tested using model 1 of PROCESS macro (version 3.5) for SPSS (version 22.0).<sup>40</sup> We assumed that working modality (office working, remote working) moderated the effect of chronotype on bedtime and get-up time (Figure S1). Both the models included the covariance of age and gender.




**Figure S1.** The two additional moderation models tested (M4, M5). Chronotype is hypothesized to predict bedtime and get-up time, while the working modality (office working, remote working) moderated this relationship.

Abbreviations: MEQr, Morningness-Eveningness Questionnaire-reduced version.

The regressions on bedtime and get-up time were significant (Model 4:  $R^2 = .247$ , F = 56.852, p < .001; Model 5:  $R^2 = .274$ , F = 65.685, p < .001). The covariate effect of "gender" emerged significant for the bedtime variable (B = 14.51 min, t = 2.630, p = .009), indicating that female respondents went to bed earlier. Age was a significant predictor of get-up time (B = -1.58 min, t = -8.509, p < .001), indicating that older age was associated with earlier get-up time.

The conditional direct effects at both values of the moderator (office working, remote working) were significant in both the models. In particular, MEQr score predicted significantly bedtime (office working: B = -9.80 min, t = -10.029, p < .001; remote working: B = -7.92 min, t = -11.154, p < .001) and get-up time (office working: B = -8.50 min, t = -9.317, p < .001; remote working: B = -6.05 min, t = -9.221, p < .001), confirming the tendency to delayed sleep time of evening-type people in both the working modality conditions. The "working modality" moderator was significant in both the models (M4: B = 51.55 min, t = 2.682, p = .008; M5: B = 69.27 min, t = 3.874, p = .001), showing that people working from home went to bed and got-up later. Finally, the significant interaction between "working modality" and MEQr score

was limited to the get-up time variable (M4: B = -1.87 min, t = -1.564, p = .118; M5: B = -2.30 min, t = -2.130, p = .033), highlighting that the relationship between chronotype and rising time was stronger in those who worked from home than in the office working condition (Figure S2).



**Figure S2.** Simple slope analysis of the interaction between MEQr scores and working modality [office working (blue line), remote working (green line)] on get-up time (hh:mm). Gray bands identify chronotypes according to the validated cut-off scores.

*Abbreviations:* ET, evening type; NT, neither type; MT, morning type; MEQr, Morningness-eveningness questionnaire-reduced version.