

## **Plasma p217+tau vs NAV4694 amyloid and MK6240 tau PET across the Alzheimer continuum.**

Vincent Doré<sup>1,2</sup>, James D. Doecke<sup>\*3</sup>, Ziad S. Saad<sup>4</sup>, Gallen Triana-Baltzer<sup>4</sup>, Randy Slemmon<sup>4</sup>, Natasha Krishnadas<sup>2</sup>, Pierrick Bourgeat<sup>3</sup>, Kun Huang<sup>2</sup>, Samantha Burnham<sup>1</sup>, Christopher Fowler<sup>5</sup>, Stephanie R. Rainey-Smith<sup>6</sup>, Ashley Bush<sup>5,7</sup>, Larry Ward<sup>5</sup>, Jo Robertson<sup>5</sup>, Ralph N. Martins<sup>6,8,9</sup>, Colin L. Masters<sup>5</sup>, Victor L. Villemagne<sup>2,10</sup>, Jurgen Fripp<sup>3</sup>, Hartmuth C. Kolb<sup>4</sup>, Christopher C. Rowe<sup>2,5,7</sup>

*1 The Australian e-Health Research Centre, CSIRO, Melbourne, Victoria, Australia;*

*2 Department of Molecular Imaging & Therapy, Austin Health, Melbourne, Victoria, Australia;*

*3 The Australian e-Health Research Centre, CSIRO, Brisbane, Queensland, Australia;*

*4 Neuroscience Biomarkers, Janssen Research and Development, La Jolla, CA, USA;*

*5 The Florey Institute of Neuroscience and Mental Health, Melbourne, Victoria, Australia;*

*6 Edith Cowan University, Perth, Australia;*

*7 Florey Department of Neuroscience and Mental Health, The University of Melbourne, Melbourne, Victoria, Australia;*

*8 Centre for Healthy Ageing, Health Futures Institute, Murdoch University, Murdoch, WA, Australia*

*9 McCusker Alzheimer's Research Foundation, Nedlands, Perth, Australia;*

*10 Department of Psychiatry, University of Pittsburgh, Pittsburgh, PA, USA;*

\*Vincent Doré and James Doecke contributed equally to this work

Corresponding author: Vincent Doré

ORCID: 0000-0002-8051-0558

Address: Dept of Molecular Imaging & Therapy, Austin Hospital, LVL1 Harrold STOKES Block, 145 Studley Road, 3084 Heidelberg, Victoria Australia

Tel: +61 (0)3 9496 3321

Email: Vincent.Dore@csiro.au

Word count: 3,500

## ABSTRACT.

**INTRODUCTION:** We evaluated a new Simoa plasma assay for phosphorylated tau at aa217 enhanced by additional ptau sites (p217+tau).

**METHODS:** Plasma p217+tau levels were compared to <sup>18</sup>F-NAV4694 amyloid-beta (A $\beta$ ) PET and <sup>18</sup>F-MK6240 tau PET in 174 cognitively impaired (CI) and 223 cognitively unimpaired (CU) participants.

**RESULTS:** Compared to A $\beta$ - CU, the plasma levels of p217+tau increased two-fold in A $\beta$ + CU and 3.5-fold in A $\beta$ + CI. In A $\beta$ - the p217+tau levels did not significantly differ between CU, MCI or dementia. P217+tau correlated with A $\beta$  centiloids  $\rho=0.67$  (CI 0.64; CU 0.45) and tau SUVR<sub>MT</sub>  $\rho=0.63$  (CI 0.69; CU 0.34). Area under curve (AUC) for AD vs A $\beta$ - CU was 0.94, for AD vs other dementia was 0.93, for A $\beta$ + vs A $\beta$ - PET was 0.89 and for tau+ vs tau- PET was 0.89.

**DISCUSSION:** Plasma p217+tau levels elevate early in the AD continuum and correlate well with A $\beta$  and tau PET.

### **Keywords:**

*Plasma p-tau217; Tau imaging; A $\beta$ -amyloid imaging; Alzheimer's disease; positron emission tomography; blood biomarkers, blood diagnostic for Alzheimer's disease, phosphorylated tau, amyloid plaque, paired helical filaments*

### **Declarations**

**Funding.** The research was supported by the Australian Federal Government through NHMRC grants APP1132604, APP1140853 and APP1152623 and by a grant from Enigma Australia. Janssen Pharmaceuticals paid a commercial data access fee to the AIBL study of aging.

**Conflicts of interest/Competing interests.** Christopher C. Rowe has received research grants from NHMRC, Enigma Australia, Biogen, Eisai and Abbvie. He is on the scientific

advisory board for Cerveau Technologies and consulted for Prothena, Eisai, Roche and Biogen Australia. Victor Villemagne is and has been a consultant or paid speaker at sponsored conference sessions for Eli Lilly, Life Molecular Imaging, GE Healthcare, Abbvie, Lundbeck, Shanghai Green Valley Pharmaceutical Co Ltd, and Hoffmann La Roche. Ziad Saad, Gallen Triana-Baltzer, Randy Slemmon and Hartmuth Kolb are employees of Janssen R&D. The other authors did not report any conflict of interest.

**Ethics approval and Consent to participate** This study was approved by the Austin Health Human Research Ethics Committee (HREC/18/Austin/201)

**Consent for publication** All participants gave written consent for publication of de-identified data.

**Availability of data and material:** The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

**Acknowledgement:** Some of the data used in the preparation of this article was obtained from the Australian Imaging Biomarkers and Lifestyle flagship study of aging (AIBL), funded by the Commonwealth Scientific and Industrial Research Organization (CSIRO), National Health and Medical Research Council (NHMRC), and participating institutions. AIBL researchers are listed at [www.aibl.csiro.au](http://www.aibl.csiro.au). The authors thank all participants who took part in the study, as well as their families.

### **Research in Context:**

1. Systematic review: The authors reviewed the literature using PubMed, meeting abstracts and presentations. Plasma phospho-tau measures compare well to PET and post-mortem across the continuum of AD but accuracy varies across ptau target sites and assay methods. There are no reports comparing PET to plasma assays targeting multiple sites of tau phosphorylation as typically found in AD. The p217+tau assay targets p217 with binding enhanced by phosphorylation at additional sites such as aa212.
2. Interpretation: Plasma p217+tau elevates early and correlates with both A $\beta$  and tau as measured by PET indicating that tau phosphorylation is an early event in AD and occurs with A $\beta$  deposition. Plasma p217+tau measurement should assist both selection for trials and diagnosis of AD.
3. Future directions: Further validation studies, head-to-head comparison to other assays, assessing the influence of co-morbidities and the ability to measure change in brain A $\beta$  and tau levels are required.

## Introduction

Assessing brain levels of A $\beta$ -amyloid (A $\beta$ ) plaque and tau aggregation has a central role in the selection of individuals for clinical trials and assessing the efficacy of therapeutic compounds. It may also aid earlier and more accurate diagnosis of Alzheimer's disease (AD) [1].

Despite the extremely low concentration of A $\beta$  and tau in plasma, progress in detection methodologies has recently allowed reliable measurement of plasma A $\beta_{42/40}$  and phosphorylated tau (P-tau). Measurement of plasma tau phosphorylated at threonine 181 (P-tau181) was the first to demonstrate good accuracy [2-4]. P-tau181 can differentiate AD from cognitively unimpaired (CU) subjects [5] and from other dementias [6]. P-tau181 was also a good predictor of elevated brain A $\beta$  and tau as measured by PET [7]. Subsequently, plasma P-tau217 and P-tau231 have also shown strong association with AD pathology with accuracy comparable to CSF and PET measures [8, 9] with some evidence of superior performance to plasma P-tau181 [10, 11].

More recently a high sensitivity Simoa assay has been developed using a capture antibody (pT3) that was raised against tau in paired helical filaments (PHF) of AD brain [12,13]. The core requirement for this antibody binding is phosphorylation at aa217 (p217) with enhanced binding when other nearby phosphorylated sites are present, predominantly at aa212 [12]. The recognised epitope is thus referred to as "p217+tau" and its measurement in plasma has demonstrated good concordance with CSF markers of AD in a validation cohort of 227 subjects with Area Under the Curve (AUC) of 0.90 vs CSF A $\beta_{42/40}$  and 0.95 vs CSF P-tau181 [13].

Plasma p217+tau has not been compared to PET measures.  $^{18}\text{F}$ -NAV4694 and  $^{18}\text{F}$ -MK6240 are PET tracers for imaging A $\beta$  plaque and PHF 3R/4R tau aggregates respectively. These

recent generation tracers have high target to background ratios giving a wide dynamic range that may improve sensitivity for detection of low levels of A $\beta$  and tau [14, 15].

In this study we evaluated the performance of the novel blood-based biomarker p217+tau against latest generation A $\beta$  and tau PET agents in participants of the Australian Imaging, Biomarkers and Lifestyle study of aging (AIBL) and the Australian Dementia Network (ADNeT) trial screening program.

## **METHODS**

### **Participants**

Three hundred and ninety-seven participants in the AIBL and ADNeT cohorts had both <sup>18</sup>F-NAV4694 A $\beta$  PET and <sup>18</sup>F-MK6240 tau PET. The AIBL cohort recruitment and evaluation is detailed elsewhere [16]. The ADNeT cohort all had mild cognitive impairment (MCI) or mild dementia and were evaluated as per AIBL. A multi-disciplinary panel, blind to all imaging and blood results, classified all participants as cognitively unimpaired (CU) or cognitively impaired (CI) with MCI, Alzheimer's disease (AD) or non-AD dementia. A diagnosis of CU required performance within 1.5 standard deviations of the published norms for their age group on neuropsychological assessment. A diagnosis of MCI [17] or dementia [18] were assigned according to internationally agreed criteria. For this study, only individuals with a clinical diagnosis of AD and a positive A $\beta$  PET scan were classified as AD. Those with dementia but a negative A $\beta$  scan were classified as non-AD dementia.

Institutional ethical review committees approved the AIBL and ADNeT studies and written informed consent was obtained from all participants.

### **Image Acquisition**

A 20-minute A $\beta$  PET acquisition was conducted 50 minutes post-injection of 200 MBq of <sup>18</sup>F-NAV4694. On a separate day a 20-minute Tau PET acquisition was conducted 90 minutes post-injection of 185 MBq of <sup>18</sup>F-MK6240.

A $\beta$  PET scans were spatially normalized using CapAIBL [19] and the standard Centiloid (CL) method was applied [20, 21]. A CL value of 25 was selected to determine a positive A $\beta$  scan (A $\beta$ +) [22-24]. <sup>18</sup>F-MK6240 tau PET scans were spatially normalized using a CapAIBL PCA-based approach [25] and scaled using the cerebellar cortex as the reference region. <sup>18</sup>F-MK6240 SUVR values were estimated in two composite regions of interest (ROI); a mesial temporal ROI (Me) comprising entorhinal cortex, hippocampus, parahippocampus and amygdala; and a meta temporal ROI (MT) composed of the Me ROI plus inferior and middle temporal and fusiform gyri. We also estimated <sup>18</sup>F-MK6240 SUVR in Braak stages IV, V and VI with ROIs derived from the Freesurfer Desikan-Killiany Atlas (see Supplementary Figures 3-4). In the CU group, we additionally estimated <sup>18</sup>F-MK6240 SUVR in the entorhinal cortex, amygdala, hippocampus and inferior temporal cortex separately. The threshold for tau positivity was the 95<sup>th</sup> percentile of the A $\beta$ - CU group in each composite ROI and individual mesial temporal regions.

### **Plasma Analysis and p217+tau assay design**

Fasted blood sampling was performed 3.3 $\pm$ 6 months from the time of A $\beta$  PET scan and 1.7 $\pm$ 2 months from the tau scan. Plasma from K<sub>2</sub>-EDTA tubes (7.5 mL S-monovette 01.1605.008, Sarstedt) containing pre-added prostaglandin E1 (33 ng/mL of whole blood, Sapphire Biosciences) to prevent platelet activation, a potential source of peripheral amyloid- $\beta$ , was centrifuged at room temperature at 200g for 10 mins to collect platelet-rich plasma, and then at 800g for 10 mins to provide plasma that was snap frozen within 2 hr of collection was stored in vapour phase liquid nitrogen prior to shipping on dry ice from Australia to Janssen R&D, La Jolla, CA, USA. Plasma ptau217+ assay was performed on a Single MOlecular Array (Simoa) HD-X platform blinded to all subject data using the technique previously described [13]. Data analysis was then performed by AIBL investigators.

## Statistical analysis

Demographic and clinical characteristics were compared between A $\beta$  status within clinical classifications using standard Generalised Linear Modelling (GLM) for quantitative features and the Chi-square test for categorical comparisons. Correlation between the plasma p217+tau and CL or tau SUVR was performed using Spearman's rank-based correlation (Spearman's Rho [ $\rho$ ]). Receiver Operating Characteristic AUC analyses were used to explore the discriminative performance of p217+tau. Optimal p217+tau thresholds for detecting A $\beta$ + and tau+ PET were calculated both using Youden's Index and two standard deviations above the mean p217+tau levels in A $\beta$ - CU group. Presented are AUC values with 95% confidence intervals (shown in square brackets), sensitivity, and specificity with 95% confidence intervals and positive and negative predictive values (PPV, NPV, not disease prevalence adjusted). Model based performance of p217+tau accounting for age, gender and *APOE*  $\epsilon$ 4 allele status was done using GLM. AUC values between models were compared using DeLong's method. Confidence intervals were computed using bootstrap. False discovery rate (FDR) corrected P-values lower than 0.05 were considered significant. Vertex and voxel-based correlations between p217+tau and A $\beta$  and tau images were performed using Spearman's rank-based correlation.

## Results

### 1. Demographic and Clinical Characteristics

Three participants had a low CL but high Braak stage V/VI tau on <sup>18</sup>F-MK6240 PET suggesting that, similar to some PiB case reports [26], the A $\beta$  PET were falsely negative.

These three subjects all showed clearly elevated plasma p217+tau levels and were kept in the analysis but are displayed with a diamond shape in the scatter plots.

Table 1 presents demographic and clinical characteristic. Of the 397 participants, 56% were cognitively unimpaired (CU), 22% had MCI, and 21% had dementia of which 82% in the dementia group were A $\beta$ +. Overall, 170 (43%) had A $\beta$ + PET and 143 (36%) had high tau PET in the meta temporal region (T<sub>MT+</sub>). Participants with dementia were approximately 4 years younger than MCI and CU.

Stratification by A $\beta$  status within each clinical classification is provided in Supplementary Table 1.

## **2. Plasma p217+tau Levels in Clinical Groups by A $\beta$ PET Status**

Figure 1A shows the concentration of plasma p217+tau in the different clinical groups, subdivided into A $\beta$ - and A $\beta$ +. The mean p217+tau concentration in CU A $\beta$ + participants (162.7fg/ml (SD: 85.8)) was approximately twice that of the CU A $\beta$ - participants (71.2fg/ml (SD: 46.6),  $p < 10^{-18}$ ). Mean p217+tau concentration in MCI A $\beta$ + (252.0 fg/ml (SD: 141.0)) were not significantly different to those with A $\beta$ + dementia (AD) (259.3 fg/ml (SD: 147.7)) but both groups showed significantly higher concentration than in both CU A $\beta$ + participants and A $\beta$ - CU, MCI and dementia participants. Levels of p217+tau in the MCI A $\beta$ - (82.1 fg/ml (SD:55.8)) and the A $\beta$ - non-AD dementia (93.4 fg/ml (SD: 45.6)) were not significantly higher than CU A $\beta$ - participants.

When subdividing the CL scale into 5 levels (Figure 1B), the ptau217+tau concentration was progressively higher from 25 CL. This progressive increase remained in those who were tau PET negative (see Supplementary Figure 1). Figure 1C shows the Centiloid level with



increasing p217+tau level and figure 1D shows the probability of having a positive A $\beta$  scan vs p217+tau level.

### **3. Correlations of plasma p217+tau with A $\beta$ PET and Tau PET**

Vertex based analysis (Figure 2 and supplementary figures 6-10) and scatter plots (Figures 3-5) demonstrate moderate to strong correlations between plasma p217+tau and A $\beta$  and tau PET. The correlation in all subjects between p217+tau and A $\beta$  PET CL gave a Spearman's  $\rho$  of 0.67 ( $p < 10^{-53}$ ). The correlation was significantly stronger in CI than in CU (CI:  $\rho = 0.64$ ,  $p < 10^{-21}$ ; CU:  $\rho = 0.45$ ,  $p < 10^{-12}$ ; Z-score difference 2.6) while Figure 2 shows the regional correlation matches the early and predominant locations of A $\beta$  accumulation in AD.

The correlation between p217+tau and tau SUVR was moderate to strong in the meta temporal ROI (Spearman  $\rho = 0.63$ ,  $p < 10^{-46}$ ) and the mesial temporal ROI (Me) (Spearman  $\rho = 0.60$ ,  $p < 10^{-40}$ ) but was progressively lower when the ROI was restricted to higher Braak stage regions. In CI subjects, the correlation was again strongest in the meta temporal ROI ( $\rho = 0.69$ ). In CU the correlation was lower; meta temporal ROI ( $\rho = 0.34$ ); mesial temporal ROI ( $\rho = 0.33$ ) while Figure 2 shows the regional correlation matches the early and predominant regions of tau accumulation in AD. The correlations persisted between ptau217+ and tau SUVR<sub>MT</sub> when analysis only included the A $\beta$ + participants ( $\rho = 0.52$  in CI;  $\rho = 0.37$  in CU). There was no correlation between p217+tau and tau PET in A $\beta$ - individuals ( $p > 0.05$ ). Surface projection of the Pearson correlation available in supplementary Figure 6.

### **4. Performance of plasma p217+tau vs A $\beta$ PET.**

For discriminating the diagnostic groups, the AUC of p217+tau for A $\beta$ + AD vs A $\beta$ - CU was 0.94 [0.90-0.97] and for A $\beta$ + AD vs all CU was 0.88 [0.84-0.92]. For A $\beta$ + AD vs A $\beta$ - dementia the AUC was 0.93 [0.90-0.96].

For discriminating A $\beta$ + PET from A $\beta$ - PET in the entire cohort, the AUC was 0.89 [0.86 - 0.93] or 0.90 if the three A $\beta$ -/tau+ PET individuals (all three had elevated plasma p217+tau) were removed. Youden Index provided a threshold concentration of 126.7fg/ml [100.0–134.4fg/ml] that yielded an accuracy of 0.85, sensitivity 0.79 [0.75-0.89], specificity 0.89 [0.80-0.92], PPV 0.84, and NPV 0.85. P217+tau accurately identified A $\beta$ + individuals among CI participants (AUC=0.89, [0.84-0.93]) and among CU (AUC=0.84 [0.76-0.91]). The Youden threshold was also 126.7fg/ml [97.4-177.2] for the CI giving sensitivity 0.82 [0.66-0.92] and specificity 0.82 [0.75-0.98], PPV 0.92 and NPV 0.65. Youden's index provided an optimal threshold of 100.3fg/ml [100.3-137.3] for CU participants, with a sensitivity of 0.80 [0.78-0.95], specificity of 0.83 [0.67-0.90], PPV 0.49 and NPV 0.95. Applying the whole cohort p217+tau threshold of 126.7fg/ml to CU gave sensitivity of 0.72 [0.6 –0.83], specificity of 0.90 [0.86–0.94], PPV 0.66 and NPV 0.92.

The Z-score threshold for p217+tau derived from CU with <15 CL on A $\beta$  PET was higher at 164 fg/ml resulting in lower sensitivity of 0.60, higher specificity of 0.96, and PPV 0.91 and NPV 0.76 for all subjects to detect positive A $\beta$  PET. In CI, the higher threshold gave sensitivity 0.66, specificity 0.94, PPV 0.96 and NPV 0.53. In CU it gave sensitivity 0.43, specificity 0.96, PPV 0.74 and NPV 0.87.

Altering the CL threshold for A $\beta$ + PET between 25 to 50 CL had no effect on AUC. However, at 15 and 20 CL thresholds the AUC decreased due to more p217+tau false negative results (Figure 3).

## **5. Performance of plasma p217+tau vs <sup>18</sup>F-MK6240 tau PET.**

Plasma p217+tau accurately discriminated individuals with elevated tau in all examined ROIs with the highest AUC in the meta temporal ROI (AUC=0.89 [0.86-0.92]). The scatter plots and ROC curves are shown in Figure 4 for the mesial temporal and meta temporal ROI. The Youden Index threshold was 126.7fg/ml, exactly the same as for A $\beta$  PET and provided an accuracy of 0.81, sensitivity of 0.80 [0.71-0.93] and specificity of 0.82 [0.71-0.93]).

In CU, the highest AUC (0.88 [0.83-0.92]) was observed in the meta temporal region with a Youden's index threshold of 97.8fm/mL giving sensitivity of 0.9 [0.77-0.97] and specificity of 0.75 [0.71-0.88]. Applying the 126.7fg/ml threshold to the CU group gave sensitivity of 0.71 [0.51-0.84] and specificity of 0.85 [0.80-0.89]. Correlation between p217+tau and tau PET was only significant ( $p < 0.05$ ) in the CU subjects who were A $\beta$ + and was strongest in the meta temporal ROI ( $\rho = 0.37$ ) (Figure 4 blue lines).

In CI the highest AUC were observed in the meta temporal and the inferior temporal regions, with AUC's of 0.86 [0.81-0.91]. The mesial temporal AUC was lower at 0.81 [0.75-0.86]. The Youden's index threshold for the meta temporal ROI provided a threshold of 148.4fg/mL with a sensitivity of 0.73 [0.65-0.83] and specificity of 0.90 [0.82-0.97]. Applying the threshold of 126.7fg/ml to the CI cohort gave sensitivity of 0.82 [0.77-0.89] and specificity of 0.71 [0.6-0.79].

### **5.1 p217+tau vs tau PET in subregions of the mesial temporal in cognitively unimpaired.**

The AUC for p217+tau to detect regional tau+ PET in the CU was higher for the meta temporal region at 0.88 [0.83-0.92] than the amygdala 0.81 [0.74 - 0.88], hippocampus 0.81 [0.74-0.88], parahippocampus 0.86 [0.77-0.93] or entorhinal cortex 0.78 [0.71-0.83]. Plotting of the data points (see supplementary Figure 12) shows that discordance was predominantly due to p217+tau+ve/tau PET- subjects with a disproportionately high prevalence of A $\beta$ + in this category compared to p217+tau-ve/tau PET-.

## 6. p217+tau predicts A $\beta$ and Tau pathology independent of confounders

AUC values from base models incorporating age, sex and *APOE*  $\epsilon$ 4 allele status to predict A $\beta$ + and meta temporal tau+ PET were 0.72 [0.67-0.77] and 0.69 [0.64-0.76] respectively. Adding p217+tau to the base model significantly improved prediction (p-value via DeLong's ROC test <0.0001) compared to the base model alone and produced AUC of 0.91 [0.86-0.93] for A $\beta$ + PET and AUC of 0.89 [0.86-0.92] for meta temporal tau+ PET. The combined model, however, did not perform significantly better than p217+tau alone that gave AUC of 0.89 [0.86-0.93] to predict A $\beta$ + and 0.89 [0.86-0.92] for meta temporal tau+ PET.

## 7. Modelling of <sup>18</sup>F-MK6240 SUVR and p217+tau as a function of Centiloid

To unveil the chronological order of the emergence of elevation in biomarkers, we normalised <sup>18</sup>F-MK6240 SUVR and p217+tau concentration to the same scale, using two different methods and mapped against A $\beta$  CL level used as a surrogate for duration of disease development. Firstly, we linearly normalised the range of biomarkers from zero to 100, with zero being the mean of the CU A $\beta$ - individuals with less than 15 CL to increase certainty of A $\beta$  negativity [24], and 100 being the mean of the 30 individuals with the highest biomarker value. When normalising the range of values between 0 and 100, (Figure 5A), the fitted polynomial curves of <sup>18</sup>F-MK6240 in amygdala and plasma p217+tau were nearly identical and rose very early when A $\beta$  level was quite low and before other tau PET regions began to rise. Secondly, we normalized each biomarker to a Z-score, again using those CU with <15 CL to provide the normal range. Figure 5B shows that although we have demonstrated in Figure 5A that plasma p217+tau rises early, it takes until 70CL to exceed two standard deviations of the normal range.

## Discussion

We determined the performance of a plasma ptau assay that utilizes an antibody raised against the paired helical filaments (PHF) of Alzheimer's disease. Human PHF tau is phosphorylated at multiple sites [27]. Plasma p217+tau measures tau phosphorylated at aa217 with binding enhanced by the simultaneous presence of phosphorylation at aa212 so may better reflect the tau of AD than single binding site ptau assays [12]. To evaluate this assay, we employed PET tracers that may give greater sensitivity than those previously used for comparison to plasma ptau [7, 21, 28].

Plasma p217+tau displayed high accuracy in detecting A $\beta$ + individuals across the clinical spectrum. Compared to A $\beta$ - PET CU, plasma p217+tau concentration was two-fold higher in A $\beta$ + CU and 3.5-fold higher in A $\beta$ + MCI and dementia. The AUC for A $\beta$ +AD vs A $\beta$ - CU was 0.94, for A $\beta$ +AD vs A $\beta$ - dementia was 0.93 while the AUC for all A $\beta$ + vs A $\beta$ - and for tau+ vs tau- were equal at 0.89. In the CU group, the AUC was 0.84 for A $\beta$ + vs A $\beta$ - PET and 0.88 for tau+ vs tau- PET. Adding age, sex and *APOE* $\epsilon$ 4 did not significantly improve the prediction of A $\beta$ + PET (AUC 0.89 vs 0.91) or tau+ PET (unchanged at 0.89). These results compare favourably to other plasma ptau assays though the use of different assay methods, different anti-tau antibodies, and cohort variation prevent direct comparison. The range of AUC reported for A $\beta$ + vs A $\beta$ - PET for plasma ptau measures has ranged from 0.76 to 0.92 [2, 3, 7-9, 29, 30]. Our results with p217+tau are at the high end of reported plasma ptau assay performance and used Simoa technology that is kit based, fully automated and has a large installation base and therefore is well suited to widespread deployment.

Our data (Figures 3-5) suggest plasma p217+tau begins to rise i) soon after brain A $\beta$  levels begin to trend up as assessed by PET, ii) concordant with the rise in tau in the amygdala region on PET, and iii) prior to the rise in meta temporal tau PET. This is supported by the vertex-based analysis of correlation in CU, that p217+tau and A $\beta$  are highly associated in

brain regions where early A $\beta$  deposition occurs and with tau in the anteromedial temporal lobe (Figure 2). However, the plasma p217+tau level does not reach +2 standard deviations above the mean found in A $\beta$ - CU until A $\beta$  reaches moderate levels (approximately 70 CL). These findings are consistent with reports for plasma ptau measures of p181, p217 and p231 [6, 9, 31, 32]. Wide intersubject variability in plasma ptau in persons without evidence of AD pathology on amyloid PET remains a challenge and warrants further investigation.

Our data also suggests that tau aggregates in the amygdala region and plasma p217+tau increase together and very early in the development of AD when A $\beta$  load is beginning to rise (Figure 5A). The amygdala performed best in our comparison of p217+tau with tau PET mesial temporal subregions. This may be due to reduced partial volume effect compared to smaller structures and less spill-over effect from off-target binding in the meninges giving a more robust signal. The effect of primary age related tauopathy (PART) where tau aggregates are present without A $\beta$  plaques [33] and predominantly in the entorhinal cortex, on plasma p217+tau levels is unclear and requires further investigation.

Only longitudinal studies will determine if what appear to be false positive p217+tau results in CU compared to PET are due to variation and limitation in the plasma measure or are detecting very early stage AD prior to PET becoming significantly positive and this work is in progress under the AIBL study umbrella. In contrast to the CU, the CI show more “false negative” p217+tau results when compared to both A $\beta$  and tau PET. Many of these false negative p217+tau results were A+/T+ on PET so this requires further investigation.

Reflecting the impact of disease prevalence in our study population (71% A $\beta$ + in CI vs 21% in CU), the p217+tau positive and negative predictive values of p217+tau for A $\beta$ + PET were very different for the CI and CU cohorts. The PPV were high in the CI group for predicting A $\beta$  (0.94) and tau (0.93), suggesting that p217+tau may be a good test to confirm AD in

patients with objective cognitive impairment. However, the NPV were relatively low at 0.66 and 0.65 suggesting that a negative p217+tau does not exclude AD. Lowering the p217+tau threshold to 100 fg/ml in the CI group only improved the NPV to 0.73 while lowering PPV to 0.88 and at 75 fg/ml the NPV was 0.78, PPV 0.84. P217+tau has potential as a diagnostic tool and combination with other biomarkers, as discussed by Schindler et al [34], may increase accuracy and provide an affordable and accurate diagnostic tool. Conversely, in the CU group the NPV were high (>0.94), suggesting that a negative p217+tau test can largely exclude AD pathology in persons with normal cognitive test results. However, the associated lower PPV (0.50) in CU indicates that positive results would need confirmation by PET or CSF analysis. These findings show that triage with p217+tau may substantially reduce the number of screening A $\beta$  PET needed to identify A $\beta$ + PET participants for preclinical AD trials resulting in considerable recruitment cost savings.

This study has several limitations. First the results need validation in other cohorts and more diverse populations. Secondly this is a cross-sectional study and longitudinal analysis is required to confirm the predictive value of p217+tau as a marker of disease progression and to clearly determine when abnormal levels of p217+tau can be measured along the disease trajectory.

In conclusion, an elevated level of plasma p217+tau is associated with both elevated A $\beta$  and tau across the clinical spectrum of Alzheimer's disease. Elevated p217+tau strongly supports a diagnosis of AD in persons with MCI or dementia while a low level in cognitively unimpaired persons is strong evidence against preclinical AD.

## REFERENCES

1. Sevigny J, Suhy J, Chiao P, Chen T, Klein G, Purcell D, et al. Amyloid PET Screening for Enrichment of Early-Stage Alzheimer Disease Clinical Trials: Experience in a Phase 1b Clinical Trial. *Alzheimer Dis Assoc Disord*. 2016;30:1-7. doi:10.1097/wad.000000000000144.
2. Karikari TK, Pascoal TA, Ashton NJ, Janelidze S, Benedet AL, Rodriguez JL, et al. Blood phosphorylated tau 181 as a biomarker for Alzheimer's disease: a diagnostic performance and prediction modelling study using data from four prospective cohorts. *The Lancet Neurology*. 2020;19:422-33. doi:[https://doi.org/10.1016/S1474-4422\(20\)30071-5](https://doi.org/10.1016/S1474-4422(20)30071-5).
3. Thijssen EH, La Joie R, Wolf A, Strom A, Wang P, Iaccarino L, et al. Diagnostic value of plasma phosphorylated tau181 in Alzheimer's disease and frontotemporal lobar degeneration. *Nature Medicine*. 2020;26:387-97. doi:10.1038/s41591-020-0762-2.
4. Lantero Rodriguez J, Karikari TK, Suárez-Calvet M, Troakes C, King A, Emersic A, et al. Plasma p-tau181 accurately predicts Alzheimer's disease pathology at least 8 years prior to post-mortem and improves the clinical characterisation of cognitive decline. *Acta Neuropathologica*. 2020;140:267-78. doi:10.1007/s00401-020-02195-x.
5. Yang C-C, Chiu M-J, Chen T-F, Chang H-L, Liu B-H, Yang S-Y. Assay of Plasma Phosphorylated Tau Protein (Threonine 181) and Total Tau Protein in Early-Stage Alzheimer's Disease. *Journal of Alzheimer's Disease*. 2018;61:1323-32. doi:10.3233/JAD-170810.
6. Janelidze S, Mattsson N, Palmqvist S, Smith R, Beach TG, Serrano GE, et al. Plasma P-tau181 in Alzheimer's disease: relationship to other biomarkers, differential diagnosis, neuropathology and longitudinal progression to Alzheimer's dementia. *Nature Medicine*. 2020;26:379-86. doi:10.1038/s41591-020-0755-1.
7. Mielke MM, Hagen CE, Xu J, Chai X, Vemuri P, Lowe VJ, et al. Plasma phospho-tau181 increases with Alzheimer's disease clinical severity and is associated with tau- and amyloid-positron emission tomography. *Alzheimer's & Dementia*. 2018;14:989-97. doi:<https://doi.org/10.1016/j.jalz.2018.02.013>.
8. Palmqvist S, Janelidze S, Quiroz YT, Zetterberg H, Lopera F, Stomrud E, et al. Discriminative Accuracy of Plasma Phospho-tau217 for Alzheimer Disease vs Other Neurodegenerative Disorders. *JAMA*. 2020;324:772-81. doi:10.1001/jama.2020.12134.
9. Ashton NJ, Pascoal TA, Karikari TK, Benedet AL, Lantero-Rodriguez J, Brinkmalm G, et al. Plasma p-tau231: a new biomarker for incipient Alzheimer's disease pathology. *Acta Neuropathologica*. 2021. doi:10.1007/s00401-021-02275-6.
10. Thijssen EH, La Joie R, Strom A, Fonseca C, Iaccarino L, Wolf A, et al. Plasma phosphorylated tau 217 and phosphorylated tau 181 as biomarkers in Alzheimer's disease and frontotemporal lobar degeneration: a retrospective diagnostic performance study. *Lancet Neurol*. 2021;20:739-52. doi:10.1016/s1474-4422(21)00214-3.
11. Mielke MM, Frank RD, Dage JL, Jeromin A, Ashton NJ, Blennow K, et al. Comparison of Plasma Phosphorylated Tau Species With Amyloid and Tau Positron Emission Tomography, Neurodegeneration, Vascular Pathology, and Cognitive Outcomes. *JAMA Neurology*. 2021;78:1108-17. doi:10.1001/jamaneurol.2021.2293.
12. Van Kolen K, Malia TJ, Theunis C, Nanjunda R, Teplyakov A, Ernst R, et al. Discovery and Functional Characterization of hPT3, a Humanized Anti-Phospho Tau Selective Monoclonal Antibody. *J Alzheimers Dis*. 2020;77:1397-416. doi:10.3233/jad-200544.
13. Triana-Baltzer G, Moughadam S, Slemmon R, Van Kolen K, Theunis C, Mercken M, et al. Development and validation of a high-sensitivity assay for measuring p217+tau in plasma. *Alzheimers Dement (Amst)*. 2021;13:e12204. doi:10.1002/dad2.12204.
14. Rowe CC, Pejoska S, Mulligan RS, Jones G, Chan JG, Svensson S, et al. Head-to-Head Comparison of <sup>11</sup>C-PiB and <sup>18</sup>F-AZD4694 (NAV4694) for  $\beta$ -Amyloid Imaging in Aging and Dementia. *Journal of Nuclear Medicine*. 2013;54:880-6.



15. Pascoal TA, Shin M, Kang MS, Chamoun M, Chartrand D, Mathotaarachchi S, et al. In vivo quantification of neurofibrillary tangles with [(18)F]MK-6240. *Alzheimers Res Ther*. 2018;10:74. doi:10.1186/s13195-018-0402-y.
16. Ellis KA, Bush AI, Darby D, De Fazio D, Foster J, Hudson P, et al. The Australian Imaging, Biomarkers and Lifestyle (AIBL) study of aging: methodology and baseline characteristics of 1112 individuals recruited for a longitudinal study of Alzheimer's disease. *Int Psychogeriatr*. 2009;21:672-87. doi:10.1017/s1041610209009405.
17. Petersen RC, Caracciolo B, Brayne C, Gauthier S, Jelic V, Fratiglioni L. Mild cognitive impairment: a concept in evolution. *J Intern Med*. 2014;275:214-28. doi:10.1111/joim.12190.
18. McKhann GM, Knopman DS, Chertkow H, Hyman BT, Jack CR, Jr., Kawas CH, et al. The diagnosis of dementia due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement*. 2011;7:263-9. doi:10.1016/j.jalz.2011.03.005.
19. Bourgeat P, Doré V, Fripp J, Ames D, Masters CL, Salvado O, et al. Implementing the centiloid transformation for 11c-PiB and  $\beta$ -amyloid 18f-PET tracers using CapAIBL. *NeuroImage*. 2018;183:387-93.
20. Klunk WE, Koeppe RA, Price JC, Benzinger TL, Devous MD, Sr., Jagust WJ, et al. The Centiloid Project: standardizing quantitative amyloid plaque estimation by PET. *Alzheimers Dement*. 2015;11:1-15.e1-4. doi:10.1016/j.jalz.2014.07.003.
21. Rowe CC, Jones G, Dore V, Pejoska S, Margison L, Mulligan RS, et al. Standardized expression of 18F-NAV4694 and 11C-PiB  $\beta$ -amyloid PET results with the Centiloid Scale. *Journal of Nuclear Medicine*. 2016;57:1233-7.
22. Amadoru S, Doré V, McLean CA, Hinton F, Shepherd CE, Halliday GM, et al. Comparison of amyloid PET measured in Centiloid units with neuropathological findings in Alzheimer's disease. *Alzheimer's Research & Therapy*. 2020;12:22. doi:10.1186/s13195-020-00587-5.
23. Doré V, Bullich S, Rowe CC, Bourgeat P, Konate S, Sabri O, et al. Comparison of (18)F-florbetaben quantification results using the standard Centiloid, MR-based, and MR-less CapAIBL(®) approaches: Validation against histopathology. *Alzheimers Dement*. 2019;15:807-16. doi:10.1016/j.jalz.2019.02.005.
24. La Joie R, Ayakta N, Seeley WW, Borys E, Boxer AL, DeCarli C, et al. Multisite study of the relationships between antemortem [(11)C]PIB-PET Centiloid values and postmortem measures of Alzheimer's disease neuropathology. *Alzheimers Dement*. 2019;15:205-16. doi:10.1016/j.jalz.2018.09.001.
25. Dore V, Bourgeat P, Burnham SC, Martins RN, Masters CL, Ames D, et al. AUTOMATED REPORTING OF TAU PET QUANTIFICATION ON THE BRAIN SURFACE. *Alzheimer's & Dementia: The Journal of the Alzheimer's Association*. 2019;15:P1269.
26. Cairns NJ, Ikonomic MD, Benzinger T, Storandt M, Fagan AM, Shah AR, et al. Absence of Pittsburgh Compound B Detection of Cerebral Amyloid  $\beta$  in a Patient With Clinical, Cognitive, and Cerebrospinal Fluid Markers of Alzheimer Disease: A Case Report. *Archives of Neurology*. 2009;66:1557-62. doi:10.1001/archneurol.2009.279.
27. Hanger DP, Betts JC, Loviny TL, Blackstock WP, Anderton BH. New phosphorylation sites identified in hyperphosphorylated tau (paired helical filament-tau) from Alzheimer's disease brain using nanoelectrospray mass spectrometry. *J Neurochem*. 1998;71:2465-76. doi:10.1046/j.1471-4159.1998.71062465.x.
28. Navitsky M, Joshi AD, Kennedy I, Klunk WE, Rowe CC, Wong DF, et al. Standardization of amyloid quantitation with florbetapir standardized uptake value ratios to the Centiloid scale. *Alzheimer's & Dementia*. 2018;14:1565-71. doi:<https://doi.org/10.1016/j.jalz.2018.06.1353>.
29. Brickman AM, Manly JJ, Honig LS, Sanchez D, Reyes-Dumeyer D, Lantigua RA, et al. Plasma p-tau181, p-tau217, and other blood-based Alzheimer's disease biomarkers in a multi-ethnic, community study. *Alzheimer's & Dementia*. 2021;17:1353-64. doi:<https://doi.org/10.1002/alz.12301>.

30. Barthélemy NR, Horie K, Sato C, Bateman RJ. Blood plasma phosphorylated-tau isoforms track CNS change in Alzheimer's disease. *Journal of Experimental Medicine*. 2020;217. doi:10.1084/jem.20200861.
31. Moscoso A, Grothe MJ, Ashton NJ, Karikari TK, Rodriguez JL, Snellman A, et al. Time course of phosphorylated-tau181 in blood across the Alzheimer's disease spectrum. *Brain*. 2021;144:325-39. doi:10.1093/brain/awaa399.
32. Palmqvist S, Insel PS, Stomrud E, Janelidze S, Zetterberg H, Brix B, et al. Cerebrospinal fluid and plasma biomarker trajectories with increasing amyloid deposition in Alzheimer's disease. *EMBO Molecular Medicine*. 2019;11:e11170. doi:<https://doi.org/10.15252/emmm.201911170>.
33. Crary JF, Trojanowski JQ, Schneider JA, Abisambra JF, Abner EL, Alafuzoff I, et al. Primary age-related tauopathy (PART): a common pathology associated with human aging. *Acta Neuropathologica*. 2014;128:755-66. doi:10.1007/s00401-014-1349-0.
34. Schindler SE, Bateman RJ. Combining blood-based biomarkers to predict risk for Alzheimer's disease dementia. *Nature Aging*. 2021;1:26-8. doi:10.1038/s43587-020-00008-0.

*Table 1: Demographic and clinical characteristics*

	<b>CU</b> (n=223)	<b>MCI</b> (n=91)	<b>Dementia</b> (n=83)
Age	75.2 (5.7)	73.6 (8.0)	70.7 (7.9) <sup>*+</sup>
Gender N male (%)	101 (45.3%)	51 (56.0%)	49 (59.0%)
Education (years)	14 (3.0)	12.5 (3.1) <sup>*</sup>	12 (3.0) <sup>*</sup>
APOE ε4+ (%)	72 (32.3%)	46 (50.5%) <sup>*</sup>	46 (55.4%) <sup>*</sup>
MMSE Median (IQR)	29 (2)	27 (4) <sup>*</sup>	23 (4) <sup>*+</sup>
CDR SoB Median (IQR)	0.0 (0.0)	1.0 (1.5) <sup>*</sup>	4.0 (1) <sup>*+</sup>
AIBL PACC Mean (SD)	-0.4 (0.8)	-2.4 (1.1) <sup>*</sup>	-4.3 (2.4) <sup>*+</sup>
Centiloid Mean (SD)	19.5 (38.1)	72.8 (66.2) <sup>*</sup>	93.9 (55.6) <sup>*+</sup>
Aβ+ N (%)	46 (20.6%)	56 (61.4%) <sup>*</sup>	68 (81.9%) <sup>*+</sup>
MK6240 SUVR <sub>MT</sub> Mean (SD)	1.02 (0.23)	1.48 (0.61) <sup>*</sup>	1.93 (0.77) <sup>*+</sup>
Tau <sub>MT</sub> + N(%)	31 (13.9%)	48 (52.7%) <sup>*</sup>	64 (77.1%) <sup>*+</sup>
Adjusted HV (SD)	2.98 (0.26)	2.78 (0.40) <sup>*</sup>	2.57 (0.42) <sup>*+</sup>
Plasma p217+tau Mean	87.8 (67.8)	183.7 (141.0) <sup>*</sup>	229.9 (150.7) <sup>*+</sup>

(SD)			
------	--	--	--

\**p*-value<0.05 compared to CU, +*p*-value<0.05 compared to MCI. AIBL PACC is the AIBL preclinical Alzheimer's cognitive composite;  $T_{MT+}$  means positive for tau in the tau PET meta temporal ROI; HV is hippocampal volume

## Figure Legends

Figure 1: Plasma p217+tau concentrations between A) clinical classification and A $\beta$  PET status and B) Centiloid levels of A $\beta$ . The dashed line corresponds to the threshold derived by Youden Index. C) Centiloid results vs p217+tau level in 25 fg/ml intervals, D) probability of being A $\beta$  PET positive vs p217+tau level. Vertical lines in D) are Youden index derived from cognitively unimpaired (CU) (100.3 fg/ml), Youden index derived from total cohort (126.7 fg/ml) and +2.0 SD of the A $\beta$ -ve CU (164 fg/ml).





Figure 2: Vertex-based analysis of regional Spearman correlation between plasma p217+tau and Centiloid (left column) and  $^{18}\text{F}$ -MK6240 SUVR (right column). CU is cognitively unimpaired.

Figure 3: Plasma p217+tau vs Centiloid measure of A $\beta$ . Scatter plot and ROC curve (A&B) with ROC curves using different Centiloid thresholds to define A $\beta$ + PET; full cohort (A&B), cognitively unimpaired sub-cohort (C&D) and cognitively impaired sub-cohort (E&F). Clinical groups are colour-coded with red for dementia, green for MCI and blue for CU. Solid circles are tau PET positive ( $T_{MT+}$ ). The black dashed horizontal line corresponds to the p217+tau threshold derived from Youden's index. In C) and D) the CU specific threshold is shown. The CI group threshold was the same as the whole cohort threshold. The diamond shapes are the three A $\beta$ - /  $T_{MT+}$  subjects.


Figure 4: Plasma p217+tau vs PET SUVR measures of tau. Scatter plot and ROC curve for mesial temporal ROI (A&B) and meta temporal ROI (C&D) and in CU alone (E&F). Clinical

groups are colour-coded with red for dementia, green for MCI and blue for CU. Solid circles are A $\beta$  PET positive. The black dashed horizontal line corresponds to the p217+tau threshold derived from Youden's index. Linear correlation and Spearman co-efficient are shown for the A $\beta$ + (dark blue) and A $\beta$ - CU (light blue) in part E.


Figure 5: Modelling of <sup>18</sup>F-MK6240 quantification and p217+tau as a function of CL using polynomial curves; A) normalised by linear transform of p217+tau levels and each tau PET ROI SUVR to a scale of zero to 100 where zero is the mean of the results for each marker in A $\beta$ - CU (<15CL) and 100 is the mean of the results for each marker from the 30 individuals with the highest values for each marker. B) normalised by Z-score using the results from A $\beta$ - CU to define the normal range for p217+tau level and each tau PET ROI SUVR. The horizontal line is +2 S.D. A) suggests that plasma p217+tau rises early, similar to amygdala tau at low levels of A $\beta$  while B) shows that the wide normal range for p217+tau delays reaching a 2 S.D. threshold for significance.





P217+tau vs Centiloid




P217+tau vs  $^{18}\text{F}$ -MK6240




Full cohort




Spearman  $\rho$




P217+tau vs Centiloid









P217+tau vs  $^{18}\text{F}$ -MK6240





CU





## Mesial Temporal



## Meta Temporal

