

1 **Supplementary Table 1.** Complete cases population and infection determinants: description and results of the univariable and multivariable
 2 conditional logistic regression analyses adjusted to the week of inclusion.

Characteristics	Cases n = 2001	Controls n = 2001	aOR (95% CI)	
			Univariable analysis	Multivariable analysis
Characteristics				
Age categories, years				
[18–28]	269 (13)	269 (13)		
[29–38]	618 (31)	618 (31)		
[39–48]	595 (30)	595 (30)		
[49+]	519 (26)	519 (26)		
Female sex	1708 (85)	1708 (85)		
At least one comorbidity ^a	301 (15)	223 (11)	1.47 (1.16–1.87)	1.26 (0.89–1.78)
Smoker	355 (18)	334 (17)	1.04 (0.85–1.28)	0.68 (0.50–0.91)
COVID-19 immunization				
None	1471 (74)	761 (38)	Reference	Reference
Partial	316 (16)	520 (26)	0.33 (0.26–0.41)	0.32 (0.24–0.44)
Complete	206 (10)	712 (36)	0.21 (0.16–0.26)	0.16 (0.11–0.23)
Healthcare sector				
Hospital	711 (36)	802 (40)	—	—
Long-term care facility	379 (19)	261 (13)	1.78 (1.39–2.29)	1.41 (0.97–2.05)
Primary care	911 (46)	938 (47)	0.99 (0.83–1.18)	1.55 (1.15–2.08)
HCW Professional category				
Medical professions	171 (8)	559 (28)	—	—
Nurses	454 (23)	386 (19)	5.77 (4.25–7.83)	5.16 (3.37–7.89)
Nurse’s aides	364 (18)	115 (6)	15.2 (10.4–22.2)	9.32 (5.38–16.1)
Others	1012 (51)	941 (47)	4.00 (3.08–5.19)	2.43 (1.70–3.48)
Exposures during the 10 days preceding inclusion				
Regular COVID-19 patient-facing activities	407 (20)	288 (14)	1.64 (1.32–2.04)	1.93 (1.34–2.79)
Exposure to an infected colleague	351 (18)	108 (5)	3.59 (2.70–4.79)	2.92 (1.95–4.38)
Exposure to an infected person outside of work	444 (22)	44 (2)	15.5 (10.1–23.8)	26.8 (16.0–44.7)
Professional cluster (patients and/or HCWs)	391 (20)	177 (9)	2.53 (1.98–3.23)	1.94 (1.36–2.77)
For COVID-19 patients^b care, systematic use of				

Mask type				
Surgical facemask	340 (30)	255 (26)	—	—
Cloth mask	8 (1)	2 (<1)	1.89 (0.34–10.7)	1.26 (0.12–12.8)
N95 respirator	770 (69)	742 (74)	0.65 (0.51–0.83)	0.84 (0.54–1.31)
Gloves	905 (81)	717 (72)	1.50 (1.16–1.94)	1.32 (0.79–2.20)
Eye protection (goggles or faceshield)	668 (60)	691 (69)	0.54 (0.43–0.68)	0.47 (0.31–0.73)
Gown	834 (75)	800 (80)	0.67 (0.51–0.87)	0.65 (0.38–1.11)
Apron	642 (57)	523 (52)	1.43 (1.15–1.78)	1.62 (1.09–2.40)
Did not care for COVID-19 patients	883 (44)	1002 (50)	—	—

3 Results are expressed as n (%) and adjusted odds ratios (aOR) (95% confidence intervals (CI)).

4 ^aComorbidity among: diabetes, arterial hypertension, myocardial infarction and/or chronic pulmonary disease.

5 ^bFor personal protective equipment use, percentages were calculated on the number of HCWs who cared to COVID-19 patients in the ten past
6 days (1118 cases and 999 controls). HCWs, healthcare worker ; PPE, personal protective equipment.

7

8 **Supplementary Table 2** Healthcare workers' use of personal protective equipment for COVID-19-patient care across the three healthcare
 9 settings during the 10 days preceding inclusion.

	Hospital		Long-term-care facility		Primary care	
	Cases n = 694	Controls n = 694	Cases n = 279	Controls n = 279	Cases n = 921	Controls n = 921
Personal protective equipment						
Mask						
Surgical facemask	83 (18)	102 (22)	67 (44)	64 (43)	151 (40)	78 (25)
N95 respirator	383 (82)	355 (78)	85 (56)	84 (57)	226 (60)	237 (75)
Gloves	409 (88)	351 (77)	127 (84)	113 (76)	258 (68)	205 (65)
Eye protection (goggles or faceshield)	346 (74)	351 (77)	86 (57)	113 (76)	173 (46)	167 (53)
Gown	408 (88)	388 (85)	102 (67)	118 (80)	228 (60)	214 (68)
Apron	324 (70)	277 (61)	92 (61)	110 (74)	138 (37)	101 (32)
Did not care for COVID-19 patients	228 (33)	238 (34)	127 (45)	131 (47)	544 (59)	606 (66)

10 Results are expressed as number (%).

11

12

13

14

15

16

17

18

19

20

21 **Supplementary Table 3.** Infection determinants of subgroup multivariable conditional logistic regression analysis according to healthcare
 22 setting.

	Hospital n = 694 cases/694 controls	Long-term-care facility n = 279 cases/279 controls	Primary care n = 921 cases/921 controls
Characteristic			
At least one comorbidity ^a	1.76 (0.99–3.12)	1.26 (0.40–3.96)	0.90 (0.50–1.62)
Smoker	0.77 (0.46–1.28)	0.60 (0.24–1.50)	1.04 (0.63–1.71)
COVID-19 immunization			
None	Reference	Reference	Reference
Partial	0.15 (0.08–0.27)	0.09 (0.02–0.37)	0.34 (0.22–0.53)
Complete	0.12 (0.06–0.22)	0.13 (0.04–0.50)	0.16 (0.09–0.30)
Exposures during the 10 days preceding inclusion			
Regular COVID-19-patient-facing activities	1.95 (1.15–3.28)	0.19 (0.04–0.97)	3.64 (1.59–8.34)
Exposure to an infected colleague	1.18 (0.66–2.09)	4.34 (0.97–19.4)	6.94 (2.45–19.6)
Exposure to an infected person outside of work	15.7 (7.38–33.5)	18.4 (4.53–74.3)	19.8 (9.51–41.3)
Professional cluster (patients and/or HCWs)	2.08 (1.19–3.63)	3.85 (0.79–18.7)	1.97 (0.97–3.99)
For COVID-19-patients care, systematic use of			
Mask type			
Surgical facemask	Reference	Reference	Reference
N95 respirator	1.68 (0.79–3.56)	1.23 (0.30–5.01)	0.61 (0.28–1.32)
Gloves	1.29 (0.58–2.87)	0.97 (0.11–8.69)	2.50 (0.97–6.45)
Eye protection (goggles or faceshield)	0.90 (0.46–1.77)	0.30 (0.04–2.12)	1.03 (0.50–2.11)
Gown	0.70 (0.30–1.62)	0.33 (0.03–3.37)	0.61 (0.27–1.41)
Apron	1.18 (0.66–2.11)	1.28 (0.25–6.47)	1.58 (0.83–3.01)

23 Results are presented as n (%), and adjusted Odds Ratios [aOR] (95% Confidence Intervals [CI]).

24 ^a Comorbidity among: diabetes, arterial hypertension, myocardial infarction and/or chronic pulmonary disease. HCWs, healthcare worker ; PPE,
 25 personal protective equipment.