

Title page

Cross-cultural adaptation and validation of the “COVID Stress Scales” in Greek

Petros Galanis, Assistant Professor, Clinical Epidemiology Laboratory, Faculty of Nursing, National and Kapodistrian University of Athens, Athens, Greece

Irene Vraka, MD, PhD, Department of Radiology, P & A Kyriakou Children's Hospital, Athens, Greece

Aglaia Katsiroumpa, RN, MSc, Faculty of Nursing, National and Kapodistrian University of Athens, Athens, Greece

Olympia Konstantakopoulou, Economist, MSc, PhD, Center for Health Services Management and Evaluation, Faculty of Nursing, National and Kapodistrian University of Athens, Athens, Greece

Olga Siskou, RN, MSc, PhD, University of Piraeus, Piraeus, Greece

Eleftheria Zogaki, Midwife, MSc, Faculty of Midwifery, Technological Education Institution of Western Macedonia, Kozani, Greece

Daphne Kaitelidou, Professor, Center for Health Services Management and Evaluation, Faculty of Nursing, National and Kapodistrian University of Athens, Athens, Greece

Corresponding author: Petros Galanis, Assistant Professor, Clinical Epidemiology Laboratory, Faculty of Nursing, National and Kapodistrian University of Athens, 123 Papadiamantopoulou street, GR-11527, Athens, Greece, e-mail: pegalan@nurs.uoa.gr, <https://orcid.org/0000-0002-1308-5782>

Abstract

Background: During the COVID-19 pandemic, several instruments were developed to measure the psychological impact of COVID-19, such as fear, anxiety, post-traumatic stress, phobia, etc.

Objective: To adapt cross-cultural and validate the “COVID Stress Scales” in Greek.

Methods: We conducted a cross-sectional study with 200 participants between November 2021 to February 2022. All participants were adults, and a convenience sample was obtained. We applied the forward-backward translation method to create a Greek version of the “COVID Stress Scales”. We assessed reliability of the questionnaire with test-retest method in a 10-day window, and we assessed validity of the questionnaire with exploratory factor analysis.

Results: Our five-factor model explained 72% of the variance and totally confirmed the factors of the initial “COVID Stress Scales”. In particular, we found the following five factors: (a) COVID-19 danger and contamination (eleven items), (b) COVID-19 socioeconomic consequences (six items), (c) COVID-19 xenophobia (six items), (d) COVID-19 traumatic stress (six items), and (e) COVID-19 compulsive checking (six items). Cronbach’s coefficients alpha for the five factors that emerged from the exploratory factor analysis were greater than 0.89 indicating excellent internal reliability.

Conclusions: We found that the “COVID Stress Scales” is a reliable and valid tool to measure stress due to the COVID-19 in the Greek population.

Key words: COVID-19; COVID Stress Scales; questionnaire; validation; Greece

Introduction

Distress, anxiety, fear, and several other psychological symptoms are common during the COVID-19 pandemic (Bareeqa et al., 2021; Cénat et al., 2021; Galanis, Andreadaki, et al., 2021; Galanis, Petrogianni, et al., 2021; Patelarou et al., 2022; Yuan et al., 2021; Zhang et al., 2021). For instance, according to studies in China, more than 25% of the general population experienced moderate to severe levels of stress and anxiety during the COVID-19 pandemic (Qiu et al., 2020; Wang et al., 2020). The psychological management of the COVID-19 is essential since higher levels of perceived threat from the COVID-19 and fear of COVID-19 are associated with increased willingness to get vaccinated against COVID-19 (Galanis et al., 2022; Galanis, Vraka, et al., 2021; Patelarou et al., 2021). On the other hand, increased anxiety is related with socially disruptive behaviors, such as panic buying and surging unnecessarily into hospitals when people misinterpret their minor symptoms as signs of serious disease (Asmundson & Taylor, 2020a, 2020b).

Anxiety is important in shaping behavioral responses to the COVID-19 pandemic and COVID-19 stressors affect almost everyone in the world to different extent. During the COVID-19 pandemic, several instruments were developed to measure the psychological impact of COVID-19, such as fear, anxiety, post-traumatic stress, phobia, etc. (Ahorsu et al., 2020; Ahuja, 2021; Arpaci et al., 2020; Forte et al., 2020; Geldsetzer, 2020; Kira et al., 2021; Mertens et al., 2020; Pérez-Fuentes et al., 2020; Petzold et al., 2020; Varshney et al., 2020; Zhong et al., 2020). Many scales have a unidimensional structure, while others consist of factors, such as trauma, phobia, post-traumatic stress, anxiety, fear, etc. Moreover, construct and content validity of some scales is not evaluated, thus limiting the validity of these tools.

Among others, the “COVID Stress Scales” are one of the first scales worldwide that are developed to assess COVID-19 related stress (Taylor et al., 2020). The aim of our study was to adapt cross-cultural and validate the “COVID Stress Scales” in Greek.

Methods

Study design

We conducted a cross-sectional study with 200 participants between November 2021 to February 2022. All participants were adults, and a convenience sample was

obtained. We applied the forward-backward translation method to create a Greek version of the “COVID Stress Scales” (Galanis, 2019). First, we conducted a pilot study with 30 participants to assess the face validity of the questionnaire (Galanis, 2013). Minimal corrections were made, resulting in the final version of the “COVID Stress Scales” in Greek. We assessed reliability of the questionnaire with test-retest method in a 10-day window. Also, we assessed validity of the questionnaire with exploratory factor analysis. Data were collected in an anonymous way and study protocol was approved from the Faculty of Nursing, National and Kapodistrian University of Athens (reference number; 370, 02-09-2021).

Questionnaire

The “COVID Stress Scales” comprised 36 items, while responses ranged from 0 (never) to 4 (almost always). Higher values indicate greater stress. According to the creators of the questionnaire, the “COVID Stress Scales” consist of five factors: (a) danger and contamination fears, (b) fears about economic consequences, (c) xenophobia, (d) compulsive checking and reassurance seeking, and (e) traumatic stress symptoms about COVID-19 (Taylor et al., 2020).

Statistical analysis

We used exploratory factor analysis to assess the construct validity of the “COVID Stress Scales”. We used the varimax rotation method with an acceptable level for factor loading of 0.4. We set the minimum acceptable value for the eigenvalues at 1. We calculated the Kaiser-Meyer-Olkin measure to assess the adequacy of the sample size considering values >0.8 as acceptable. Also, we performed Bartlett’s test of sphericity considering a p-value <0.05 as acceptable. We assessed reliability of the “COVID Stress Scales” with test-retest method in a 10-day window. In that case, we calculated, Pearson’s correlation coefficient for the 36 items with values >0.7 considered to be exceptional. Also, we calculated Cronbach’s coefficient alpha for the factors that emerged from the exploratory factor analysis with values greater than 0.7 indicate an acceptable level of internal reliability. We used numbers and percentages to present categorical variables and mean, standard deviation, median, minimum value, and maximum value to present continuous variables. We calculated mean scores for each factor. In that case, we added the answers in the items of each factor and then we divided the sum with the total number of items. Thus, each factor had a

total score from 0 to 4 with higher values indicate a higher level of stress. All tests of statistical significance were two-tailed, and p -values < 0.05 were considered as statistically significant. Statistical analysis was performed with the IBM SPSS 21.0 (IBM Corp. Released 2012. IBM SPSS Statistics for Windows, Version 21.0 Armonk, NY, USA).

Results

Demographic characteristics

Detailed demographic characteristics of the participants is shown in Table 1. Most of the participants were females (65%), were married (89%), had children (36%) and had a university degree (73%). Mean age of the participants was 34.1 years old. Among the participants, 26% were diagnosed with COVID-19 and 79% a family member/friend was diagnosed. Twenty-three-point five percent were vaccinated for seasonal influenza and 59.5% were vaccinated for COVID-19.

Factor analysis

Detailed results of the exploratory factor analysis are presented in the Table 2. The p -value for Bartlett's test of sphericity was < 0.001 and the Kaiser–Meyer–Olkin measure was 0.87, indicating that our sample was adequate to perform the exploratory factor analysis. We found five factors including 35 out of items of the “COVID Stress Scales”. Our five-factor model explained 72% of the variance and totally confirmed the factors of the initial “COVID Stress Scales”. In particular, we found the following five factors: (a) COVID-19 danger and contamination (eleven items), (b) COVID-19 socioeconomic consequences (six items), (c) COVID-19 xenophobia (six items), (d) COVID-19 traumatic stress (six items), and (e) COVID-19 compulsive checking (six items). The only item that was not included in our five-factor model was the following: “I am worried that my mail has been contaminated by mail handlers”.

Reliability analysis

Cronbach's coefficients alpha for the five factors that emerged from the exploratory factor analysis were greater than 0.89 indicating excellent internal reliability. In particular, Cronbach's coefficient alpha for the factor “COVID-19 danger and

contamination” was 0.93, for the factor “COVID-19 socioeconomic consequences” was 0.94, for the factor “COVID-19 xenophobia” was 0.94, for the factor “COVID-19 traumatic stress” was 0.93, and for the factor “COVID-19 compulsive checking” was 0.89. Moreover, Pearson’s correlation coefficients for the 36 items and the five factors were greater than 0.77 (p-value<0.001 in all cases) indicating very good reliability of the “COVID Stress Scales”.

Descriptive statistics

Descriptive statistics for the five factors of the “COVID Stress Scales” is presented in Table 3. Stress due to COVID-19 danger and contamination was the highest (mean=2.2), and then stress due to COVID-19 xenophobia (mean=1.1), stress due to COVID-19 compulsive checking (mean=0.9), stress due to COVID-19 socioeconomic consequences (mean=0.6), and stress due to COVID-19 traumatic stress (mean=0.5). In general, stress due to COVID-19 was low to moderate.

Discussion

We translated and validated the “COVID Stress Scales” in Greek. We found that reliability and validity of the scale was excellent, thus it could be used in Greek samples to measure stress due to the COVID-19. The scale comprised 36 items that are simple and easy to answer them. Our five-factor model totally confirms the five-factor model of the initial “COVID Stress Scales” with the following five factors: (a) danger and contamination, (b) socioeconomic consequences, (c) xenophobia, (d) traumatic stress, and (e) compulsive checking.

One of the factors that we identified through factor analysis was COVID-19 danger and contamination. The COVID-19 pandemic has become a source of fear worldwide and continues to lead people to high levels of fear, anxiety, and insecurity (Heiat et al., 2021; Luo et al., 2021; Muller et al., 2021). Measuring the level of fear, anxiety, and insecurity in different countries and populations could help policy makers to identify high-risk groups in need of public health and education campaigns. For instance, fear is higher in females, in hospital staff, in college students, and in participants in studies in Asia (Luo et al., 2021). Misconceptions and misunderstandings mainly due to media, public awareness level, contradictory data

and non-scientific speculations could be the main reasons for the development of fear and corona-phobia (Heiat et al., 2021).

The second factor that we found in our model was COVID-19 socioeconomic consequences. The COVID-19 pandemic has made many poor countries worldwide face challenging socioeconomic consequences (Buheji et al., 2020; Patterson et al., 2021; Sharma et al., 2021; Singh & Misra, 2020). Lack of vaccines and shortcomings in medical infrastructure are the main reasons that affect negatively the poor societies widening the bridge between the poor and the rich, thus the poor are becoming poorer and the rich are becoming richer. Policymakers should apply immediate control plans and strategies to minimize the impact of the COVID-19 pandemic on the socio-economic activities of the poor. Low-income countries with fewer resources should adopt policies to minimize the impact of the COVID-19 pandemic on the livelihood of the poor.

Also, we found that COVID-19 traumatic stress is another factor in our five-model factor. The prevalence of post-traumatic stress disorder due to the COVID-19 pandemic is high not only to health professionals but also to the general population (Caruso et al., 2021; d’Ettorre et al., 2021; Li et al., 2021; Nagarajan et al., 2022; Sahebi et al., 2021). The prevalence of post-traumatic stress disorder during the COVID-19 pandemic reaches 15% in the general population and 21.5% in healthcare workers. The COVID-19 pandemic has placed people under tremendous psychological stress and prevention of post-traumatic stress disorder is a challenge worldwide. There is a need for urgent appropriate support, interventions to protect people from the psychological impact of the COVID-19 pandemic and healthcare policies to prevent and manage post-traumatic stress disorder.

Finally, we found that COVID-19 compulsive checking is one of the factors that create stress in people. Extreme focus on hygiene measures and contamination during the COVID-19 pandemic could exacerbate obsessive-compulsive symptoms (Cunning & Hodes, 2022). The COVID-19 pandemic appears to be related with a worsening of obsessive-compulsive symptoms especially during the early stages of the pandemic (Guzick et al., 2021; Maye et al., 2022). Moreover, the COVID-19 pandemic has been an enormous stressor for patients with contamination-related obsessive-compulsive

disorder. Treatment personalization and effective evidence-based treatment for obsessive-compulsive disorder should be an urgent public health priority.

Limitations

Our study has several limitations. According to our knowledge, this is the first study that assesses the validity of the “COVID Stress Scales”. We found that reliability and validity of the scale was excellent but further studies in different populations and settings should test the psychometric properties of the scale. Also, our sample comprised 200 participants, thus studies with larger and more representative samples could add more information regarding the “COVID Stress Scales”. More sophisticated analysis such as confirmatory factor analysis could be applied to confirm or not our results.

Conclusions

We found that the “COVID Stress Scales” is a reliable and valid tool to measure stress due to the COVID-19 in Greek populations. Our five-factor model totally confirms the five-factor model of the initial “COVID Stress Scales” with the following five factors: (a) danger and contamination, (b) socioeconomic consequences, (c) xenophobia, (d) traumatic stress, and (e) compulsive checking. Accurate measurement of individuals’ stress is essential to understand feelings and responses during a major public health threaten as the COVID-19 pandemic. Further studies could expand our evidence and add valuable data regarding the “COVID Stress Scales”.

Table 1. Demographic characteristics of the participants.

Characteristics	N	%
Gender		
Females	130	65
Males	70	35
Age (years) ^a	34.1	8.1
Marital status		
Singles	22	11
Married	178	89
Children		
No	128	64
Yes	72	36
Educational level		
High school	54	27
University degree	146	73
Previous COVID-19 diagnosis		
No	148	74
Yes	52	26
Previous COVID-19 diagnosis for family members/friends		
No	42	21
Yes	158	79
Vaccinated for seasonal influenza		
No	153	76.5

Yes	47	23.5
Vaccinated for COVID-19		
No	81	40.5
Yes	119	59.5

^a mean, standard deviation

Table 2. Exploratory factor analysis for the 36 items of the “COVID Stress Scales”.

The following questions ask about various kinds of worries that you might have experienced over the past seven days about the coronavirus and the COVID-19	Factors				
	COVID-19 danger and contamination	COVID-19 socioeconomic consequences	COVID-19 xenophobia	COVID-19 traumatic stress	COVID-19 compulsive checking
I am worried about catching the virus	0.42				
I am worried that basic hygiene (e.g., handwashing) is not enough to keep me safe from the virus	0.67				
I am worried that our healthcare system is unable to keep me safe from the virus	0.77				
I am worried that I can't keep my family safe from the virus	0.83				
I am worried that our healthcare system won't be able to protect my loved ones	0.83				
I am worried that social distancing is not enough to keep me safe from the virus	0.73				
I am worried about grocery stores running out of food		0.84			
I am worried about grocery stores running out of cold or flu remedies		0.81			
I am worried about pharmacies running out of prescription medicines		0.83			
I am worried about grocery stores running out of water		0.91			
I am worried about grocery stores running out of cleaning or disinfectant supplies		0.80			
I am worried that grocery stores will close down		0.83			
I am worried that foreigners are spreading the virus in my country			0.83		
If I met a person from a foreign country, I'd be worried that they might have the virus			0.87		
I am worried about coming into contact with foreigners because they might have the virus			0.88		

I am worried that foreigners are spreading the virus because they're not as clean as we are			0.78		
If I went to a restaurant that specialized in foreign foods, I'd be worried about catching the virus			0.62		
If I was in an elevator with a group of foreigners, I'd be worried that they're infected with the virus			0.68		
I am worried that people around me will infect me with the virus	0.54				
I am worried that if I touched something in a public space (e.g., handrail, door handle), I would catch the virus	0.63				
I am worried that if someone coughed or sneezed near me, I would catch the virus	0.75				
I am worried that I might catch the virus from handling money or using a debit machine	0.49				
I am worried about taking change in cash transactions	0.50				
I am worried that my mail has been contaminated by mail handlers					
I had trouble sleeping because I worried about the virus				0.78	
I had bad dreams about the virus				0.82	
I thought about the virus when I didn't mean to				0.82	
Disturbing mental images about the virus popped into my mind against my will				0.86	
had trouble concentrating because I kept thinking about the virus				0.86	
Reminders of the virus caused me to have physical reactions, such as sweating or a pounding heart				0.83	
Social media posts concerning COVID-19					0.77
Seeing YouTube videos about COVID-19					0.67
Seeking reassurance from friends or family about COVID-19					0.62

Checking your own body for signs of infection (e.g., taking your temperature)					0.57
Asking health professionals (e.g., doctors or pharmacists) for advice about COVID-19					0.80
Searched the Internet for treatments for COVID-19					0.89
Cronbach's alpha coefficient	0.93	0.94	0.94	0.93	0.89

Values express loadings.

Table 3. Descriptive statistics for the five factors of the “COVID Stress Scales”.

Factor	Mean	Standard deviation	Median	Minimum value	Maximum value
COVID-19 danger and contamination	2.2	1.1	2.5	0	4
COVID-19 socioeconomic consequences	0.6	1.0	0	0	4
COVID-19 xenophobia	1.1	1.2	0.5	0	4
COVID-19 traumatic stress	0.5	0.8	0	0	3
COVID-19 compulsive checking	0.9	0.9	0.7	0	3.8

References

- Ahorsu, D. K., Lin, C.-Y., Imani, V., Saffari, M., Griffiths, M. D., & Pakpour, A. H. (2020). The Fear of COVID-19 Scale: Development and Initial Validation. *International Journal of Mental Health and Addiction*. <https://doi.org/10.1007/s11469-020-00270-8>
- Ahuja, K. K. (2021). Scanning the VIRUS: A study of dimensions of stress and coping with COVID-19. *Current Psychology*. <https://doi.org/10.1007/s12144-021-01369-4>
- Arpaci, I., Karataş, K., & Baloğlu, M. (2020). The development and initial tests for the psychometric properties of the COVID-19 Phobia Scale (C19P-S). *Personality and Individual Differences, 164*, 110108. <https://doi.org/10.1016/j.paid.2020.110108>
- Asmundson, G. J. G., & Taylor, S. (2020a). Coronaphobia: Fear and the 2019-nCoV outbreak. *Journal of Anxiety Disorders, 70*, 102196. <https://doi.org/10.1016/j.janxdis.2020.102196>
- Asmundson, G. J. G., & Taylor, S. (2020b). How health anxiety influences responses to viral outbreaks like COVID-19: What all decision-makers, health authorities, and health care professionals need to know. *Journal of Anxiety Disorders, 71*, 102211. <https://doi.org/10.1016/j.janxdis.2020.102211>
- Bareeqa, S. B., Ahmed, S. I., Samar, S. S., Yasin, W., Zehra, S., Monese, G. M., & Gouthro, R. V. (2021). Prevalence of depression, anxiety and stress in china during COVID-19 pandemic: A systematic review with meta-analysis. *The International Journal of Psychiatry in Medicine, 56*(4), 210–227. <https://doi.org/10.1177/0091217420978005>
- Buheji, M., da Costa Cunha, K., Beka, G., Mavrić, B., Leandro do Carmo de Souza, Y., Souza da Costa Silva, S., Hanafi, M., & Chetia Yein, T. (2020). The Extent of COVID-19 Pandemic Socio-Economic Impact on Global Poverty. A Global Integrative Multidisciplinary Review. *American Journal of Economics, 10*(4), 213–224. <https://doi.org/10.5923/j.economics.20201004.02>
- Caruso, R., Annaloro, C., Arrigoni, C., Ghizzardi, G., Dellafiore, F., Magon, A., Maga, G., Nania, T., Pittella, F., & Villa, G. (2021). Burnout and post-traumatic stress disorder in frontline nurses during the COVID-19 pandemic: A systematic literature review and meta-analysis of studies published in 2020: COVID-19, burnout, and

PTSD in nurses. *Acta Biomedica Atenei Parmensis*, 92(S2), e2021428.
<https://doi.org/10.23750/abm.v92iS2.11796>

Cénat, J. M., Blais-Rochette, C., Kokou-Kpolou, C. K., Noorishad, P.-G., Mukunzi, J. N., McIntee, S.-E., Dalexis, R. D., Goulet, M.-A., & Labelle, P. R. (2021). Prevalence of symptoms of depression, anxiety, insomnia, posttraumatic stress disorder, and psychological distress among populations affected by the COVID-19 pandemic: A systematic review and meta-analysis. *Psychiatry Research*, 295, 113599.
<https://doi.org/10.1016/j.psychres.2020.113599>

Cunning, C., & Hodes, M. (2022). The COVID-19 pandemic and obsessive-compulsive disorder in young people: Systematic review. *Clinical Child Psychology and Psychiatry*, 27(1), 18–34. <https://doi.org/10.1177/13591045211028169>

d’Ettorre, G., Ceccarelli, G., Santinelli, L., Vassalini, P., Innocenti, G. P., Alessandri, F., Koukopoulos, A. E., Russo, A., d’Ettorre, G., & Tarsitani, L. (2021). Post-Traumatic Stress Symptoms in Healthcare Workers Dealing with the COVID-19 Pandemic: A Systematic Review. *International Journal of Environmental Research and Public Health*, 18(2), 601. <https://doi.org/10.3390/ijerph18020601>

Forte, G., Favieri, F., Tambelli, R., & Casagrande, M. (2020). COVID-19 Pandemic in the Italian Population: Validation of a Post-Traumatic Stress Disorder Questionnaire and Prevalence of PTSD Symptomatology. *International Journal of Environmental Research and Public Health*, 17(11), 4151. <https://doi.org/10.3390/ijerph17114151>

Galanis, P. (2013). Validity and reliability of questionnaires in epidemiological studies. *Arch Hellen Med*, 30(1), 97–110.

Galanis, P. (2019). Translation and cross-cultural adaptation methodology for questionnaires in languages other than Greek. *Arch Hellen Med*, 36(1), 124–135.

Galanis, P., Andreadaki, E., Kleanthous, E., Georgiadou, A., Evangelou, E., Kallergis, G., & Kaitelidou, D. (2021). Determinants of Psychological Distress during the Covid-19 Pandemic and the Lockdown Measures in Greece and Cyprus. *Int J Caring Sci*, 14(2), 837–853.

Galanis, P., Petrogianni, E., Vraka, I., Konstantakopoulou, O., Siskou, O., Bilali, A., & Kaitelidou, D. (2021). *Fear of COVID-19 among nurses in mobile COVID-19 testing units in Greece* [Preprint]. Nursing. <https://doi.org/10.1101/2021.07.05.21260037>

- Galanis, P., Vraka, I., Fragkou, D., Bilali, A., & Kaitelidou, D. (2021). Intention of healthcare workers to accept COVID-19 vaccination and related factors: A systematic review and meta-analysis. *Asian Pacific Journal of Tropical Medicine*, *14*(12), 543. <https://doi.org/10.4103/1995-7645.332808>
- Galanis, P., Vraka, I., Siskou, O., Konstantakopoulou, O., Katsiroumpa, A., & Kaitelidou, D. (2022). Willingness, refusal and influential factors of parents to vaccinate their children against the COVID-19: A systematic review and meta-analysis. *Preventive Medicine*, *157*, 106994. <https://doi.org/10.1016/j.ypmed.2022.106994>
- Geldsetzer, P. (2020). Knowledge and Perceptions of COVID-19 Among the General Public in the United States and the United Kingdom: A Cross-sectional Online Survey. *Annals of Internal Medicine*, *173*(2), 157–160. <https://doi.org/10.7326/M20-0912>
- Guzick, A. G., Candelari, A., Wiese, A. D., Schneider, S. C., Goodman, W. K., & Storch, E. A. (2021). Obsessive–Compulsive Disorder During the COVID-19 Pandemic: A Systematic Review. *Current Psychiatry Reports*, *23*(11), 71. <https://doi.org/10.1007/s11920-021-01284-2>
- Heiat, M., Heiat, F., & Halaji, M. (2021). Phobia and Fear of COVID-19: Origins, complications and management, a narrative review. *Annali Di Igiene Medicina Preventiva e Di Comunità*, *4*, 360–370. <https://doi.org/10.7416/ai.2021.2446>
- Kira, I. A., Shuwiekh, H. A. M., Rice, K. G., Ashby, J. S., Elwakeel, S. A., Sous, M. S. F., Alhuwailah, A., Baali, S. B. A., Azdaou, C., Oliemat, Enas. M., & Jamil, H. J. (2021). Measuring COVID-19 as Traumatic Stress: Initial Psychometrics and Validation. *Journal of Loss and Trauma*, *26*(3), 220–237. <https://doi.org/10.1080/15325024.2020.1790160>
- Li, Y., Scherer, N., Felix, L., & Kuper, H. (2021). Prevalence of depression, anxiety and post-traumatic stress disorder in health care workers during the COVID-19 pandemic: A systematic review and meta-analysis. *PLOS ONE*, *16*(3), e0246454. <https://doi.org/10.1371/journal.pone.0246454>
- Luo, F., Ghanei Gheshlagh, R., Dalvand, S., Saedmoucheshi, S., & Li, Q. (2021). Systematic Review and Meta-Analysis of Fear of COVID-19. *Frontiers in Psychology*, *12*, 661078. <https://doi.org/10.3389/fpsyg.2021.661078>
- Maye, C. E., Wojcik, K. D., Candelari, A. E., Goodman, W. K., & Storch, E. A. (2022). Obsessive compulsive disorder during the COVID-19 pandemic: A brief

- review of course, psychological assessment and treatment considerations. *Journal of Obsessive-Compulsive and Related Disorders*, 100722. <https://doi.org/10.1016/j.jocrd.2022.100722>
- Mertens, G., Gerritsen, L., Duijndam, S., Salemink, E., & Engelhard, I. M. (2020). Fear of the coronavirus (COVID-19): Predictors in an online study conducted in March 2020. *Journal of Anxiety Disorders*, 74, 102258. <https://doi.org/10.1016/j.janxdis.2020.102258>
- Muller, A. E., Himmels, J. P. W., & Van de Velde, S. (2021). Instruments to measure fear of COVID-19: A diagnostic systematic review. *BMC Medical Research Methodology*, 21(1), 82. <https://doi.org/10.1186/s12874-021-01262-5>
- Nagarajan, R., Krishnamoorthy, Y., Basavarachar, V., & Dakshinamoorthy, R. (2022). Prevalence of post-traumatic stress disorder among survivors of severe COVID-19 infections: A systematic review and meta-analysis. *Journal of Affective Disorders*, 299, 52–59. <https://doi.org/10.1016/j.jad.2021.11.040>
- Patelarou, E., Galanis, P., Mechili, E. A., Argyriadi, A., Argyriadis, A., Asimakopoulou, E., Brokaj, S., Bucaj, J., Carmona-Torres, J. M., Cobo-Cuenca, A. I., Doležel, J., Finotto, S., Jarošová, D., Kalokairinou, A., Mecugni, D., Pulomenaj, V., Saliáj, A., Sopjani, I., Zahaj, M., & Patelarou, A. (2021). Factors influencing nursing students' intention to accept COVID-19 vaccination: A pooled analysis of seven European countries. *Nurse Education Today*, 104, 105010. <https://doi.org/10.1016/j.nedt.2021.105010>
- Patelarou, E., Galanis, P., Mechili, E. A., Argyriadi, A., Argyriadis, A., Asimakopoulou, E., Kicaj, E., Bucaj, J., Carmona-Torres, J. M., Cobo-Cuenca, A. I., Doležel, J., Finotto, S., Jarošová, D., Kalokairinou, A., Mecugni, D., Pulomenaj, V., Malaj, K., Sopjani, I., Zahaj, M., & Patelarou, A. (2022). Assessment of COVID-19 Fear in Five European Countries before Mass Vaccination and Key Predictors among Nurses and Nursing Students. *Vaccines*, 10(1), 98. <https://doi.org/10.3390/vaccines10010098>
- Patterson, G. E., McIntyre, K. M., Clough, H. E., & Rushton, J. (2021). Societal Impacts of Pandemics: Comparing COVID-19 With History to Focus Our Response. *Frontiers in Public Health*, 9, 630449. <https://doi.org/10.3389/fpubh.2021.630449>
- Pérez-Fuentes, M. del C., Molero Jurado, M. del M., Oropesa Ruiz, N. F., Martos Martínez, Á., Simón Márquez, M. del M., Herrera-Peco, I., & Gázquez Linares, J. J.

- (2020). Questionnaire on Perception of Threat from COVID-19. *Journal of Clinical Medicine*, 9(4), 1196. <https://doi.org/10.3390/jcm9041196>
- Petzold, M. B., Bendau, A., Plag, J., Pyrkosch, L., Maricic, L. M., Rogoll, J., Betzler, F., Große, J., & Ströhle, A. (2020). Development of the COVID-19-Anxiety Questionnaire and first psychometric testing. *BJPsych Open*, 6(5), e91. <https://doi.org/10.1192/bjo.2020.82>
- Qiu, J., Shen, B., Zhao, M., Wang, Z., Xie, B., & Xu, Y. (2020). A nationwide survey of psychological distress among Chinese people in the COVID-19 epidemic: Implications and policy recommendations. *General Psychiatry*, 33(2), e100213. <https://doi.org/10.1136/gpsych-2020-100213>
- Sahebi, A., Yousefi, A., Abdi, K., Jamshidbeigi, Y., Moayedi, S., Torres, M., Wesemann, U., Sheikhbardsiri, H., & Golitaleb, M. (2021). The Prevalence of Post-traumatic Stress Disorder Among Health Care Workers During the COVID-19 Pandemic: An Umbrella Review and Meta-Analysis. *Frontiers in Psychiatry*, 12, 764738. <https://doi.org/10.3389/fpsy.2021.764738>
- Sharma, A., Ghosh, D., Divekar, N., Gore, M., Gochhait, S., & Shireshi, S. S. (2021). Comparing the socio-economic implications of the 1918 Spanish flu and the COVID-19 pandemic in India: A systematic review of literature. *International Social Science Journal*, 71(S1), 23–36. <https://doi.org/10.1111/issj.12266>
- Singh, A. K., & Misra, A. (2020). Impact of COVID-19 and comorbidities on health and economics: Focus on developing countries and India. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*, 14(6), 1625–1630. <https://doi.org/10.1016/j.dsx.2020.08.032>
- Taylor, S., Landry, C. A., Paluszek, M. M., Fergus, T. A., McKay, D., & Asmundson, G. J. G. (2020). Development and initial validation of the COVID Stress Scales. *Journal of Anxiety Disorders*, 72, 102232. <https://doi.org/10.1016/j.janxdis.2020.102232>
- Varshney, M., Parel, J. T., Raizada, N., & Sarin, S. K. (2020). Initial psychological impact of COVID-19 and its correlates in Indian Community: An online (FEEL-COVID) survey. *PLOS ONE*, 15(5), e0233874. <https://doi.org/10.1371/journal.pone.0233874>
- Wang, C., Pan, R., Wan, X., Tan, Y., Xu, L., Ho, C. S., & Ho, R. C. (2020). Immediate Psychological Responses and Associated Factors during the Initial Stage of the 2019 Coronavirus Disease (COVID-19) Epidemic among the General

Population in China. *International Journal of Environmental Research and Public Health*, 17(5), 1729. <https://doi.org/10.3390/ijerph17051729>

Yuan, K., Gong, Y.-M., Liu, L., Sun, Y.-K., Tian, S.-S., Wang, Y.-J., Zhong, Y., Zhang, A.-Y., Su, S.-Z., Liu, X.-X., Zhang, Y.-X., Lin, X., Shi, L., Yan, W., Fazel, S., Vitiello, M. V., Bryant, R. A., Zhou, X.-Y., Ran, M.-S., ... Lu, L. (2021). Prevalence of posttraumatic stress disorder after infectious disease pandemics in the twenty-first century, including COVID-19: A meta-analysis and systematic review. *Molecular Psychiatry*, 26(9), 4982–4998. <https://doi.org/10.1038/s41380-021-01036-x>

Zhang, L., Pan, R., Cai, Y., & Pan, J. (2021). The Prevalence of Post-Traumatic Stress Disorder in the General Population during the COVID-19 Pandemic: A Systematic Review and Single-Arm Meta-Analysis. *Psychiatry Investigation*, 18(5), 426–433. <https://doi.org/10.30773/pi.2020.0458>

Zhong, B.-L., Luo, W., Li, H.-M., Zhang, Q.-Q., Liu, X.-G., Li, W.-T., & Li, Y. (2020). Knowledge, attitudes, and practices towards COVID-19 among Chinese residents during the rapid rise period of the COVID-19 outbreak: A quick online cross-sectional survey. *International Journal of Biological Sciences*, 16(10), 1745–1752. <https://doi.org/10.7150/ijbs.45221>