

#	Authors and Date	Country	Study Design	Population	Natural environment(s)	Intervention or exposure	Key findings	Barriers (B) or Facilitators (F)
1	<i>Sprague N., Berrigan D., and Ekenga C. (2020)</i>	USA	Interventional study (non-randomized experiment) – with mixed-methods	Children (n=122; age range: 10-15-year-old)	Green spaces – urban forest parks, camping trips, urban farms, cave trips	NBE - Nature-Based Education	<ul style="list-style-type: none"> - Statistically significant positive changes in STEM capacity (+44%) and HRQoL (+46%) for participating students. - Qualitative data highlighted the intervention's educational and health benefits. 	<ul style="list-style-type: none"> - Age (F) - the older the child, the more active they will be, and the more benefits they will have on HRQoL. No effect for STEM. - Duration - the longer tended to have more beneficial effects on STEM capacity. - Stressful life events - the more one experiences, the less likely they will have benefits from NOEs.
2	<i>Arnberger A., Eder R., Allex B., Ebenberger M., Hutter H.P., Wallner P., Bauer N., Zaller J. and Frank T.(2018)</i>	Austria & Switzerland	Interventional study (field experiment) – using mixed-methods	Adults (workers and university students) (n=22, age mean=26.7; SD=4.1)	Both - one urban city, two meadows (managed vs unmanaged) and river in mountain area	Wilderness expedition – walking and interacting (i.e. viewing) nature	<ul style="list-style-type: none"> - While differences measured on the physiological level between urban built and natural sites were marginal (on DBP not SBP), psychological measures showed higher health benefits of the natural environments 	<ul style="list-style-type: none"> - Environment type (B/F) - river and alpine mountain meadow had the highest health benefits in terms of restoration, BP, and perception of beauty; but all sites recorded an improvement on calming and positive effect post-intervention.

							compared to the built one.	
3	Gargiulo I., Benages-Albert M., Garcia X. & Vall-Casas P. (2020)	Spain	Observational study (exploratory fieldwork) – using qualitative methods	Adults stream users (N=30; F=14 and M=16, age range: 27-65+ years old)	Green – urban stream corridor	Leisure-Time Physical Activity (LTPA)	<ul style="list-style-type: none"> - Social and physical factors of the environment are perceived as either barriers or facilitators, with different nuances and importance, depending on each type of user. - Also, for the same type of user, factors perception also depends on gender; whereby safety was important for women engaging in LTPA. 	<ul style="list-style-type: none"> - Safety (B) – women reported lower level of use of blue/green spaces during LTPA if safety was a concern. - Environmental design (B/F) - itineraries with enhanced visibility, higher attendance, pruning of dense vegetation and provisioning of assistance in case of need, all promoted engagement in LTPA. - Accessibility (F) - stream accessibility and proximity to environments is conducive for LTPA. - Presence of others (B/F) - having someone to share the experience with facilitated engagement. But

								<p>can be a barrier when perceived as safety risk (i.e. walkers vs runners/cyclists and vice-versa).</p> <ul style="list-style-type: none"> - Environment type (F)
4	Denton H. and Aranda K. (2019)	UK	Observational study (ethnography) – using qualitative methods	Adults (regular swimmers and existing club members) (n=6; F=3 and M=6, age range: 38-73 years old)	Blue space - sea swimming club (Brighton, UK)	Sea swimming	<ul style="list-style-type: none"> - The swimmers found sea swimming transformative, (resulting in changes in the swimmer's experience of themselves); connecting (experiencing a sense of connection to nature, place and others); and re-orientating (as swimmers seemed to use this disruption to reconnect to what they consider is important), through the disruption to the sense of time, space and body, swimmers find alternative and 	<ul style="list-style-type: none"> - Physical activity - engaging actively in the sea by swimming was critical to gaining health benefits (i.e. emotional and physical health) - Fear and stigma - negative body image can impact one's engagement in sea swimming; but also fear of the challenges from the sea.

							expanded perspectives about themselves and their world.	
5	Finlay J., Franke T., McKay H. and Sims-Gould J. (2015)	Canada	Observational study (participant observation) – using qualitative methods	Older adults (community dwellers; T1: N=27; T2: N=19, age range: 65-86 years old)	Both - urban parks in neighbourhood, with green and blue features	Walking	<ul style="list-style-type: none"> - Older adults have distinct therapeutic relationships with landscapes. - Nature can promote the physical, mental, and social health of older adults. - Blue space in particular embodies important therapeutic qualities for older adults. 	<ul style="list-style-type: none"> - Safety (B/F)- can be experienced differently by people. - Accessibility (B/F) - the least accessible, the worst the experience - Personal perception (B/F) - the same place could evoke feelings from enjoyment to indifference to concern due to traffic, park maintenance, walkability, etc.

6	<p>McEwan K., Richardson M., Sheffield D., Ferguson F.J. and Brindley P. (2019)</p>	UK	Interventional study (RCT) – quantitative methods	<p>Adults (18+) residing in Sheffield and owning a smartphone (N=148)</p>	Greenspace – urban park vs control (urban built)	Social prescription app on smartphone design to make people notice nature (while being in nature)	<ul style="list-style-type: none"> - Using a social prescription using a Smartphone app (noticing nature) resulted in statistically significant improvements in wellbeing for adults in general, and clinically significant improvements in wellbeing for those classed as having a mental health difficulty. - These improvements were more pronounced in the green space condition, despite improvements still in control. 	<ul style="list-style-type: none"> - Environment type (F) – both built and green environment yielded short-term benefits on wellbeing through nature connectedness; but only green space condition had sustained effects after one-month follow-up. - Previous experience with nature (F) – from childhood or in the last year, both have positive effects on wellbeing. - Positive affect (F) – predictor of wellbeing in green condition.
7	<p>Nicolosi V., Wilson J., Yoshino A. & Viren P. (2020)</p>	USA	Interventional study (field experiment) – using quantitative methods	<p>Adults (undergraduates) (n=63; F=31 and M=32;</p>	Blue spaces – the coast vs control (urban side-road)	Coastal and urban walk	<ul style="list-style-type: none"> - Significantly higher average perceived restoration scores were associated with the natural (coastal) walk. - Coastal exposure, sound quality 	<ul style="list-style-type: none"> - Perceived sound level (F) - higher perceived sound level was a significant predictor of a restorative experience. - Environment type (F) – if in natural environment, then increased

				age mean=20.4)			and type were rated as very good and more natural than the sidewalk respectively and were significant predictors of a restorative experience.	perceived restoration.
8	<i>Cheesbrough A.E., Garvin T., Nykiforuk C.I.J. (2019)</i>	Canada	Observational study (case study)– using qualitative methods	Adults and older adults (residents around one of the five selected NAP); (n=33; F=18 and M=15; age range=29-87 years)	Both – within five Natural Area Parks (blue and green features present)	Nature photography and reflection in nature	<ul style="list-style-type: none"> - Proximity to natural areas facilitated frequent and spontaneous visits. - Repeat visits fostered intimacy with the space over time. - Participants felt 'away from the city' while in the middle of the city. - Participants reported physical, spiritual and psychological therapeutic impacts. - Natural areas facilitated connections to nature, self, companions, and others. 	<ul style="list-style-type: none"> - Proximity (F) - increase engagement in physical activity and therefore promotes improved health benefits. - Topography - more difficult terrain were motivational for users of NAPs. - Sensory qualities (F) - facilitated a positive experience of being in nature, and therefore allowed for this visit to be a restorative experience. - Safety (B) - main barrier to using these areas (whether for fear of humans or wildlife).

9	<p><i>Barton J., Bragg R., Pretty J., Roberts J., and Wood C. (2016)</i></p>	<p>South Africa & Scotland</p>	<p>Interventional study (field experiment) – using quantitative methods</p>	<p>Adolescents (n=130; F=74 and M=57; age range: 11-18 years old)</p>	<p>Both – in a game reserve and a loch</p>	<p>Wilderness expedition in natural environments</p>	<ul style="list-style-type: none"> - Environment, gender, and the length and location of expeditions significantly contributed to PPTs' changes in self-esteem (SE) and nature connectedness (NC). - PPTs living in urban environments and going to local wilderness for a short duration will receive the same amount of benefits SE and NC as PPTs who live in a rural location and are immersed in a remote wilderness for longer. 	<ul style="list-style-type: none"> - Gender (F) - males had higher self-esteem at start, but significant increase in SE for females at the end. - Duration (F)- even short durations of expeditions can have benefits on nature connectedness and self-esteem.
---	--	------------------------------------	---	---	--	--	--	--

1 0	<p>Lanki T., Siponen T., Ojala A., Korpela K., Pennanen A., Tiittanen P., Tsunetsugu Y., Kagawa T. and Tyrväinen L. (2017)</p>	Finland	Interventional study (field experiment) – using quantitative methods	Adults (female volunteers in Helsinki) (n=36; age range: 30-60 years old)	Green space (vs control) - an urban forest, an urban park, and a built-up city centre	Each visit: 15 min of sedentary viewing; and 30min of walking	<ul style="list-style-type: none"> - Beneficial changes in cardiovascular physiology were observed in green environments. - Specifically, lower blood pressure (viewing period only), lower heart rate, and higher indices of heart rate variability. - Large urban park and extensively managed urban woodland had positive influence, but the overall perceived restorativeness was higher in the woodland. - This may be explained by stress relief and lower air pollution and noise exposure. 	<ul style="list-style-type: none"> - Stress relief (F) – the more relaxed one is in NOE, the better their health outcomes. - Air pollution (B) – the higher the air pollution, the worst the health outcomes. - Noise exposure (B) – higher noise engenders higher stress, and therefore lower health benefits.
--------	--	---------	--	---	---	---	--	---

1 1	Marselle M.R., Warber S.L. and Irvine K.N. (2019)	UK	Observational study – using quantitative methods	Adults (volunteers) (N=1,516; age range: 55 years or older)	Both - natural environment (i.e., natural and semi-natural places, farmland, green corridor, coastal area, urban green space, or any mixture of the above)	Nature group walks	<ul style="list-style-type: none"> - Neither nature group walking, nor doing this frequently, moderated the effects of stressful life events on mental health. - The positive associations of group walks in nature were at a greater magnitude than the negative associations of stressful life events on depression, positive affect, and mental well-being, suggesting an 'undoing' effect of nature group walks. 	<ul style="list-style-type: none"> - Stressful life events (B) – walking can help un-do stress associated with stressful life events by reducing depression and increasing positive affect and wellbeing. - Presence of others (B) - can dampen buffering effect of nature on mental health. - Physical activity (F) – mechanism by which individuals gain benefits from nature.
1 2	PÁLSDÓTTIR A.M., STIGMAR K., NORRVING B., PETERSSON I.F., ÅSTRÖM M. and PESSAH-RASMUSSEN H. (2020)	Sweden	Interventional study (RCT) – using quantitative methods	Adults and older adults (stroke survivors) (n=101; F=	Green space - Alnarp Rehabilitation Garden (Nature Area (informal and non-cultivated) and the Cultivation and Gardening	Nature-based rehabilitation (NBR) using horticultural therapy	<ul style="list-style-type: none"> - The patients with sub-acute stroke were highly compliant with the intervention. The participants in both the intervention and control groups improved. 	<ul style="list-style-type: none"> - Weather (B) - not suitable NBR in bad weather. - Access (B) - acted as barrier to participation in NBR for some PPTs due to longer travel time to the garden.

				61 and M=40; age range: 50–80 years old).	Area (formal land cultivated))		<ul style="list-style-type: none"> - However, no statistically significant differences in improvement were found between the intervention and control groups for any of the outcome measures. - Fatigue decreased to a value below the suggested cut-off for mental fatigue (< 10.5) in the intervention group, but not in the control group. 	
1 3	<i>Pratiwi P.I., Xiang Q. and Furuya K. (2019)</i>	Japan	Interventional study (field experiment) – using quantitative methods	Adults and older adults (local residents) (n=12 in spring; F=6 and M=6; mean age: 66.4) and (n=12 in summer; F=7	Green spaces – across three sites: urban city site and two viewing spots in urban park	Viewing cherry blossom trees and fresh greenery in urban parks VS urban city in Spring and August	<ul style="list-style-type: none"> - Viewing cherry blossoms and fresh greenery in urban parks led to lower blood pressure in spring and early summer than viewing city areas in spring and early summer. - The results of this study 	<ul style="list-style-type: none"> - Seasons (F) - positive mood states were higher in spring as well as lowered mood disturbances; whereas state-anxiety levels were lower in early summer. - Environment type (F) - green spaces > urban city for physiological and

				and M=5; mean age: 65.75)			<p>suggest that viewing urban parks results in physiological and psychological relaxation.</p>	<p>psychological effects of NBI.</p> <ul style="list-style-type: none"> - Biodiversity and surrounding features (B/F) (i.e. mosquitoes, sun, temperature, etc.) - thought to have increased heart rate and modified BP measurements, when viewing urban parks in both seasons. - Presence of water (F) - associated with a significant positive effect and high perceived restorativeness. - Traffic (B) (i.e. people or noises from vehicles) - could be responsible for altering BP measures in urban city.
--	--	--	--	---------------------------	--	--	--	--

1 4	Byström K., Grahn P. and Hägerhäll C. (2019)	Sweden	Interventional study (field experiment) – using qualitative methods	Children (with disabilities, i.e. autism) (n=9; mental age range: 4–6 years)	Both – immersion in a farm and surrounding nature (no specifications)	KOMSI treatment – nature therapy for treating children with disabilities (i.e. horseback riding, free play, etc.)	<ul style="list-style-type: none"> - The intervention led researchers to conclude on three key benefits of the intervention: 1) reduce stress and instill calm, 2) arouse curiosity and interest, and 3) attract attention spontaneously. - These three perceived benefits are related to vitality forms. It is argued that the vitality forms from nature and animals are favorable for effecting development-promoting interactions with a therapist. 	<ul style="list-style-type: none"> - Therapeutic environment (F/B) - if in nature it can trigger positive or negative responses for the child (i.e. not all autistic children would appreciate being out) - Presence of animals (F) - for this subgroup, animals and nature facilitated communication and alleviated stress.
1 5	Ana BY, Wanga D., Liua XJ., Guanb HM., Wei HX. and Renb ZB. (2019)	China	Interventional study (field experiment) – using quantitative methods	Adults (undergraduates in horticulture)	Green spaces – across three types of forests: 1) Maple, 2) Birch 3) Oak	Forest Bathing in three types of forests	<ul style="list-style-type: none"> - This study looked at the relationship between environmental factors (temperature, RH, light intensity, and light spectrum) 	<ul style="list-style-type: none"> - Tree species (F) - maple> oak> birch for HR improvements. Yet, birch forests still had HR improvements, and was the only one to demonstrate that at

				(n=13; M=7 and F=6; mean age: 21 years old)			<ul style="list-style-type: none"> and physiological changes (SP, DP, and HR). - Pre-forest-bathing temperature and spectrum can impact the response of blood pressure due to the "law of the initial value". - HR was influenced positively by visits to maple >oak>birch trees. - Authors recommend visitors to walk in maple forests to obtain cardiovascular and autonomic nervous system well-being. 	<p>lower levels of BP to begin with.</p> <ul style="list-style-type: none"> - Temperature (B) - Pre-forest bathing temperature can negatively impact the response of BP if PPTs felt too cool and moist. - Light spectrum (B) - pre-forest bathing spectrum can negatively impact the response of BP if high G/B ratio are too extreme.
1 6	Leavell M.A., Leiferman J. A., Gascon M., Braddick F., Gonzalez J. C. and Litt J. S. (2019)	USA	Literature review	Across age groups	Both – across several types of natural environments	Several types of NBIs being reviewed here - i.e. water rafting, horticulture, green exercise	<ul style="list-style-type: none"> - Nature-based social prescription increases social connectedness and influences physical health and mental well-being by certain 	<ul style="list-style-type: none"> - Intrapersonal processes (F) - give way to social connections and longer-term health outcomes. - Interpersonal processes (F) - improves social connections and

							<p>intrapersonal, interpersonal, and environmental pathways.</p> <ul style="list-style-type: none"> - NBI practice represents a low cost, creative intervention to strengthen social networks, reduce stress, and facilitate social connectedness among participants and providers. 	<p>health outcomes by promoting social involvement, relatedness, and shared learning.</p> <ul style="list-style-type: none"> - Environmental processes (B/F) – such as access to nature, perceived neighbourhood attachment, and perceived aesthetics.
1 7	<p>Hunter R.F., Cleland C., Cleary A., Droomers M., Wheeler B.W., Sinnette D., Nieuwenhuijse n M.J. and Braubachi M. (2019)</p>	<p>USA, Australia, UK</p>	<p>Meta-narrative evidence synthesis</p>	<p>Across groups</p>	<p>age</p>	<p>Urban green spaces – i.e. urban parks, rooftops, parking lots, etc.</p>	<p>Any NBI intervention that has only physical changes to the UGS or with health promotion to tackle inequalities</p>	<ul style="list-style-type: none"> - There was strong evidence for: 1) park-based and greenway/ trail interventions employing a dual approach (i.e. a physical change to the UGS and promotion/marketing programmes); 2) Greening of vacant lots which reduced stress and social benefits (e.g. - Changes to the built environment in parks (with dual-approach) (F) - provision of signage and community garden, improvements in existing playing fields, replacement of old playground equipment, installation of outdoor gyms, improved footpaths and clearing of rubbish and vandalism all increased

							<p>reduction in crime, increased perceptions of safety); 3) Greening of urban streets and SuDS for managing storm water had environmental benefits as well.</p>	<p>individual's park use, physical activity and the latter two improved QoL and perception of safety.</p> <ul style="list-style-type: none"> - Proximity to newly developed walking/cycling routes (F) - increased use of these UGS. - Greening of vacant lots (F) – reduced perception of unsafe environment and bolster use of these UGS.
18	<p><i>van den Bosch M. and Sang O. (2017)</i></p>	Not provided	Systematic review (of reviews)	Across age groups	Both - (i.e. green infrastructure, biodiversity, blue environments, etc.)	Interventions in urban natural environments	<ul style="list-style-type: none"> - There is strong evidence on the effect of urban nature on affect state. - There is strong evidence on the effect of urban nature on urban heat reduction. - Positive affect and heat reduction can mediate urban nature's effect on mortality. 	<ul style="list-style-type: none"> - Micro-features (F) - Conditions of natural environments (B/F) - Perceived quality (F) - Accessibility (B/F) - Safety (B)

1 9	<p>Houlden V., Weich S., de Albuquerque J.P., Jarvis S. and Rees K (2018)</p>	Europe USA Canada, Australia	Systematic review	Adolescents, adults, and older adults	Green spaces – mixed definition that encompasses vegetated areas and/or wilderness	Not interventions per se, but includes studies with walking in GS as measure for visits to GS	<ul style="list-style-type: none"> - There was adequate evidence for associations between the amount of local-area greenspace and life satisfaction (hedonic wellbeing), but not personal flourishing (eudaimonic wellbeing). - Evidence for associations between mental wellbeing and visits to greenspace, accessibility, and types of greenspace was limited. - There was inadequate evidence for associations with views of greenspace and connectedness to nature. 	<ul style="list-style-type: none"> - Views of greenspace (B/F) - if looking at unpleasant urban/rural views it will have negative association with mental health. - Connection with nature (F) - the more connected one is with nature, the more health benefits (i.e. life satisfaction, happiness, affect, QoL) one will experience. This is modulated by being actively engage in nature, however. - Visits to greenspaces (F) - active immersion in wilderness was found to lead to greater happiness, affect and attention.
--------	---	------------------------------------	----------------------	---	--	---	---	---

20	McCormick R. (2017)	USA, Spain and others not specified	Systematic review	Children (age range: 0-18)	Green spaces – wooded playgrounds, natural habitats, gardens, etc.	Not interventions per se but does include studies who used walking in NOE as measure for visits to green space.	<ul style="list-style-type: none"> - Access to green space is important to the mental well-being, overall health, and cognitive development of children. It promotes attention restoration, moderates the impacts of stress, improves behaviours and symptoms of ADHD and was even associated with higher standardized test scores. 	<ul style="list-style-type: none"> - Proximity to GS (F) - only passive exposure but the closer one lives to nature, the better health outcomes they have. - Physical activity in NOE (F) - walking in nature for children vs in urban environment led to improved attention and spatial working memory - which can help children with ADHD focus better.
21	Barrett J., Evans S. and Mapes N. (2019)	UK	Literature review	Older adults (residents of dementia care settings)	Green space – garden areas within dementia care settings	Green dementia care - includes many type of NBIs (i.e. horticulture, walking, gardening, etc.)	<ul style="list-style-type: none"> - Compelling evidence for several health and wellbeing benefits associated with green dementia care (i.e. improved wellbeing, social interactions, stress-reduction and restorative effects, self- 	<ul style="list-style-type: none"> - Safety concerns (i.e. fear of falling in garden) (B) - Staff attitudes and lack of staff education and awareness (B) - Social prejudice and stigma (B) - Limited staff to accompany residents and limited resources (B) - Weather (B)

						<p>worth and confidence.)</p> <ul style="list-style-type: none"> - Evidence base is stronger regarding the barriers and facilitators to accessing nature for this population - staff education and care culture is critical to the success and effective use of the garden for such residents. - Design of the outdoor space need to ensure that these spaces are visually and physically accessible for its residents. 	<ul style="list-style-type: none"> - Self-perception of being too old and lack of confidence (B) - Poor physical and visual access - Poor garden design (B) (i.e. lack of resting places and weather protection) - Care culture NOT person-centred (B)
2 2	<p>Otoni C.A., Sims-Gould J., Winters M., Heijnen M. and McKay H.A. (2016)</p>	Canada	<p>Observational study (participant observation) – using</p>	<p>Older adults (60+) (n=28; F=17 and M=11, age range: 61-89, in 2012; n</p>	<p>Both areas – three areas in parks in Vancouver with features of GS and BS</p>	<p>Physical activity in nature - recorded as step counts/day (mean)</p> <ul style="list-style-type: none"> - Neighbourhood environments influence health and well-being as people age. - There are strong interconnections between built 	<ul style="list-style-type: none"> - Amenities (F) – i.e. benches, seen as a necessity to promote social interactions and positive experiences. - Ability to engage in other type of activities (F) – i.e.

			<p>qualitative methods</p>	<p>=22, F=12 and M= 10; in 2014)</p>			<p>and social environments.</p> <ul style="list-style-type: none"> - Microscale features can enable older adults' to accommodate to their abilities. - Benches can promote mobility and social connectedness for older adults. - Microscale features, like benches, are a prudent investment for communities. 	<p>family or friends activities, going to the pub, going to the gym featured more prominently than benches in relation to their mobility.</p> <ul style="list-style-type: none"> - Injury (B) - to use the outdoor environments. - Wildlife (F) - promoted feelings of enjoyment and calmness. Also helped in creating routines/familiarity with these spaces. - Presence of other people (B/F) - a negative experience for older adults if too many people use benches. But seeing people around them also provided positive feelings opportunities for social interactions. - SES (B) - accessibility and availability of GS and BS was more common for older adults with higher
--	--	--	----------------------------	--------------------------------------	--	--	--	---

								social and economic status.
2 3	Howarth M., Rogers M., Withnell N. and McQuarrie C. (2018)	UK	Observational study (cross-sectional) – using mixed-methods	Adults and older adults (suffering from mental disorders (n=47; age range: 35-68 years and average age: 53.2 years)	Green space – garden area created by social enterprise	Therapeutic horticulture	<ul style="list-style-type: none"> - Quantitative findings showed that participants were working towards self-reliance. Qualitative data found similar results. - Mental health recovery programme enabled participant integration into the community through providing a space to grow and build self-confidence while reengaging with society. - The results suggest that using therapeutic horticulture as 	<ul style="list-style-type: none"> - Positive staff attitudes (F) – welcoming and non-judgmental attitudes promoted wellbeing and social connection for this population. It also helped people feel safer. - Activities as a new purpose (F) – engaging in nature itself improved wellbeing and allowed people to feel a sense of purpose. By developing new skills people felt more confident in their own self and their employability. - Presence of others (F) – improved sense of purpose and recovery, as everybody shared

							an intervention within the mental health recovery programme can support people with mental health problems to re-engage socially.	same/similar experiences; this helped them move beyond their diagnosis. It also provided opportunities to re-engage with society.
2 4	<i>Kabisch N., Matilda van den Bosch M. and Lafortezae R. (2017)</i>	U.S., Germany France Spain Denmark, Bulgaria, Austria, Sweden, UK, Japan, Canada and China	Systematic Review	Children and the elderly	Both – features of both GS and BS	Some studies included interventions / active engagement with nature	<ul style="list-style-type: none"> - There is a tendency for a positive association between urban green and blue spaces and reduced risk factors related to urbanization for children and the elderly as well as the promotion of health-related behaviours and subsequent positive health outcomes. - But the evidence is weak and the results are somewhat inconsistent. 	<ul style="list-style-type: none"> - Socioeconomic factors (B) (i.e. deprivation, income, educational level, unemployment) - the lower one's household, the worst their health outcomes, and the lower the relationship between health and nature. - Air pollution (B) - act as mediator of the relationship between nature and health - but not if elderly engage actively in NOE (i.e. gardening). - Heat-related pollution (B) - the higher the heat in parks, the less use and the worst health outcomes,

								specifically for the elderly. - Proximity/distance (F/B) - proximity can modify effectiveness of NBIs.
2 5	<i>Shin J.C., Parab K.V., An R. and Grigsby-Toussaint D.S. (2020)</i>	USA, Australia, Canada, Spain, UK, Netherlands, Lithuania	Systematic review	Across age groups	Green spaces – neighbourhood greenness, visits to GS, engagement in activities related to GS	Several interventions included: walking, gardening, work environment	- Green space exposure (through active engagement) is associated with better sleep quality and quantity. - Authors suggest green exercise and therapeutic gardening for future interventions.	- Time of day for activity (F) – afternoon walking > morning walks. - Type of environment (F) - outdoor > indoor interventions on sleep latency. - Behavioural preferences (F) - people had better sleep on weekdays with lower exposure to green spaces. Also preferred vaster expenses of greenspace on weekends and for longer than during weekdays.
2 6	<i>Lakhani A., Norwood M., Watling D.P., Zeeman H. and Kendall E. (2019)</i>	USA, Norway, Netherlands, Australia, Korea, Japan	Systematic review	Adults and older adults (suffering from neurological disability:	Both – includes studies with features from both green and blue environments	Several interventions included: gardening, green care farming, wilderness therapy,	- Given the limited research to date, and the diversity of nature specific activities, it is not possible to establish	- Environment type (F) - for care farming, evidence is mixed on social health. - Garden design (F) - gardens often well arranged, walled

				dementia, stroke, and brain injury)		forest therapy	<p>definitive conclusions around the efficacy of engaging with nature specific activities on the psychosocial health of people with neurological disability.</p> <ul style="list-style-type: none"> - At best, findings clarify that engaging with natural environments contribute to favourable emotional health outcomes and social health outcomes for people with dementia. 	<p>and preferably in connection with a shielded dementia care unit, may also improve agitation among people with dementia.</p> <ul style="list-style-type: none"> - Mobility (B) - impact of wander gardens on agitation reduction was lower if PPT had ambulatory issues. - Presence of caregiver (F) - brought positive emotional health outcomes in patients with dementia, when entering garden. - Short-term plants (F) –associated with improved social health vs long-term plants. Possibly due to faster harvesting capacity. - Active engagement (F) - psychological health has been favourably impacted only when activities in nature
--	--	--	--	-------------------------------------	--	----------------	--	--

								involve active engagement.	
2 7	<p>Kondo M.C., Fluehr J.M., McKeon T. and Charles C. Branas C.C. (2018)</p>	<p>USA, UK, Netherlands , Canada, Lithuania, Denmark, Germany, Finland, Japan, Italy, Spain</p>	<p>Systematic review</p>	<p>Across groups</p>	<p>age</p>	<p>Green spaces – natural environment</p>	<p>Several types of interventions including: viewing nature, walking, exercising, gardening</p>	<ul style="list-style-type: none"> - This review of experimental, quasi-experimental, and longitudinal studies found evidence of a positive association between urban green space and attention, mood, and physical activity, and negative association with mortality, short-term cardiovascular markers (heart rate), and violence. - In most cases, it is not possible to observe patterns of findings of association between urban green space exposure and health outcomes (i.e. birth outcomes, 	<ul style="list-style-type: none"> - Environment type (F) - natural environment > urban built environment for attention, general health, cardiovascular outcomes (i.e. HR, HRV), mood and emotions (i.e. specifically urban woodlands for restoration). - Biodiversity (F) - found to improve mood and emotions but is mediated by length of park visit and perceived restoration. - Physical activity (F) - engagement in PA in nature was positively associated with health outcomes in experimental studies VS observational studies.

							stress, BP, cancer, diabetes, etc.).	
28	<p>Callaghan A., McCombe G., Harrold A., McMeel C., Mills G., Moore-Cherryb N. and Cullen W. (2020)</p>	<p>Australia, USA, UK, Bulgaria, Belgium, Denmark, Netherlands, Serbia</p>	<p>Scoping review</p>	<p>Across age groups</p>	<p>Green spaces – urban parks, neighbourhood greenness</p>	<p>Several interventions included: horticultural therapy, walking, viewing from indoors</p>	<ul style="list-style-type: none"> - The majority of studies found a positive association between GS and mental health. - Policies to increase urban green space may have sustainable public health benefits. 	<ul style="list-style-type: none"> - Ethnicity (B) - South Asian children living in more deprived areas and with lower access and quality of GS had more behavioural difficulties VS white British children. - Deprivation (B) - quality and access to greenspaces is lower in deprived and lower-income communities,

								leading to worse health outcomes.
29	<p>Koselka E.P.D., Weidner L.C., Minasov A., Berman M.G., Leonard W.R., Santoso M.V., de Brito J.N., Pope Z.C., Pereira M.A. and Horton T.H. (2019)</p>	USA	Interventional study (pilot study) – using quantitative methods	Adults (undergraduates, graduates and employees) (n=37; 18–35 years; age mean=22.9 and SD=4.6)	Green	Walking in nature across 3 settings: forest; along roadside, activities of daily living	<ul style="list-style-type: none"> - This study has found that moderate-intensity walking in a forested environment had a positive impact on psychological health. - This suggests that completing physical activity in greenspaces amplifies beneficial acute psychological responses and yields greater improvements in mental health than does activity completed indoors or in a built urban environment. 	<ul style="list-style-type: none"> - Type of environment (F) - forest walking >roadside> daily activities for positive/negative affect, perceived stress, and anxiety. - Physical activity (F) - walking in general brought improved mental health but amplified in forest environment.
30	Zufferey J. (2016)	Japan, Australia, China, USA, New			Both – GS and BS elements (not specific)	Not specified per se, but includes studies with walking/	<ul style="list-style-type: none"> - This literature review shows moderate to strong empirical evidence for the 	<ul style="list-style-type: none"> - Age (F) - children and young adults seemed to benefit more from exposure to GS and

		Zealand, Canada	Systematic review	Across groups	age	exercise in GS and BS	<p>positive influence of contact with green and blue spaces and mental and physical health and low evidence for influences on social cohesion.</p> <ul style="list-style-type: none"> - It also shows that health impacts may vary according to the population group considered (e.g. children, people with low socio-economic status who benefit more from exposure to these environments). 	<p>BS, especially via physical activity; which had combined effects on physical and mental health.</p> <ul style="list-style-type: none"> - SES (B/F) - lower SES households tended to have more health benefits associated with exposure to green and blue spaces. - Type of environment (F) - natural environments > urban built on emotional wellbeing.
3 1	Costello L., McDermott M- L., Patela P. and Dare J. (2019)	Australia			Blue – the ocean and surrounding beaches	Ocean swimming	<ul style="list-style-type: none"> - All the ocean swimming groups studied were united by their routine of beach swimming, by their love of the ocean, and their conviction that 	<ul style="list-style-type: none"> - Type of environment (F) - swimming in the ocean VS public/private pool. - Biodiversity (B/F) - when seeing fishes, dolphins, whales, etc. people experienced

			Observational study (ethnography) – using qualitative methods	Older adults (self-organised ocean swimmers) (n=10; F=7 and M=10; age range: 55-80+ years)			their ocean swimming practice as part of a group was beneficial for their social connectedness, wellbeing and physical and mental health.	positive emotions and pleasurable experiences. However fear of sharks led to negative emotions, despite increasing social cohesion. - Type of activity (F) - swimming > other type of outdoor exercise, as it was low-impact. It would also help in alleviating stress. - Weather (F) - despite cold and rainy weather, PPTs would still engage in swimming, as their commitment to the group was the priority. - Group membership (F) - PPTs recognised that they would not derive the same enjoyment, pleasure and health benefits w/o group.
3 2	<i>Birch J., Rishbeth C. and Payne S.R. (2020)</i>	UK		Adolescents and adults (n=24;	Both – urban parks in Sheffield (UK)	Arts workshop and interviews in nature	- Deteriorating landscapes, young people's shifting identities and perceived time pressures	- Poor quality of urban environment (B) – or urban deprivation, was more important than

			Observational study (case study) – using qualitative methods	F=14 and M=10; age range: 17-27 years, with n=9 experiencing mental difficulties and n=15 living in an area of urban deprivation).			<p>disrupted support.</p> <ul style="list-style-type: none"> - Overall young people expressed how urban nature encounters were experienced as accepting and relational, offering a stronger sense of self; feelings of escape connection and care with the human and non-human world. 	<p>ethnicity and SES across PPTs.</p> <ul style="list-style-type: none"> - Presence of others (F/B) - having someone with you during a visit to an urban environment was positively experienced (i.e. wanting someone to share experience with), or negatively experienced (i.e. wanting to be alone). - Individual factors (B/F) - pressures, changing priorities and changing friendships all have their mediating role.
3 3	Wood E., Harsant A., Dallimer M., ronin de Chavez A., McEachan R.R.C. and Christopher Hassall (2018)	UK	Observational study (cross-sectional) – using quantitative methods	Adults and older adults (users of local parks in deprived areas) (n=128; F=59 and M=69; age range: 18-76+ years)	Green	<p>Visits to greenspaces</p> <ul style="list-style-type: none"> - survey conducted at the park entrance 	<ul style="list-style-type: none"> - Authors found that biodiversity and site facilities were positively correlated within urban parks. However, we found that only biodiversity was related to perceptions of psychological restoration amongst a multi- 	<ul style="list-style-type: none"> - Biodiversity (F) - the more biodiversity a park had; the more people would benefit from psychological restoration. - Amenities - positively correlated with urban parks, but no effects on psychological restoration.

							<p>ethnic group of PPTs.</p> <ul style="list-style-type: none"> - These findings suggest that urban planners should aim to enhance ecological diversity in urban green spaces. 	<ul style="list-style-type: none"> - Ethnicity - no effects found.
3 4	<p>Corazon S.S., Sidenius U., Poulsen D.V. , Gramkow M.C. and Stigsdotter U.K. (2019)</p>	<p>Europe, Asia, Australia</p>	<p>Systematic Review (without meta-analysis)</p>	<p>Adults and older adults</p>	<p>Both – all types of outdoors natural green environments</p>	<p>All types of sedentary and light exercise activities, in all time durations in nature</p>	<ul style="list-style-type: none"> - The synthesis of the results points towards outdoor, nature-based exposure having a positive effect on different emotional parameters, related to stress relief. The studies into physiological measures showed more equivocal results. - The general use of self-referred individuals imposes a potential strong bias. 	<ul style="list-style-type: none"> - Type of environments (F) – natural environments vs control had positive association with emotional outcomes (i.e. positive affect, perceived stress and wellbeing/QoL), and negative association with negative affect. This could not be found for physiological measures – (too heterogeneous).

3 5	<p><i>Maund Pgreen.R., Irvine K.N., Reeves J., Strong E., Cromie R., Dallimer M. and Davies Z.G. (2019)</i></p>	UK	Interventional study (pilot study) – using mixed-methods	<p>Adults and older adults (already registered with the community mental wellbeing service and diagnosed with depression and/or anxiety) (n=16; F=8 and M=8)</p>	Blue wetlands	<p>– Wetland NBI – guided walking, bird watching or other activities (i.e. canoeing) done in nature over a six-week period.</p>	<ul style="list-style-type: none"> - There were significant improvements in mental health across a range of indicators, including mental wellbeing, anxiety, stress and emotional wellbeing. Participants and healthcare professionals cited additional outcomes including improved physical health and reduced social isolation. - The wetland site provided a sense of escape from participants' everyday environments, facilitating relaxation and reductions in stress. - Wetland staff knowledge of the natural world, 	<ul style="list-style-type: none"> - NBI design (F) - to be successful, NBIs need to take into account transportation, staff knowledge and group dynamics. - Biodiversity (F) - the presence of water, diverse wildlife and the inherent peacefulness of wetlands were positively experienced with the intervention. - Session content (B/F) - most PPTs preferred if there was only ONE activity vs many → less stress and anxiety that way.
--------	---	----	--	--	---------------	--	--	---

							transportation and group organisation also played a role in the intervention's success.	
36	Benjamin-Neelon S.E., Platt A., Bacardi-Gascon M., Armstrong S., Neelon B., Jimenez-Cruz A. (2019)	Mexico	Observational study (cross-sectional) – using quantitative methods	Children (n=102, age range: 3-5 years; in Ensenada (M=29 and F=21) and in Tijuana (F=27 and M=25)	Green spaces – urban parks in two cities (Tijuana and Ensenada)	Time spent in GS (measured with GPS)	<ul style="list-style-type: none"> - Greater time in greenspace was associated with decreased sedentary time. - Greater time in greenspace was associated with increased physical activity. - Associations were mainly driven by children in Tijuana compared to Ensenada. - Time spent in greenspace was not associated with body mass index (BMI) z-score. 	<ul style="list-style-type: none"> - Duration (F) - the greater the time spent in greenspace, the less sedentary time these children will experience, but also the greater their MVPA will be (physical activity).
37	Coventry P.A., Neale C. Dyke A., Pateman R.	UK		Adults and older adults (conservation volunteers) (n=45; F=20)	Green spaces – across three sites: Askam Bog, St Nicks natural reserve and a large green field with	Three interventions : group walking, conservation	<ul style="list-style-type: none"> - Undertaking purposeful activity in public green space has the potential to promote health and prevent 	<ul style="list-style-type: none"> - Location (F)- effects differed for stress across locations, meaning that the location of the GS, over the type of activity, was

	and Cinderby S. (2019)		Interventional study (field experiment) – using mixed-methods	and M=25; age mean: 43.8)	surrounding woodland, adjacent to a semi-urban housing development	, citizen science	<p>mental ill health. Undertaking such activities in locations where people have the most connection might confer additional benefits.</p> <ul style="list-style-type: none"> - Social interaction, physical activity and restoration were all implicated as potential mechanisms by which activities in public green spaces might lead to improved mental health. 	<p>an important factor in reducing stress - which was explained by an enhanced place attachment and place identity at this location.</p> <ul style="list-style-type: none"> - Type of activity (F) - although not shown quantitatively, conservation and citizen science were both associated with deeper sense of purpose by providing learning opportunities, and because it conferred co-benefits to health, wellbeing and to nature itself.
3 8	Britton E., Kindermann G., Domegan C. and Carlin C. (2018)	Europe, USA, Canada, New Zealand, Israel	Systematic review	Across all age groups – but with pre-existing condition	Blue space – wilderness, sea, urban/semi-urban areas (beach, city), or mix of these	Several interventions included in BS: surfing, Dragon Boat Racing (DBR), sailing fly fishing kayaking, canoeing, at the beach, swimming,	<ul style="list-style-type: none"> - Blue care have direct benefit mental health and psycho-social wellbeing. - There was also evidence for greater social connectedness during and after interventions, but results were inconsistent 	<ul style="list-style-type: none"> - Access (F/B) - Lack of resources/equipment (B) - Fears/stigma (B) - associated with personal abilities, level of fitness, environment, social and cultural norms, diagnosis of illnesses and level of appropriate training for those

						(as part of a kayaking intervention), and scuba diving	<ul style="list-style-type: none"> - and mixed; with very few findings for physical health. - Findings suggest how activities in BS, rather than particular qualities of BS, might contribute to rehabilitation and health promotion. 	<ul style="list-style-type: none"> - delivering intervention. - Gender - Seasickness - due to poor weather (B) - Fatigue/tiredness - post-intervention (B)
3 9	<p>Saadi D., Schnell I., Tirosh E., Basagaña X. and Agay-Shay K. (2020)</p>	Israel	Interventional study (field experiment) – using quantitative methods	<p>Adults (women specifically) (n=120; age range: 20-35, from two small cities in the north of Israel, of whom n=48 were Arab and n=24 were Jewish women (n= 72)</p>	<p>Green spaces – across Afula-urban park, Afula-city center, Afula-residential area, Nazareth urban park, Nazareth-city center and Nazareth-residential area VS home (control)</p>	<p>Viewing and waling in nature while sitting on benches across 6 different sites</p>	<ul style="list-style-type: none"> - Visits to urban parks compared to staying in the home environment had beneficial short-term changes in psychological, physiological, and cognitive responses, regardless of ethnicity. - The changes could not be attributed to the investigated mediators. - Women should be encouraged to go outdoors and specifically visit parks to 	<ul style="list-style-type: none"> - Environment type (F/B) - Arab woman demonstrated improvement in most outdoor environments, while for the Jewish woman, improvement was reported mainly in parks, but not in any other urban environment. - Socio-demographic (ethnic preferences) (F) - whereby benefits were stronger in intra-ethnic parks. - Comfort level at home (F/B) - more comfort at home for Jewish women vs

								improve their psychological and physiological health	Arab women, which could have reduced positive effects of outdoor environments considered less comfortable for this subgroup.
--	--	--	--	--	--	--	--	--	--