

Supplementary Information

The impact of human mobility and reactive case detection on malaria transmission in Zanzibar

Aatreyee M. Das^{1,2}, Manuel W. Hetzel^{1,2}, Joshua O. Yukich³, Logan Stuck⁴,
Bakar S. Fasih^{1,2,5}, Abdul-wahid H. Al-mafazy⁶, Abdullah Ali⁷,
Nakul Chitnis^{1,2}

1 Department of Epidemiology and Public Health, Swiss Tropical and Public Health Institute,
Basel, Switzerland

2 University of Basel, Basel, Switzerland

3 Center for Applied Malaria Research and Evaluation, Department of Tropical Medicine,
Tulane University School of Public Health and Tropical Medicine, New Orleans, LA, USA

4 Amsterdam Institute for Global Health and Development, Amsterdam, Netherlands

5 Health Systems, Impact Evaluation and Policy Department, Ifakara Health Institute,
Dar es Salaam, United Republic of Tanzania

6 RTI International, Dar es Salaam, United Republic of Tanzania

7 Zanzibar Malaria Elimination Programme, Zanzibar, United Republic of Tanzania

S1 Methods

S1.1 The controlled reproduction number for the whole system

The controlled reproduction number for the whole system, R_s , gives the expected number of secondary infections arising across all three patches from a primary infection when interventions are in place. R_s was calculated by taking the spectral radius of the next generation matrix [1, 2]. Note, R_s is different to the local controlled reproductive number on each island given in the main text, R_c , as that is calculated by taking the ratio of the transmission and recovery rates on each island.

The equation for the rate of change of infected individuals (equation (4) in the main text) can be linearised by decomposing the Jacobian matrix into two matrices describing the transmission events leading to new infections, \mathbf{F} , and the recovery events leading to removal from the infected class, \mathbf{V} :

$$\frac{d\vec{I}}{dt} = (\mathbf{F} - \mathbf{V})\vec{I}, \quad (1)$$

where

$$F_{ij} = \sum_{k=1}^3 \left(\frac{\beta_k N_j \theta_{kj} \theta_{ki}}{\sum_{l=1}^3 N_l \theta_{kl}} \right), \quad (2)$$

and

$$V_{ij} = \begin{cases} \mu & i = j, \\ 0 & i \neq j. \end{cases} \quad (3)$$

The next generation matrix, \mathbf{K} , is given by

$$\mathbf{K} = \mathbf{F}\mathbf{V}^{-1}. \quad (4)$$

R_s is equal to the spectral radius of \mathbf{K} .

In order to investigate the potential impact of changing transmission levels on mainland Tanzania, the value of β_{mainland} was varied between 0.004 and 0.0054 and R_s was re-calculated.

The results are shown in Fig. S1. While the value for the transmission rate on mainland Tanzania is below the transmission rate on Pemba, the spectral radius of \mathbf{K} is dominated by the ratio of the transmission and recovery rates on Pemba. Once the transmission rate on the mainland approaches that of Pemba, the mainland transmission rate becomes the dominant factor in determining the spectral radius of \mathbf{K} . After this point, we see a linearly increasing relationship between the mainland transmission rate and the controlled reproductive number of the system.

S1.2 Treatment of imported cases

Treatment of imported cases leads to a shift in the equilibrium prevalence observed on each island. Fig. S2 shows the timeseries plots for reaching equilibrium for the following proportions of treatment of outbound travellers from the mainland to Zanzibar, O , and travellers from Zanzibar on their return to Zanzibar, R , assuming the baseline level of RCD is maintained:

- $O = R = 0.25$,
- $O = R = 0.5$,
- $O = R = 0.75$,
- $O = R = 1$.

S2 Figures




Figure S1: Plot showing the relationship between the transmission rate on mainland Tanzania, and the controlled reproductive number of the whole system.




Figure S2: 7-day moving average of the median of 500 stochastic simulations for an SIS model of RCD for Pemba and Unguja comparing different levels of treating infected travellers, assuming baseline RCD is maintained. Here, the value given as the proportion of travellers treated account for both travellers from the mainland and travellers from Zanzibar.

References

- [1] Julie C Blackwood and Lauren M Childs. An introduction to compartmental modeling for the budding infectious disease modeler. *Letters in Biomathematics*, 5(1):195–221, 2018.
- [2] Odo Diekmann, JAP Heesterbeek, et al. The construction of next-generation matrices for compartmental epidemic models. *Journal of the Royal Society Interface*, 7(47):873–885, 2010.