

Supplementary material

This material accompanies the article

Telomere length and its associations with mental disorders, age and genetic risk for mental disorders

Authors: Julian Mutz and Cathryn M. Lewis

Table of contents:

GWAS summary statistics	2
T/S ratio in individuals with mental disorders	3
T/S ratio in individuals with mental disorders by antidepressant medication	4
Figure: T/S ratio by antidepressant medication	4
Table: T/S ratio by antidepressant medication	5
Polygenic risk scores in individuals with mental disorders	6
Associations between T/S ratio and polygenic risk scores	7
Scatter plots	7
Table	8
Sensitivity analysis.....	9
Figure: T/S ratio, no comorbid depression and anxiety disorders.....	9
Table: T/S ratio, no comorbid depression and anxiety disorders	10
Sensitivity analysis stratified by antidepressant medication.....	11
Figure: T/S ratio, no comorbid depression and anxiety disorders by antidepressant medication	11
Table: T/S ratio, no comorbid depression and anxiety disorders by antidepressant medication.....	12


GWAS summary statistics

Supplementary Table 1. GWAS summary statistics

Phenotype	GWAS reference	Sample size	
		Cases	Controls
Anxiety disorders	Otowa et al. (2016) doi: 10.1038/mp.2015.197	7016	14745
Major depressive disorder	Wray et al. (2018)* doi: 10.1038/s41588-018-0090-3	45591	97674
Bipolar disorder	Stahl et al. (2019) doi: 10.1038/s41588-019-0397-8	20352	31358

Note: GWAS = genome wide association study. *Summary statistics from Wray et al. (2018) excluding UK Biobank and 23andMe data.


T/S ratio in individuals with mental disorders



Supplement Figure 1. Average T/S ratio (log z adjusted) in individuals with mental disorders and non-psychiatric controls. Horizontal lines show group means. T/S ratio values below the 0.01st or above the 99.99th percentile not shown.

T/S ratio in individuals with mental disorders by antidepressant medication

Figure: T/S ratio by antidepressant medication



Supplement Figure 2. Average T/S ratio (log z adjusted) in individuals with mental disorders compared to non-psychiatric controls (reference group) stratified by antidepressant medication use. Estimates shown are ordinary least squares regression beta coefficients and 95% confidence intervals. Model 1 – adjusted for age and sex; Model 2 – adjusted for age, sex, white blood cell count, Townsend deprivation index, physical activity, smoking status, body mass index, body fat percentage and C-reactive protein.


Table: T/S ratio by antidepressant medication

Supplement Table 2. T/S ratio (log z adjusted) in individuals with mental disorders by antidepressant medication

Term	Model 1					Model 2				
	β	95% CI		$p_{\text{Bonf.}}$	p_{BH}	β	95% CI		$p_{\text{Bonf.}}$	p_{BH}
Controls	Ref	-	-	-	-	Ref	-	-	-	-
Anxiety disorder	-0.001	-0.012	0.011	>0.999	0.898	0.003	-0.008	0.014	>0.999	0.712
+ Antidepressant	-0.043	-0.063	-0.022	<0.001	<0.001	-0.024	-0.045	-0.004	0.264	0.053
Depression	-0.015	-0.024	-0.007	0.005	0.001	-0.007	-0.016	0.001	>0.999	0.188
+ Antidepressant	-0.049	-0.064	-0.035	<0.001	<0.001	-0.026	-0.041	-0.012	0.005	0.001
Bipolar disorder	-0.028	-0.067	0.011	>0.999	0.235	-0.011	-0.05	0.028	>0.999	0.712
+ Antidepressant	-0.043	-0.102	0.016	>0.999	0.235	-0.012	-0.071	0.047	>0.999	0.757

Note: β = ordinary least squares regression beta coefficient; CI = confidence interval; Ref = reference group; Bonf. = Bonferroni; BH = Benjamini & Hochberg. Model 1 – adjusted for age and sex; Model 2 – adjusted for age, sex, white blood cell count, Townsend deprivation index, physical activity, smoking status, body mass index, body fat percentage and C-reactive protein. *P*-values corrected for 12 tests.


Polygenic risk scores in individuals with mental disorders



Supplement Figure 3. Differences in polygenic risk scores for anxiety disorder (left panel), depression (middle panel) and bipolar disorder (right panel) between individuals with these disorders and non-psychiatric controls.

Associations between T/S ratio and polygenic risk scores

Scatter plots



Supplement Figure 4. Associations between average T/S ratio (log z adjusted) and polygenic risk scores for anxiety disorder, depression and bipolar disorder. T/S ratio values below the 0.01st or above the 99.99th percentile not shown.

Table


Supplement Table 3. Associations between T/S ratio (log z adjusted) and polygenic risk scores for mental disorders.

PRS	Full sample					Cases					Controls				
	β	95% CI		$p_{\text{Bonf.}}$	p_{BH}	β	95% CI		$p_{\text{Bonf.}}$	p_{BH}	β	95% CI		$p_{\text{Bonf.}}$	p_{BH}
Anxiety disorder	-0.002	-0.006	0.001	0.589	0.196	-0.006	-0.015	0.004	>0.999	0.350	-0.002	-0.006	0.002	>0.999	0.385
Depression	-0.006	-0.010	-0.003	0.001	0.001	-0.006	-0.012	0.001	0.592	0.236	-0.008	-0.012	-0.004	0.001	0.001
Bipolar disorder	0.003	-0.001	0.008	0.342	0.171	-0.003	-0.036	0.030	>0.999	0.858	0.003	-0.001	0.008	0.707	0.236

Note: PRS = polygenic risk score; β = ordinary least squares regression beta coefficient; CI = confidence interval; Ref = reference group; Bonf. = Bonferroni; BH = Benjamini & Hochberg. All analyses were adjusted for the first six ancestry-informative population principal components, batch number and assessment centre. *P*-values adjusted for three (full sample) and six (cases and controls) tests.

Sensitivity analysis

Figure: T/S ratio, no comorbid depression and anxiety disorders



Supplement Figure 5. Average T/S ratio (log z adjusted) in individuals with anxiety disorders or depression compared to non-psychiatric controls (reference group), excluding individuals with comorbid anxiety disorders and depression. Estimates shown are ordinary least squares regression beta coefficients and 95% confidence intervals. Model 1 – adjusted for age and sex; Model 2 – adjusted for age, sex, white blood cell count, Townsend deprivation index, physical activity, smoking status, body mass index, body fat percentage and C-reactive protein.

Table: T/S ratio, no comorbid depression and anxiety disorders


Supplement Table 4. Average T/S ratio (log z adjusted), no comorbid depression and anxiety disorders.

Term	Model 1					Model 2				
	β	95% CI		$p_{\text{Bonf.}}$	p_{BH}	β	95% CI		$p_{\text{Bonf.}}$	p_{BH}
Controls	Ref	-	-	-	-	Ref	-	-	-	-
Anxiety disorder	0.003	-0.013	0.019	>0.999	0.741	0.004	-0.012	0.021	>0.999	0.741
Depression	-0.028	-0.037	-0.018	<0.001	<0.001	-0.015	-0.025	-0.006	0.004	0.002

Note: β = ordinary least squares regression beta coefficient; CI = confidence interval; Ref = reference group; Bonf. = Bonferroni; BH = Benjamini & Hochberg. Model 1 – adjusted for age and sex; Model 2 – adjusted for age, sex, white blood cell count, Townsend deprivation index, physical activity, smoking status, body mass index, body fat percentage and C-reactive protein. *P*-values corrected for four tests.

Sensitivity analysis stratified by antidepressant medication

Figure: T/S ratio, no comorbid depression and anxiety disorders by antidepressant medication



Supplement Figure 6. Average T/S ratio (log z adjusted) in individuals with anxiety disorders or depression compared to non-psychiatric controls (reference group) stratified by antidepressant medication use, excluding individuals with comorbid anxiety disorders and depression. Estimates shown are ordinary least squares regression beta coefficients and 95% confidence intervals. Model 1 – adjusted for age and sex; Model 2 – adjusted for age, sex, white blood cell count, Townsend deprivation index, physical activity, smoking status, body mass index, body fat percentage and C-reactive protein.

Table: T/S ratio, no comorbid depression and anxiety disorders by antidepressant medication

Supplement Table 5. Average T/S ratio (log z adjusted), no comorbid depression and anxiety disorders, by antidepressant medication

Term	Model 1					Model 2				
	β	95% CI		$p_{\text{Bonf.}}$	p_{BH}	β	95% CI		$p_{\text{Bonf.}}$	p_{BH}
Controls	Ref	-	-	-	-	Ref	-	-	-	-
Anxiety disorder	0.005	-0.012	0.022	>0.999	0.618	0.006	-0.011	0.023	>0.999	0.618
+ Antidepressant	-0.024	-0.078	0.029	>0.999	0.601	-0.011	-0.064	0.042	>0.999	0.683
Depression	-0.021	-0.031	-0.011	<0.001	<0.001	-0.012	-0.022	-0.002	0.184	0.046
+ Antidepressant	-0.058	-0.078	-0.038	<0.001	<0.001	-0.032	-0.052	-0.013	0.010	0.003

Note: β = ordinary least squares regression beta coefficient; CI = confidence interval; Ref = reference group; Bonf. = Bonferroni; BH = Benjamini & Hochberg. Model 1 – adjusted for age and sex; Model 2 – adjusted for age, sex, white blood cell count, Townsend deprivation index, physical activity, smoking status, body mass index, body fat percentage and C-reactive protein. *P*-values corrected for eight tests.