Supplementary Material

Model specification:

$$Y_{ij} = \pi_{0i} + \pi_{1i} time_{ij} + \pi_{2i} time_{ij}^{2} + \varepsilon_{ij}$$
(Eq. 1)

$$\pi_{0i} = \gamma_{00} + \gamma_{01} group_{i} + \gamma_{01} month_{i} + \zeta_{0i}$$
(Eq. 2)

$$\pi_{1i} = \gamma_{10} + \gamma_{11} group_{i} + \zeta_{1i}$$
(Eq. 3)

$$\pi_{2i} = \gamma_{20} + \gamma_{21} group_{i} + \zeta_{2i}$$
(Eq. 4)

The growth curve model is presented mathematically in Equation (1) to (4). Equation (1) addresses intra-individual growth where Y_{ij} , depressive or anxiety symptoms for the individual i at time j, is a function of time and its quadratic term. π_{0i} represents the individual i's initial status. π_{1i} and π_{2i} represent growth rates and ε_{ij} is the residual term. Equation (2) to (4) addresses inter-individual differences in the initial status (intercept) and growth rates. γ_{00} , γ_{10} and γ_{20} represent the population average initial status (intercept) and growth rates. ζ_{0i} , ζ_{1i} and ζ_{2i} are parameter residuals.



Fig S1. Sample selection process





Fig S3. Propensity score density before and after matching



Fig S4. Standardised percentage bias for each covariate before and after matching



Fig S5. Predicted growth trajectories by Covid-19 group with 95% confidence intervals



Fig S6. Propensity score density before and after matching based on one-to-many nearest neighbour matching



Fig S7. Predicted growth trajectories by Covid-19 group with 95% confidence intervals based on one-to-many nearest neighbour matching



Fig S8. Predicted growth trajectories by Covid-19 group with 95% confidence intervals based on kernel matching (*NB these sensitivity analyses yielded less balanced matches so apparent differences in pre-infection mental health should be interpreted with caution*)



Fig S9. Predicted growth trajectories by Covid-19 group with 95% confidence intervals based on one-to-one matching excluding cases based on symptoms



Fig S10. Predicted growth trajectories by Covid-19 group with 95% confidence intervals based on one-to-one matching excluding people who were unsure about their long Covid diagnosis

		Long Covid	Short Covid
		(N=481)	(N=481)
Gender	Men	21.8%	21.0%
	Women	78.2%	79.0%
Ethnicity	White	94.2%	95.0%
	Ethnic minority	5.8%	5.0%
Age group	18-29	6.0%	7.7%
	30-45	31.6%	30.8%
	46-59	39.1%	38.9%
	60+	23.3%	22.7%
Education	GCSE or below	15.4%	16.6%
	A levels or equivalent	19.5%	18.7%
	Degree or above	65.1%	64.7%
Annual income	<£16,000	14.3%	14.8%
	≤£16,000-29,999	23.7%	22.7%
	≤£30,000-59,999	38.3%	41.0%
	≤£60,000-89,999	16.2%	15.0%
	≥£90,000	7.5%	6.7%
Employment status	Employed, keyworker	39.1%	40.1%
	Employed, non-keyworker	32.6%	31.0%
	Not employed	28.3%	28.9%
Area of living	City	32.0%	31.0%
	Town	46.8%	48.2%
	Rural	21.2%	20.8%
Living status	Living alone	14.8%	12.5%
	Living with adult only	48.2%	50.7%
	Living with children	37.0%	36.8%
Social contact	Twice a month or less	35.3%	33.7%
	Once or twice a week	33.3%	33.3%
	Three times a week or more	31.4%	33.1%
Number of close friends		4.7 (3.1)	4.8 (3.1)
Physical health condition	Yes	44.1%	44.1%
	No	55.9%	55.9%
Mental health condition	Yes	23.1%	23.9%
	No	76.9%	76.1%
Baseline depressive symptoms		8.3 (5.9)	8.2 (6.0)
Baseline anxiety symptoms		6.5 (5.4)	6.4 (5.2)
Month had COVID-19		10.5 (6.5)	10.3 (7.1)
Confirmed by testing (COVID-19	Yes	74.6%	72.8%
or antibody test)	No	25.4%	27.2%
Severe symptoms (hospitalised or	Yes	81.7%	82.5%
had to stay in bed)	No	18.3%	17.5%

Table S1 Sample characteristics by matched Covid-19 group

		Long Covid	Short Covid	Total
Test	Yes, a medical professional has formally diagnosed me	112	2	112
	with long Covid	23.3%	6	11.6%
	Yes, I have not been formally diagnosed but consider	369	9	369
	myself to have Long Covid	76.7%	6	38.4%
	No, I do not consider myself to have Long Covid	-	- 369	369
			/6./%	38.4%
	i am unsure	-	- 112	112
	Total	10	25.5% 1 191	11.0%
	lotal	40. 100%	1 401 401	100%
Severity (first 1-2 weeks)	I was hospitalised	3(י ייטט א ר 16	100%
	i was nospitaliseu	50 6 70	/ 2.20/	40
	Lownering and sumptoms and had to rest in had	0.27	0 - 5.5/0	4.0/0
	r experienced symptoms and had to rest in bed	30:	3 381	744
		/5.5%	6 /9.2%	//.3%
	I experienced symptoms but was able to carry on with	79	9 71	150
	daily activities	16.4%	6 14.8%	15.6%
· · · ·	I was asymptomatic	9	9 13	22
		1.9%	6 2.7%	2.3%
	Total	48:	1 481	962
		100%	6 100%	100%
	My symptoms were worse at the beginning (the first 1- 2 weeks) and then got better	23	3 179	202
		4.8%	6 37.6%	21.2%
Symptoms	My symptoms were worse at the beginning (the first 1-	152	2 167	319
	2 weeks) and then mostly got better but some lingered	32.0%	6 35.1%	33.5%
	After the first 1-2 weeks, my symptoms got better but	6	7 21	88
	then the same symptoms kept coming back	14.19	6 4.4%	9.3%
	After the first 1-2 weeks, my symptoms got better but I	60	0 15	75
	then developed new symptoms	12.6%	6 3.2%	7.9%
	Most of my symptoms lasted for 2-3 weeks	2	5 39	64
		5.3%	6 8.2%	6.7%
	Most of my symptoms lasted for 4-12 weeks	62	2 23	85
		13.19	6 4.8%	8.9%
	Most of my symptoms lasted for more than 12 weeks	8	1 6	87
		17 19	- 0 6 13%	9.2%
	I cannot answer this question (e.g. you had COVID yery		5 26	3.270
	recently)	1 10	20 ζ ς ς ο/	2 20/
	Total	/ ۲.۱ ا ر ۸	5.5/0 5.7/0	0.570 0E1
	iUtai	4/:	y 470	1000/
		100%	° 100%	100%

Table S2. Detailed Covid experience measures by matched Covid-19 group