

# EARLY OUTPATIENT TREATMENT OF COVID-19: A RETROSPECTIVE ANALYSIS OF 392 CASES IN ITALY

## Questionnaire for physicians

Name \_\_\_\_\_ date \_\_\_\_\_

Answer please to the following questions having in mind your attitudes and experience during the time interval included in the study (1/11/20-31/3/21).

1. How often did you prescribe **vitamins and supplements** in:

	never	sometimes	often	always
<b>Covid-19 stage 0</b>				
<b>Covid-19 stage 1</b>				
<b>Covid-19 stage 2a</b>				
<b>Covid-19 stage 2b</b>				

*(for each stage, only one choice allowed)*

2. How often did you prescribe **aspirin** in:

	never	sometimes	often	always
<b>Covid-19 stage 0</b>				
<b>Covid-19 stage 1</b>				
<b>Covid-19 stage 2a</b>				
<b>Covid-19 stage 2b</b>				

*(for each stage, only one choice allowed)*

2.1 List please criteria supporting your choice (for example presence/absence of any signs/symptoms, risk factors, indications/contraindications)

3. How often did you prescribe **antibiotics** in:

	never	sometimes	often	always
<b>Covid-19 stage 0</b>				
<b>Covid-19 stage 1</b>				
<b>Covid-19 stage 2a</b>				
<b>Covid-19 stage 2b</b>				

*(for each stage, only one choice allowed)*

3.1 List please criteria supporting your choice (for example presence/absence of any signs/symptoms, risk factors, indications/contraindications)

4. How often did you prescribe **hydroxychloroquine** in:

	<b>never</b>	<b>sometimes</b>	<b>often</b>	<b>always</b>
<b>Covid-19 stage 0</b>				
<b>Covid-19 stage 1</b>				
<b>Covid-19 stage 2a</b>				
<b>Covid-19 stage 2b</b>				

*(for each stage, only one choice allowed)*

4.1 List please criteria supporting your choice (for example presence/absence of any signs/symptoms, risk factors, indications/contraindications)

5. How often did you prescribe **ivermectin** in:

	<b>never</b>	<b>sometimes</b>	<b>often</b>	<b>always</b>
<b>Covid-19 stage 0</b>				
<b>Covid-19 stage 1</b>				
<b>Covid-19 stage 2a</b>				
<b>Covid-19 stage 2b</b>				

*(for each stage, only one choice allowed)*

5.1 List please criteria supporting your choice (for example presence/absence of any signs/symptoms, risk factors, indications/contraindications)

6. How often did you prescribe **colchicine** in:

	<b>never</b>	<b>sometimes</b>	<b>often</b>	<b>always</b>
<b>Covid-19 stage 0</b>				
<b>Covid-19 stage 1</b>				
<b>Covid-19 stage 2a</b>				
<b>Covid-19 stage 2b</b>				

*(for each stage, only one choice allowed)*

6.1 List please criteria supporting your choice (for example presence/absence of any signs/symptoms, risk factors, indications/contraindications)

7. How often did you prescribe **enoxaparin** in:

	<b>never</b>	<b>sometimes</b>	<b>often</b>	<b>always</b>
<b>Covid-19 stage 0</b>				
<b>Covid-19 stage 1</b>				
<b>Covid-19 stage 2a</b>				
<b>Covid-19 stage 2b</b>				

*(for each stage, only one choice allowed)*

7.1 List please criteria supporting your choice (for example presence/absence of any signs/symptoms, risk factors, indications/contraindications)

8. How often did you prescribe **glucocorticoids** in:

	<b>never</b>	<b>sometimes</b>	<b>often</b>	<b>always</b>
<b>Covid-19 stage 0</b>				
<b>Covid-19 stage 1</b>				
<b>Covid-19 stage 2a</b>				
<b>Covid-19 stage 2b</b>				

*(for each stage, only one choice allowed)*

8.1 List please criteria supporting your choice (for example presence/absence of any signs/symptoms, risk factors, indications/contraindications)

9. How often did you observe any suspect adverse reactions with each of the following drugs?

	never	uncommonly	commonly	very commonly
		<1%	1-10%	>10%
<b>vitamins and supplements</b>				
<b>aspirin</b>				
<b>antibiotics</b>				
<b>hydroxychloroquine</b>				
<b>ivermectin</b>				
<b>colchicine</b>				
<b>enoxaparin</b>				
<b>glucocorticoids</b>				

10. In case you observed any suspect adverse reactions, please indicate whether they were:

<b>Grade 1: Mild</b> <i>transient or mild discomfort (&lt; 48 hours); no medical intervention or therapy required</i>	
<b>Grade 2: Moderate</b> <i>mild to moderate limitation in activity; some assistance may be needed; no or minimal medical intervention or therapy required</i>	
<b>Grade 3: Severe</b> <i>marked limitation in activity, some assistance usually required; medical intervention or therapy required, hospitalizations possible</i>	
<b>Grade 4: Life threatening</b> <i>extreme limitation in activity, significant assistance required; significant medical intervention or therapy required, hospitalization or hospice care probable</i>	

11. In case you observed grade 2 or higher suspect adverse reactions, please briefly describe the case(s):