

1 **SARS-CoV-2 reinfections with BA.1 (Omicron) variant among**
2 **fully vaccinated individuals in the northeast of Brazil**

3
4 Francisco P. F. Neto^{1,2,4¶}, Diego G. Teixeira^{1¶}, Dayse C. S. da Cunha¹, Ingryd C. Moraes¹,
5 Celisa P. M. Tavares¹, Genilson P. Gurgel⁵, Sanderson D. do Nascimento⁴, David C. dos
6 Santos³, Alexandre de O. Sales⁴, Selma M.B. Jerônimo^{1*}.

7
8 ¹ Instituto de Medicina Tropical do Rio Grande do Norte, Universidade Federal do Rio Grande
9 do Norte, Natal, Rio Grande do Norte, Brazil.

10 ² Biochemistry Department, Universidade Federal de São Paulo, São Paulo, São Paulo, Brazil.

11 ³ Instituto Metr pole Digital, Universidade Federal do Rio Grande do Norte, Natal, Rio Grande
12 do Norte, Brazil.

13 ⁴ Get lio Sales Diagn sticos, Natal, Rio Grande do Norte, Brazil.

14 ⁵ Secretaria Municipal de Sa de, Apodi, Rio Grande do Norte, Brazil.

15 * Corresponding author:

16 E-mail: selma.jeronimo@ufrn.br (SMBJ)

17

18 ¶ Contributed equally to this work: Francisco P. F. Neto and Diego G. Teixeira

19 **Abstract**

20 **Background**

21 The first case of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) in
22 Rio Grande do Norte, northeast Brazil, was diagnosed on March 12, 2020; thereafter, the
23 pattern of COVID-19 followed the multiple waves as seen elsewhere. Those waves were mostly
24 due to the SARS-CoV-2 virus mutations leading to emergence of variants of concern (VoC).
25 The introduction of new VoCs in a population context of prior SARS-CoV-2 infections or after
26 vaccination has been a challenge in understanding the kinetics of the protective immune
27 response against SARS-CoV-2. The aim of this study was to investigate the outbreak of SARS-
28 CoV-2 reinfections observed in mid-January 2022 in Rio Grande do Norte state, Brazil when
29 the omicron variant was introduced.

30 **Methodology/Principal findings**

31 From a total of 172,965 individuals with mild to severe respiratory symptoms, 58,097
32 tested positive for SARS-CoV-2 between March 2020 through mid-February 2022. Of those
33 previously infected, 444 had documented a second SARS-CoV-2 infection and 9 of these
34 reinfection cases were selected for sequencing. Genomic analysis revealed that virus lineages
35 diverged between primary and the reinfection, with the latter caused by the Omicron (BA.1)
36 variant among individuals fully vaccinated against SARS-CoV-2.

37 **Conclusions/Significance**

38 Once all subjects whose samples were sequenced had prior SARS-CoV-2 infection and
39 were also fully vaccinated, our data suggest that the Omicron variant evades natural and
40 vaccine-induced immunities, confirming the continuous need to decrease transmission and to
41 develop effective blocking vaccines.

42 **Keywords:** COVID-19; SARS-CoV-2; Omicron; Reinfection; Vaccination; Natural infection.

43 **Author summary**

44 The pattern of the COVID-19 pandemic has been characterized by multiple waves of
45 cases with a variety of outcomes from asymptomatic, to moderate or to severe fatal cases. By
46 December 2021, about 75.3% of Rio Grande do Norte population, northeast Brazil, had already
47 been fully vaccinated against SARS-CoV-2 and a decrease in newer detection cases was seen
48 to about 8% of the suspected ones. Nevertheless, with the introduction of the Omicron variant
49 at the end of 2021, the number of new SARS-CoV-2 infections reached its highest peak since
50 the start of the pandemic with 75% of the suspected cases testing positive. From March 2020
51 to February 2022, we confirmed 444 reinfection cases among the ones tested, of which 62.3%
52 (n=277) occurred during the Omicron outbreak, from December 2021 to early February 2022.
53 Of the reinfection cases, 9 were sequenced and genetic analysis showed that they belong to a
54 BA.1 lineage, which seems to have been introduced multiple times into the region. The primary
55 isolates varied. Thus, our data suggest that the Omicron variant evades immunity provided from
56 either natural infection from any other SARS-CoV-2 variants or from different types of
57 vaccines.

58 **Introduction**

59 The world has followed the spread of SARS-CoV-2 since 2019 in real time. However,
60 over two years after the first cases of Coronavirus Disease 2019 (COVID-19) were reported in
61 China, much is still unknown about the mutations and the molecular evolution of SARS-CoV-2
62 over time. The unpredictability of host diversity and the selective pressure by the immune
63 system either by natural infection and/or vaccination on viruses lead us to a field of concerns
64 that gradually have been enlightened.

65 The emergence of SARS-CoV-2 variants with potentially increased transmissibility,
66 virulence, and resistance to antibody neutralization have already been reported [1,2]. The
67 dissemination of these variants, so-called variants of concern (VoC), are also closely associated
68 with higher viral load [3], and/or rates of reinfection [4] among individuals. By the fact that
69 VoCs can surpass the protective effects of humoral immunity induced either by natural
70 infection or by vaccination [5,6], this poses an additional challenge to curb the pandemic since
71 the highest number of new cases and an increased number of reinfections still have been
72 reported worldwide in 2022 [7].

73 The multiwave pandemic dynamics plays different scenarios depending on the
74 mutations present in each SARS-CoV-2 variant and its location. Some of them can provide
75 great impact in specific regions of the globe. For example, studies have shown that the Gamma
76 (P.1) variant is more prevalent in South America [8–10], however, by contrast, the Delta variant
77 and its sublineages [4,11,12] have been found all over the world.

78 Recently, Brazil faced the third wave of COVID-19 pandemic with a record upsurge of
79 cases that started in the end of December 2021 and it had been sustained by the highly
80 transmissible variant B.1.1.529, a VoC, named Omicron [13]. In Brazil, COVID-19 vaccination
81 started on January 17, 2021, however despite the high coverage of immunization in the country,
82 achieving over 79% of the population with at least one dose of the vaccine, at the end of January

83 2022 [14], there was still a sustained number of cases occurring over the year of 2021, and
84 some of the COVID-19 reinfection cases had been reported among fully vaccinated people [6].

85 From March 2020 to February 2022, 444 reinfection episodes were detected in our
86 database, in which 277 cases were diagnosed in the first two months of 2022. Here, we present
87 nine cases of reinfection with the Omicron variant in immunocompetent and fully vaccinated
88 individuals. We also provide genomic variations and clinical features between primary
89 infection and reinfection cases among residents of Rio Grande do Norte (RN) State, located in
90 the Northeastern region of Brazil.

91

92 **Materials and Methods**

93 **Data acquisition, sample collection and clinical testing**

94 All clinical data and nasopharyngeal swab specimens from the first and second
95 infection were obtained from patients in the municipalities of Natal and Apodi, located in the
96 state of Rio Grande do Norte, Brazil. The research was carried out at the Instituto de Medicina
97 Tropical-UFRN (IMT-UFRN) in a partnership with Getúlio Sales Diagnósticos and Health
98 Municipal Secretariats of Apodi and Natal.

99 Nasopharyngeal swabs were collected and placed in Viral Transport Medium.
100 Samples were transported to IMT-UFRN and tested in the same day for SARS-CoV-2 by
101 reverse transcriptase quantitative polymerase chain reaction (RT-qPCR). Remnant samples
102 were stored at -80°C . SARS-CoV-2 infection was diagnosed by RT-qPCR using CDC/EUA
103 protocol [15]. The volunteers filled a questionnaire with epidemiological questions We
104 selected 18 samples with RT-qPCR positive results, presenting a cycle threshold (Ct) value \leq
105 30, collected from July 2020 to February 2022.

106 **RNA extraction, library preparation and sequencing**

107 Viral RNA was isolated from 200- μl samples collected from people suspected of
108 SARS-CoV-2- infections using the QIAamp Viral RNA Mini kit (QIAGEN, Hilden,
109 Germany) according to the manufacturer's instructions. Virus genome sequencing was
110 carried out on all positive samples selected using a COVIDSeq protocol (Illumina Inc, USA).

111 cDNA synthesis was performed from the extracted RNA using a multiplex
112 polymerase chain reaction (PCR) protocol, producing 98 amplicons across the SARS-CoV-2
113 genome (<https://artic.network/>). The primer pool additionally had primers targeting human

114 RNA, producing an additional 11 amplicons. The PCR amplified product was later processed
115 for tagmentation and adapter ligation using IDT for Illumina Indexes. Further enrichment and
116 cleanup were performed as per protocols provided by the manufacturer (Illumina Inc). All
117 samples were processed as batches in a 16-well plate. Pooled samples were quantified using
118 Qubit 2.0 fluorometer (Invitrogen Inc.) and fragment sizes were analyzed in 4150 TapeStation
119 System (Agilent Inc). The pooled library was further normalized to 4nM concentration and
120 25 µl of each normalized pool containing index adapter set A were combined in a new
121 microcentrifuge tube to a final concentration of 100pM. The pooled libraries were loaded and
122 clustered onto iSeq 100 i1 Reagent v2 (Illumina) with flow cell and sequenced using
123 sequencing by synthesis (SBS) chemistry on the Illumina iSeq 100 sequencing system. The
124 sequencing run was created with Local Run Manager software installed on the iSeq 100
125 instrument. Using the Generate FASTQ Analysis module with the custom library preparation
126 kit options, the number of sequencing cycles was set to 151 bp and the paired-end sequencing
127 option was determined.

128 **Read mapping and Consensus calling**

129 All reads were mapped to the reference SARS-CoV-2 genome (GenBank accession
130 number MN908947.3) using the BWA-MEM V0.7.12-r1039 [16]. Next, the removal of
131 duplicate reads and the split of reads containing poorly sequenced base was proceeded by the
132 Genomic Analysis-Toolkit (GATK) V4.2.0.0 [17]; the quality of the final mapping process
133 was accessed by Mosdepth [18] and Qualimap V2.2.1 [19]. The SNV identification was
134 performed by the *HaplotypeCaller* function from the GATK, followed by the filtration using
135 the *VariantFiltration*, *FilterVcf* and *SelectVariants*, excluding variants with depth lower than
136 10. Those variants which passed by the depth threshold were annotated using the SnpEff
137 V5.0e [20].

138 The sequence consensus calling was performed by the
139 *FastaAlternateReferenceMaker* function from GATK, and for the final fasta file those
140 positions covered by fewer than 10 reads were replaced by Ns, by using an in-house script
141 written in Python 3.8 (available at GitHub). All sequences were submitted to the Pangolin
142 COVID-19 Lineage Assigner v.3.1.17 [21] (accessed on 24/Jan/2022). The whole genome
143 sequences of viral variants of the nine samples (two samples from each case of reinfection)
144 were deposited in the GISAID database (accession nos. EPI_ISL_9467461,
145 EPI_ISL_9467460, EPI_ISL_9467471, EPI_ISL_9467463, EPI_ISL_9467462,
146 EPI_ISL_9467470, EPI_ISL_9467458, EPI_ISL_9467469, EPI_ISL_9467457,
147 EPI_ISL_9467468, EPI_ISL_9467459, EPI_ISL_9467454, EPI_ISL_9467465,
148 EPI_ISL_9467464, EPI_ISL_9467456, EPI_ISL_9467467, EPI_ISL_9467455,
149 EPI_ISL_9467466) and all GISAID sequences used in this study are listed in S1 Table.

150 **Phylogenetics analysis**

151 The 18 genome sequences obtained in this work were aligned to 700 other genome sequences
152 from Rio Grande do Norte, Brazil, obtained from GISAID on January 20, 2022, using the
153 MAFFT v.7.394 [22]. We then built a maximum likelihood (ML) phylogenetics tree supported
154 by 1,000 replicates of bootstrap using IQ-TREE2 [23], under the substitution model
155 GTR+F+R4, which was inferred to be the best-fitting model by de software itself. The same
156 procedure was performed for the Alignment containing only the Brazilian BA.1 genomes,
157 under the GTR+F+R3 model. Tree topologies were inspected by Figtree v.1.4.4
158 (<http://tree.bio.ed.ac.uk/software/figtree/> last accessed 2021-11-29).

159 **Ethical Issues**

160 The study protocol was reviewed and approved by the Federal University of Rio
161 Grande do Norte Ethical committee (36287120.2.0000.5537).

163 **Results**

164

165 **Tracking reinfection cases in Rio Grande do Norte, Brazil**

166 From March 2020 to February 2022, a total of 172,965 individuals with upper
167 respiratory symptoms underwent testing by RT-qPCR for SARS-CoV-2 at IMT-RN, of which
168 58,097 tested positive. Positive tests remained high from December 2020 to July 2021 (mean
169 160.9/day, 100.4 SD), when thereafter a decrease of COVID-19 cases was observed until the
170 beginning of January 2022 (mean 27.8/day, 28.5 SD), followed by a rapid increase in positivity
171 (Fig 1A). Among those, 444 individuals had 2 episodes of symptomatic infection for COVID-
172 19 (Fig 1B). In mid-2020, reinfection cases were already reported, although scarce, and in early
173 2022, during the Omicron-predominant period, they reached a peak comprising 62.3% (n=277)
174 of all reinfection cases identified in our system. The earliest day of reinfection was 62 days
175 apart, after the primary infection, and the longest was 638.

176 **Fig 1. Timeline of COVID-19 detection cases in Rio Grande do Norte state, Brazil.** (A)
177 Positive COVID-19 cases in RN detected by RT-qPCR in IMT-RN laboratory. (B) Detection
178 of SARS-CoV-2 infection in 444 individuals. Participants with two entries in our database were
179 selected and the period of the first (gray circle) and second (red square) entry was plotted. A
180 gray line is connecting both COVID-19 tests of the same individual. The density of COVID-
181 19 cases on top of the plot. (C) SARS-CoV-2 PANGO lineages detected in RN from March
182 2020 to January 2022, obtained from GISAID.

183 **SARS-CoV-2 isolates sequencing**

184 Of the 444 individuals with documented SARS-CoV-2 reinfection, 9 immunocompetent
185 and fully vaccinated individuals were chosen to have their isolates sequenced (S1 Fig). Among
186 those, 5 were males and the mean age was 32.4 years (± 7.4). The first infection ranged from
187 June 2020 to November 2021. All reinfection cases occurred in January 2022. The time course

188 between the 2 episodes of COVID-19 ranged between 70 and 584 days, with mild clinical
189 presentation for both infections (Fig 2A). The average between the last vaccination date and
190 the second episode was 86.8 days (± 46.1). The most reported symptoms during both infections
191 were headache, nasal congestion, fatigue, rhinorrhea, myalgia, pharyngitis, fever, and cough.
192 Comorbidities, such as hypertension, diabetes mellitus, cancer, and asthma/COPD were not
193 reported among participants and none of them were hospitalized. In addition, two of the
194 subjects had already received a third dose of the vaccine.

195 In order to distinguish between reinfection and long-term viral persistence, we
196 performed a maximum-likelihood (ML) phylogenetic analysis of all SARS-CoV-2 whole
197 genomes from Rio Grande do Norte state, recovered from GISAID on January 20, 2021,
198 combining with all 18 samples sequenced in this work (Fig 2B). The earliest isolate reported
199 from Rio Grande do Norte state was obtained on July 02, 2020 and belongs to the BA.1.1.311
200 lineage (EPI_ISL_2466179), but herein we sequenced an oldest one, from June 08, 2020; which
201 was assigned to the B.1.195 lineage by the Pangolin tool, when the ML tree root was set. The
202 tree topology revealed that all sequences belonged to the variant of concern Omicron (BA.1)
203 lineage and they were clustered in a monophyletic clade while their respective pairs, from the
204 previous SARS-CoV-2 infection, were found spread through the tree. A sequence from the
205 individual #2 isolated during the first infection which was assigned to the N.1 lineage, was
206 placed close to the Omicron branch. The N.1 lineage had been detected in Brazil during the
207 early epidemic in 2020, according to the *outbreak.info* website.

208 **Fig 2. Identification of SARS-CoV-2 reinfection cases in Rio Grande do Norte, Brazil.**

209 **(A)** Primary infection and reinfection and the virus lineages. The figure also shows the time
210 of the last COVID-19 vaccine. **(B)** Maximum-likelihood tree of all 709 SARS-CoV-2 whole-
211 genome sequences available at GISAID repository, downloaded on January 20, 2022. The
212 color assigned to the branches are related to their respective lineage and sequenced samples

213 in this work are labeled according to their pair, sampling date and virus lineage, respectively.
214 The dark dots at the dichotomies represent those nodes with high bootstrap support (≥ 80).
215 The tree's root was set on the youngest isolate (from June 8, 2020), sequenced in this work,
216 which belongs to the B.1.195 lineage. (C) ML tree composed by other 1700 Omicron
217 sequences isolated in Brazil plus a B.1.1.529 reference as outgroup (EPI_ISL_6704867), with
218 the branches colored according to the Brazilian state where the virus was isolated. The red
219 dots indicate where each of the 9 sequences are placed in the tree. Each of the 9 isolates and
220 the other sequences from Brazil close to them can be seen in S2 Fig.

221 Among the 9 individuals in this study, a female individual had a reinfection episode 70
222 days apart from primary infection and 8 days after the COVID-19 vaccine booster shot. Despite
223 this short interval, the virus lineages diverged between the two infections (Figs 2A and B), the
224 primary occurring through a AY.99.2 variant, while the reinfection episode was by BA.1. It is
225 important to note the high frequency of AY lineages (Delta VOC) on the topology ML tree,
226 representing more than 30% of all sequences on Rio Grande do Norte state, and also that
227 variants arising from Delta were found on almost 100% of sequenced isolates in Brazil in
228 December 2021 [24].

229 To better characterize the Omicron variant detected during the reinfection cases, we
230 aligned it against all other Brazilian BA.1 whole genomes available in the GISAID database
231 until 20 January 2022. In addition, we also selected an African B.1.1.529 reference sequence
232 as an outgroup (accession EPI_ISL_6704867). The second ML tree (Fig 2C) revealed that in a
233 broader scale of this single lineage, the genomes isolated from the reinfection are spread
234 through the tree, emphasizing the multiple introductions of the BA.1 lineage into the Rio
235 Grande do Norte state. The relationship of the 9 BA.1 isolates sequenced here can be observed
236 closer on S2 Fig, with a remarkable presence of isolates from the states of São Paulo,
237 Pernambuco and Ceará.

238 **Discussion**

239 SARS-CoV-2 infection episodes by emerging variants in previously naturally infected
240 or vaccinated individuals are already a real phenomenon worldwide which poses an important
241 issue to understand human immune response to the virus and vaccines [7]. Reinfection cases
242 have been rising profoundly since the Omicron variant of SARS-CoV-2 started to disseminate
243 in November 2021 [25]. To date, such a fact was not observed with the same rate in previously
244 reported variants of concern of SARS-CoV-2. In our study, we demonstrate that 62.3%
245 (277/444) of all reinfection cases detected over 22 months occurred during the Omicron-wave
246 (Fig 1). To confirm those reinfection events, we proposed to investigate a small prospective
247 cohort study from 9 individuals with reinfection of COVID-19 in the state of Rio Grande do
248 Norte, Brazil. This state receives a high number of tourists during the end of the year, and this
249 may have been one of the reasons for the diversity of introductions in the region. From early-
250 December 2021 to mid-January 2022, São Paulo was the Brazilian state with the most
251 expressive number of flights to Natal (S3 Fig), which may explain the similarities between the
252 majority of the isolates sequenced in this work and the ones sequenced in São Paulo. This,
253 together with the fact that São Paulo is one of the states most affected by the omicron variant
254 [26], due to being a hub for intercontinental flights, provides, therefore, an increased risk of
255 arrival of new variants as the omicron case shows.

256 The Brazilian Health Ministry criteria for a SARS-CoV-2 reinfection case is two
257 positive RT-qPCR confirmations within at least 90 days apart, clinical manifestations
258 characteristics to COVID-19, positive viral culture, with an additional viral RNA sequencing
259 from both episodes showing different strains [6],[27]. Our work lacks the viral culture,
260 nevertheless, we demonstrated one case of an individual with two infection episodes, 70 days
261 apart, and the sequence of the two isolates showed clearly different lineages. Reinfection cases
262 under 90 days apart from the episodes have already been described [28].

263 Recent studies have demonstrated that Omicron-infected cases tended to cause mild
264 symptoms such as pharyngitis, headache, nasal [29] which corroborates with our data. Despite
265 the fact that three subjects have a booster, with the third/additional dose, they developed
266 symptomatic COVID-19, as others with complete vaccination and previous history of COVID-
267 19 [7]. These data support the findings that neither the current vaccines available nor previous
268 history of COVID-19 are sufficient to establish a robust response toward the Omicron variant
269 and to promote a reduced risk of infection. Nonetheless, a recent study has shown that booster
270 vaccination increases the neutralization titers to Omicron [5].

271 Although with the data presented herein, we have barely scratched the surface of the
272 many possibilities in which the Omicron lineage or other variants are capable of causing
273 reinfections, further studies are needed to understand the role of SARS-CoV-2 mutations which
274 lead the evasion of the immune response generated by prior infection or vaccination. The
275 kinetics of the protective immunity is quite not ascertained. The currently available vaccines
276 have apparently reduced severity of the COVID-19, even though an increase in hospitalization
277 and deaths occurred with the introduction of the apparent more benign omicron variant, when
278 compared to the Gamma one. This indicates the need to develop more effective vaccines, with
279 potential for reducing transmission and a lengthier protective immune response. Genomic
280 surveillance is a key for assessing new variants and therefore suitable for planning new
281 strategies to battle COVID-19.

282 Our data suggest that the Omicron variant evades immunity provided from different
283 types of vaccines or from natural infection from any other earlier SARS-CoV-2 variants of
284 concern and may be related mostly to the mild symptoms of the upper respiratory tract for the
285 majority of the subjects (S1 Fig). Thus, along with vaccination, the reassurance of protective
286 measures needs to be held to prevent further spread of SARS-CoV-2 variants or risk of
287 appearance of new variants.

288 **Acknowledgements:** We would like to thank the Núcleo de Processamento de Alto
289 Desempenho da Universidade Federal do Rio Grande do Norte - NPAD/UFRN for allowing us
290 to access their computer facilities.

291 **References**

- 292 1. Wang P, Nair MS, Liu L, Iketani S, Luo Y, Guo Y, et al. Antibody resistance of SARS-
293 CoV-2 variants B.1.351 and B.1.1.7. *Nature*. 2021;593: 130–135. doi:10.1038/s41586-
294 021-03398-2
- 295 2. Garcia-Beltran WF, Lam EC, St Denis K, Nitido AD, Garcia ZH, Hauser BM, et al.
296 Multiple SARS-CoV-2 variants escape neutralization by vaccine-induced humoral
297 immunity. *Cell*. 2021;184: 2372-2383.e9. doi:10.1016/j.cell.2021.03.013
- 298 3. Lyngse FP, Mølbak K, Skov RL, Christiansen LE, Mortensen LH, Albertsen M, et al.
299 Increased transmissibility of SARS-CoV-2 lineage B.1.1.7 by age and viral load. *Nat*
300 *Commun*. 2021;12: 7251. doi:10.1038/s41467-021-27202-x
- 301 4. Dhar MS, Marwal R, Vs R, Ponnusamy K, Jolly B, Bhoyar RC, et al. Genomic
302 characterization and epidemiology of an emerging SARS-CoV-2 variant in Delhi, India.
303 *Science*. 2021;374: 995–999. doi:10.1126/science.abj9932
- 304 5. Dejnirattisai W, Huo J, Zhou D, Zahradník J, Supasa P, Liu C, et al. SARS-CoV-2
305 Omicron-B.1.1.529 leads to widespread escape from neutralizing antibody responses.
306 *Cell*. 2022;185: 467-484.e15. doi:10.1016/j.cell.2021.12.046
- 307 6. Resende PC, Bezerra JF, Teixeira Vasconcelos RH, Arantes I, Appolinario L, Mendonça
308 AC, et al. Severe Acute Respiratory Syndrome Coronavirus 2 P.2 Lineage Associated
309 with Reinfection Case, Brazil, June-October 2020. *Emerging Infect Dis*. 2021;27: 1789–
310 1794. doi:10.3201/eid2707.210401
- 311 7. Rahman S, Rahman MM, Miah M, Begum MN, Sarmin M, Mahfuz M, et al. COVID-19
312 reinfections among naturally infected and vaccinated individuals. *Sci Rep*. 2022;12:
313 1438. doi:10.1038/s41598-022-05325-5

- 314 8. Stefanelli P, Trentini F, Guzzetta G, Marziano V, Mammone A, Sane Schepisi M, et al.
315 Co-circulation of SARS-CoV-2 Alpha and Gamma variants in Italy, February and March
316 2021. *Euro Surveill.* 2022;27. doi:10.2807/1560-7917.ES.2022.27.5.2100429
- 317 9. Jaspe RC, Loureiro CL, Sulbaran Y, Moros ZC, D'Angelo P, Rodríguez L, et al.
318 Introduction and rapid dissemination of SARS-CoV-2 Gamma Variant of Concern in
319 Venezuela. *Infect Genet Evol.* 2021;96: 105147. doi:10.1016/j.meegid.2021.105147
- 320 10. Faria NR, Mellan TA, Whittaker C, Claro IM, Candido D da S, Mishra S, et al. Genomics
321 and epidemiology of the P.1 SARS-CoV-2 lineage in Manaus, Brazil. *Science.* 2021;372:
322 815–821. doi:10.1126/science.abh2644
- 323 11. Elliott P, Haw D, Wang H, Eales O, Walters CE, Ainslie KEC, et al. Exponential growth,
324 high prevalence of SARS-CoV-2, and vaccine effectiveness associated with the Delta
325 variant. *Science.* 2021;374: eabl9551. doi:10.1126/science.abl9551
- 326 12. Naveca FG, Nascimento V, Souza V, Corado A de L, Nascimento F, Resende PC, et al.
327 The SARS-CoV-2 variant Delta displaced the variants Gamma and Gamma plus in
328 Amazonas, Brazil. 1 Oct 2021 [cited 11 Mar 2022]. Available:
329 [https://virological.org/t/the-sars-cov-2-variant-delta-displaced-the-variants-gamma-and-](https://virological.org/t/the-sars-cov-2-variant-delta-displaced-the-variants-gamma-and-gamma-plus-in-amazonas-brazil/765)
330 [gamma-plus-in-amazonas-brazil/765](https://virological.org/t/the-sars-cov-2-variant-delta-displaced-the-variants-gamma-and-gamma-plus-in-amazonas-brazil/765)
- 331 13. World Health Organization. Classification of Omicron (B.1.1.529): SARS-CoV-2
332 Variant of Concern. 26 Nov 2021 [cited 28 Jan 2022]. Available:
333 [https://www.who.int/news/item/26-11-2021-classification-of-omicron-\(b.1.1.529\)-sars-](https://www.who.int/news/item/26-11-2021-classification-of-omicron-(b.1.1.529)-sars-cov-2-variant-of-concern)
334 [cov-2-variant-of-concern](https://www.who.int/news/item/26-11-2021-classification-of-omicron-(b.1.1.529)-sars-cov-2-variant-of-concern)
- 335 14. Mathieu E, Ritchie H, Ortiz-Ospina E, Roser M, Hasell J, Appel C, et al. A global
336 database of COVID-19 vaccinations. *Nat Hum Behav.* 2021;5: 947–953.
337 doi:10.1038/s41562-021-01122-8

- 338 15. Centers for Disease Control and Prevention. CDC Diagnostic Tests for COVID-19. 7
339 Aug 2021 [cited 29 Sep 2021]. Available: [https://www.cdc.gov/coronavirus/2019-](https://www.cdc.gov/coronavirus/2019-ncov/lab/testing.html)
340 [ncov/lab/testing.html](https://www.cdc.gov/coronavirus/2019-ncov/lab/testing.html)
- 341 16. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.
342 *Bioinformatics*. 2009;25: 1754–1760. doi:10.1093/bioinformatics/btp324
- 343 17. Van der Auwera GA, Carneiro MO, Hartl C, Poplin R, Del Angel G, Levy-Moonshine
344 A, et al. From FastQ data to high confidence variant calls: the Genome Analysis Toolkit
345 best practices pipeline. *Curr Protoc Bioinformatics*. 2013;11: 11.10.1-11.10.33.
346 doi:10.1002/0471250953.bi1110s43
- 347 18. Pedersen BS, Quinlan AR. Mosdepth: quick coverage calculation for genomes and
348 exomes. *Bioinformatics*. 2018;34: 867–868. doi:10.1093/bioinformatics/btx699
- 349 19. Okonechnikov K, Conesa A, García-Alcalde F. Qualimap 2: advanced multi-sample
350 quality control for high-throughput sequencing data. *Bioinformatics*. 2016;32: 292–294.
351 doi:10.1093/bioinformatics/btv566
- 352 20. Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, et al. A program for
353 annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs
354 in the genome of *Drosophila melanogaster* strain w1118; iso-2; iso-3. *Fly (Austin)*.
355 2012;6: 80–92. doi:10.4161/fly.19695
- 356 21. O’Toole Á, Scher E, Underwood A, Jackson B, Hill V, McCrone JT, et al. Assignment
357 of epidemiological lineages in an emerging pandemic using the pangolin tool. *Virus Evol*.
358 2021;7: veab064. doi:10.1093/ve/veab064
- 359 22. Nakamura T, Yamada KD, Tomii K, Katoh K. Parallelization of MAFFT for large-scale
360 multiple sequence alignments. *Bioinformatics*. 2018;34: 2490–2492.
361 doi:10.1093/bioinformatics/bty121

- 362 23. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective
363 stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol.*
364 2015;32: 268–274. doi:10.1093/molbev/msu300
- 365 24. Gularte JS, da Silva MS, Mosena ACS, Demoliner M, Hansen AW, Filippi M, et al.
366 EARLY INTRODUCTION, DISPERSAL AND EVOLUTION OF DELTA SARS-
367 CoV-2 IN SOUTHERN BRAZIL, LATE PREDOMINANCE OF AY.99.2 AND AY.101
368 RELATED LINEAGES. *Virus Res.* 2022; 198702. doi:10.1016/j.virusres.2022.198702
- 369 25. Mallapaty S. COVID reinfections surge during Omicron onslaught. *Nature.* 2022.
370 doi:10.1038/d41586-022-00438-3
- 371 26. Estado de São Paulo - Instituto Butantan. BOLETIM EPIDEMIOLÓGICO DA REDE
372 DEALERTA DAS VARIANTES DO SARS-COV-2ESTADO DE SÃO PAULO -
373 INSTITUTO BUTANTAN. 1 Jan 2022 [cited 19 Jan 2022]. Available:
374 [https://butantan.gov.br/assets/arquivos/1_1_22_Relat%C3%B3rio_Sequenciamento.pdf](https://butantan.gov.br/assets/arquivos/1_1_22_Relat%C3%B3rio_Sequenciamento.pdf#page=2)
375 #page=2
- 376 27. Yahav D, Yelin D, Eckerle I, Eberhardt CS, Wang J, Cao B, et al. Definitions for
377 coronavirus disease 2019 reinfection, relapse and PCR re-positivity. *Clin Microbiol*
378 *Infect.* 2021;27: 315–318. doi:10.1016/j.cmi.2020.11.028
- 379 28. Tang CY, Wang Y, McElroy JA, Li T, Hammer R, Ritter D, et al. Reinfection with two
380 genetically distinct SARS-CoV-2 viruses within 19 days. *J Med Virol.* 2021;93: 5700–
381 5703. doi:10.1002/jmv.27154
- 382 29. Al-Tawfiq JA, Hoang V-T, Le Bui N, Chu D-T, Memish ZA. The Emergence of the
383 Omicron (B.1.1.529) SARS-CoV-2 Variant: What is the Impact on the Continued
384 Pandemic? *J Epidemiol Glob Health.* 2022. doi:10.1007/s44197-022-00032-w

385

386 **Supporting Information Captions**

387 **S1 Table.** GISAID sequences used in this study.


388 **S1 Fig.** Schematic representation of different types of vaccine against SARS-CoV-2 in all
389 reinfection cases.

390 **S2 Fig.** Subset of the Omicron (BA.1) ML phylogenetic tree indicating the closest isolates from
391 different areas of Brazil. The Omicron genomes sequenced here are indicated with the red tips.


392 **S3 Fig.** Cumulative sum of flights according to the Brazilian federative unit or Country of
393 origin. From December 1, 2021 to December 31, 2021 the data was obtained from National
394 Civil Aviation Agency (ANAC) - Consultation of past flights
395 (<https://sas.anac.gov.br/sas/bav/view/frmConsultaVRA>, accessed on January 24, 2022). Data
396 from January 1 2022 to January 20, 2022 the data was obtained from ANAC - Consultation of
397 planned flights
398 ([https://siros.anac.gov.br/SIROS/\(S\(4gkz4ekn0yp1rg5lv3gvrq41\)\)/view/registro/frmConsulta](https://siros.anac.gov.br/SIROS/(S(4gkz4ekn0yp1rg5lv3gvrq41))/view/registro/frmConsultaVoos)
399 [Voos](#), January 24, 2022). The order from the Brazilian federative unit with the most flight to
400 Natal (from December 01, 2021 to January 20, 2022) from, to the lesser one, is: São Paulo (604
401 flights); Pernambuco (227 flights); Brasília (199 flights); Minas Gerais (85 flights); Rio de
402 Janeiro (78 flights); Bahia (53 flights); Ceará (52 flights); Rio Grande do Norte (20 flights);
403 Paraná (4 flights); Goiás (1 flight); and also Portugal with 22 international flights.

COVID-19 detected by IMT-RN

A



B



C




Figure 1






Figure 2

SUPPLEMENTARY RESULTS




Supplementary Figure S1. Schematic representation of different types of vaccine against SARS-CoV-2 in all reinfection cases.



Supplementary Figure S2. Subset of the Omicron (BA.1) ML phylogenetic tree

indicating the closest isolates from different areas of Brazil. The Omicron genomes sequenced here are indicated with the red tips.



Supplementary Figure S3. Cumulative sum of flights according to the Brazilian federative unit or Country of origin. From December 1, 2021 to December 31, 2021 the data was obtained from National Civil Aviation Agency (ANAC) - Consultation of past flights (<https://sas.anac.gov.br/sas/bav/view/frmConsultaVRA>, accessed on January 24, 2022). Data from January 1 2022 to January 20, 2022 the data was obtained from ANAC - Consultation of planned flights ([https://siros.anac.gov.br/SIROS/\(S\(4gkz4ekn0yp1rg5lv3gvrq41\)\)/view/registro/frmConsultaVoos](https://siros.anac.gov.br/SIROS/(S(4gkz4ekn0yp1rg5lv3gvrq41))/view/registro/frmConsultaVoos), January 24, 2022). The order from the Brazilian federative unit with the most flight to Natal (from December 01, 2021 to January 20, 2022) from, to the lesser one, is: São Paulo (604 flights); Pernambuco (227 flights); Brasília (199 flights); Minas Gerais (85 flights); Rio de Janeiro (78 flights); Bahia (53 flights);

Ceará (52 flights); Rio Grande do Norte (20 flights); Paraná (4 flights); Goiás (1 flight); and also Portugal with 22 international flights.