

Supplementary Material:

		Hazard ratios				
Group	Clinical Endpoint	p	HR _{80/20}	HR _{20/50}	HR _{80/50}	HR _{95/50}
All (n=590,750)	Fatal Prostate Cancer	<10 ⁻¹⁶	4.42 [3.91-5.02]	0.48 [0.45-0.51]	2.12 [2.0-2.27]	3.0 [2.75-3.3]
Non-Hispanic White (n=420,473)		<10 ⁻¹⁶	4.37 [3.77-5.05]	0.48 [0.45-0.52]	2.11 [1.96-2.27]	3.0 [2.69-3.34]
Black or African American (n=102,203)		<10 ⁻¹⁶	2.37 [1.73-3.29]	0.66 [0.56-0.77]	1.57 [1.33-1.85]	1.9 [1.5-2.42]
Hispanic White (n=27,651)		0.001	3.52 [1.72-6.95]	0.53 [0.38-0.76]	1.87 [1.31-2.64]	2.54 [1.49-4.23]
Asian (n=6,644)		0.03	20.33 [2.4-503.26]	0.21 [0.04-0.64]	4.37 [1.52-21.31]	9.51 [1.89-96.34]
Native American (n=5,835)		0.008	10.94 [1.58-85.77]	0.31 [0.11-0.8]	3.41 [1.26-9.62]	6.03 [1.41-26.97]
Pacific Islander (n=3,246)		0.43	3.06 [0.31-175.49]	0.57 [0.08-1.8]	1.75 [0.55-13.44]	2.32 [0.4-51.41]
Unknown (n=16,472)		0.09	2.05 [0.9-4.54]	0.7 [0.48-1.05]	1.44 [0.95-2.16]	1.71 [0.93-3.07]
Other (n=8,226)		0.002	7.66 [2.05-33.35]	0.36 [0.17-0.7]	2.78 [1.43-5.88]	4.58 [1.71-13.8]
All (n=590,750)		Metastatic Prostate Cancer	<10 ⁻¹⁶	4.89 [4.57-5.21]	0.46 [0.44-0.47]	2.23 [2.16-2.31]
Non-Hispanic White (n=420,473)	<10 ⁻¹⁶		4.62 [4.27-5.01]	0.47 [0.45-0.49]	2.17 [2.08-2.26]	3.13 [2.95-3.32]
Black or African American (n=102,203)	<10 ⁻¹⁶		3.03 [2.62-3.51]	0.59 [0.55-0.63]	1.78 [1.65-1.92]	2.28 [2.05-2.54]
Hispanic White (n=27,651)	<10 ⁻⁷		2.92 [1.96-4.43]	0.58 [0.47-0.71]	1.71 [1.4-2.11]	2.22 [1.65-3.01]
Asian (n=6,644)	<10 ⁻⁴		7.61 [2.64-19.07]	0.35 [0.22-0.61]	2.7 [1.61-4.22]	4.56 [2.07-9.22]
Native American (n=5,835)	<10 ⁻⁴		4.98 [2.16-12.06]	0.46 [0.3-0.69]	2.28 [1.48-3.57]	3.34 [1.78-6.38]
Pacific Islander (n=3,246)	0.4		1.77 [0.55-6.43]	0.75 [0.39-1.35]	1.33 [0.74-2.55]	1.53 [0.64-4.07]

Unknown (n=16,472)	Prostate Cancer	<10 ⁻⁹	4.74 [2.97-7.47]	0.46 [0.37-0.58]	2.2 [1.74-2.78]	3.19 [2.25-4.51]
Other (n=8,226)		<10 ⁻⁶	5.95 [3.02-11.48]	0.41 [0.3-0.58]	2.45 [1.74-3.43]	3.8 [2.27-6.26]
All (n=590,750)		<10 ⁻¹⁶	5.2 [5.09-5.31]	0.44 [0.44-0.45]	2.31 [2.28-2.33]	3.39 [3.33-3.44]
Non-Hispanic White (n=420,473)		<10 ⁻¹⁶	5.12 [4.99-5.26]	0.45 [0.44-0.45]	2.29 [2.26-2.32]	3.38 [3.31-3.44]
Black or African American (n=102,203)		<10 ⁻¹⁶	3.22 [3.06-3.39]	0.57 [0.56-0.58]	1.83 [1.79-1.88]	2.39 [2.3-2.48]
Hispanic White (n=27,651)		<10 ⁻¹⁶	4.24 [3.74-4.85]	0.49 [0.45-0.52]	2.06 [1.93-2.2]	2.92 [2.66-3.22]
Asian (n=6,644)		<10 ⁻¹⁶	5.32 [4.04-7.09]	0.43 [0.37-0.49]	2.27 [1.98-2.61]	3.49 [2.83-4.34]
Native American (n=5,835)		<10 ⁻¹⁶	5.1 [4.04-6.55]	0.45 [0.4-0.51]	2.31 [2.05-2.62]	3.4 [2.85-4.12]
Pacific Islander (n=3,246)		<10 ⁻¹⁰	3.58 [2.5-5.37]	0.53 [0.43-0.63]	1.9 [1.57-2.33]	2.6 [1.98-3.57]
Unknown (n=16,472)		<10 ⁻¹⁶	5.27 [4.39-6.3]	0.44 [0.4-0.48]	2.32 [2.12-2.55]	3.45 [3.0-3.94]
Other (n=8,226)		<10 ⁻¹⁶	5.52 [4.45-6.89]	0.43 [0.39-0.48]	2.36 [2.12-2.64]	3.59 [3.06-4.25]

Supplementary Table 1: Association of PHS290 with any, metastatic and fatal prostate cancer with self-reported ethnicity/race groups. Cox Proportional-Hazards results from association with age at prostate cancer, metastatic prostate cancer and death from prostate cancer. *P*-values reported are from univariable models using PHS290 as the sole predictor variable. Hazard ratios (HRs) are shown comparing men in various percentiles of genetic risk. HR_{80/20}: highest 20% (≥80th percentile of PHS290, using previously published thresholds for men <70 years old and no diagnosis of cancer) vs. average risk (30-70th percentile). HR_{20/50}: lowest 20% (≤20th percentile) vs. average risk. HR_{80/50}: highest 20% vs. average risk. HR_{95/50}: highest 5% (≥95th percentile) vs. average risk. Numbers in brackets are 95% confidence intervals.

	All	European	African	Admixed American	Middle Eastern	Central/South Asian	East Asian
<i>Participants</i>							
All participants	586,608 (375,763)	420,672 (297,152)	104,997 (48,308)	50,586 (24,961)	694 (437)	567 (225)	9,092 (4,680)
Prostate cancer	68,706 (49,073)	48,178 (37,602)	16,178 (8,717)	3,775 (2,358)	75 (59)	13 (8)	487 (329)
Metastases from prostate cancer	6,373 (4,250)	4,312 (3,181)	1,650 (822)	361 (215)	5 (3)	2 (0)	43 (29)
Death from prostate cancer	1,843 (1,302)	1,364 (1,051)	377 (196)	90 (46)	4 (4)	0 (0)	8 (5)
<i>Age demographics</i>							
Age at diagnosis, median and IQR	67 [62-72]	68 [63-73]	63 [58-68]	65 [60-70]	73 [67.5-78]	67 [62-69]	67 [61-72]
Age at last follow-up, median and IQR	664 [59-74]	68 [62-75]	62 [55-70]	59 [47-71]	66 [49.5-81]	49 [37-60]	56 [42-69]

Supplementary Table 2 Participant characteristics for genetic ancestry groups, n=586,608. Genetic ancestry groups were made based on PCA analysis on individual genotypes from Wendt et al. 4142 individuals were not assigned a genetic ancestry group and are excluded here.

Group	Clinical Endpoint	p	Hazard ratios			
			HR80/20	HR20/50	HR80/50	HR95/50
All (n=586,608)	Fatal Prostate Cancer	<10-16	4.39 [3.84-5.01]	0.48 [0.45-0.52]	2.12 [1.98-2.26]	2.99 [2.71-3.3]
European (n=420,672)		<10-16	4.3 [3.7-4.99]	0.49 [0.45-0.52]	2.09 [1.94-2.26]	2.97 [2.65-3.31]
African (n=104,997)		<10-7	2.35 [1.71-3.22]	0.66 [0.57-0.77]	1.56 [1.32-1.84]	1.89 [1.49-2.39]
Admixed American (50,586)		<10-5	4.3 [2.37-7.71]	0.48 [0.36-0.65]	2.08 [1.54-2.78]	2.96 [1.9-4.57]
East Asian (n=9,092)		-	-	-	-	-
Central/South Asian (n=567)		-	-	-	-	-
Middle Eastern (n=694)		-	-	-	-	-
All (n=586,608)		Metastatic Prostate Cancer	<10-16	4.87 [4.56-5.18]	0.46 [0.44-0.47]	2.23 [2.16-2.3]
European (n=420,672)	<10-16		4.6 [4.26-4.98]	0.47 [0.45-0.49]	2.16 [2.08-2.25]	3.12 [2.95-3.31]
African (n=104,997)	<10-16		2.96 [2.58-3.44]	0.59 [0.55-0.63]	1.76 [1.64-1.9]	2.25 [2.03-2.51]
Admixed American (50,586)	<10-16		4.01 [2.94-5.31]	0.5 [0.43-0.58]	2.01 [1.72-2.32]	2.81 [2.24-3.47]
East Asian (n=9,092)	<10-4		6.51 [2.51-17.29]	0.38 [0.23-0.63]	2.51 [1.57-4.06]	4.06 [1.99-8.42]
Central/South Asian (n=567)	-		-	-	-	-

Middle Eastern (n=694)		-	-	-	-	-
All (n=586,608)	Prostate Cancer	<10-16	5.19 [5.09-5.31]	0.44 [0.44-0.45]	2.3 [2.28-2.33]	3.38 [3.33-3.44]
European (n=420,672)		<10-16	5.13 [4.99-5.26]	0.45 [0.44-0.45]	2.29 [2.26-2.32]	3.38 [3.31-3.44]
African (n=104,997)		<10-16	3.21 [3.06-3.37]	0.57 [0.56-0.58]	1.84 [1.79-1.88]	2.39 [2.3-2.47]
Admixed American (50,586)		<10-16	4.36 [3.99-4.82]	0.48 [0.46-0.5]	2.09 [2.0-2.2]	3.0 [2.8-3.23]
East Asian (n=9,092)		<10-16	4.23 [3.27-5.47]	0.48 [0.42-0.55]	2.03 [1.79-2.3]	2.94 [2.42-3.58]
Central/South Asian (n=567)		0.24	2.84 [0.52-13.39]	0.59 [0.27-1.37]	1.67 [0.72-3.6]	2.22 [0.61-7.17]
Middle Eastern (n=694)		0.05	1.93 [1.01-3.77]	0.72 [0.52-1.0]	1.4 [1.0-1.95]	1.64 [1.0-2.74]

Supplementary Table 3: Association of PHS290 with any, metastatic and fatal prostate cancer using genetic ancestry groups. Cox Proportional-Hazards results from association with age at prostate cancer, metastatic prostate cancer and death from prostate cancer. *P*-values reported are from univariable models using PHS290 as the sole predictor variable. Hazard ratios (HRs) are shown comparing men in various percentiles of genetic risk. HR_{80/20}: highest 20% ($\geq 80^{\text{th}}$ percentile of PHS290, using previously published thresholds for men <70 years old and no diagnosis of cancer) vs. average risk (30-70th percentile). HR_{20/50}: lowest 20% ($\leq 20^{\text{th}}$ percentile) vs. average risk. HR_{80/50}: highest 20% vs. average risk. HR_{95/50}: highest 5% ($\geq 95^{\text{th}}$ percentile) vs. average risk. Numbers in brackets are 95% confidence intervals. For subgroup analyses with less than 10 events of the endpoint, the box is marked '-'. Genetic ancestry groups are the same as described in the caption for Supplementary Table 2.

Clinical Endpoint	Black or African American	Asian	Hispanic White	Native American	Other	Pacific Islander	Unknown
Fatal Prostate Cancer	2.53 [2.14-2.92]***	0.48 [0.12-0.98]	1.02 [0.68-1.43]	1.04 [0.34-2.0]	1.87 [1.04-2.8]	1.83 [0.0-4.87]	0.72 [0.0-1.88]
Metastatic Prostate Cancer	2.85 [2.64-3.08]***	0.94 [0.59-1.38]	1.29 [1.08-1.52]*	1.47 [1.02-2.02]	1.65 [1.28-2.07]*	1.19 [0.23-2.33]	1.49 [0.73-2.31]
Any Prostate Cancer	2.33 [2.28-2.38]***	0.9 [0.8-1.01]	1.06 [1.0-1.12]	1.1 [0.98-1.21]	1.18 [1.09-1.27]*	1.0 [0.76-1.29]	0.87 [0.7-1.04]



Supplementary Table 4: Association of self-reported race/ethnicity with three prostate cancer clinical endpoints. Hazard ratios (HRs) are shown from Cox proportional hazards models with race/ethnicity as the only predictor variable and Non-Hispanic White as the reference for each group. Numbers in brackets are 95% confidence intervals.

Clinical Endpoint	p Value	Family History
Fatal Prostate Cancer	1.83 [1.53-2.17]	<10 ⁻¹²
Metastatic Prostate Cancer	1.67 [1.52-1.83]	<10 ⁻¹⁶
Any Prostate Cancer	1.93 [1.88-1.99]	<10 ⁻¹⁶

Supplementary Table 5: Association of family history with three prostate cancer clinical endpoints. Hazard ratios (HRs) are shown from Cox proportional hazards models with family history as the only predictor variable. This analysis was limited to the 374,455 participants who provided family history information in baseline survey data. Numbers in brackets are 95% confidence intervals.

Clinical Endpoint	HR _{80/20}	Family History	Non-Hispanic White	Black or African American	Hispanic White	Asian	Native American	Other	Pacific Islander	Unknown
Fatal Prostate Cancer	4.07 [3.39-4.74]***	1.67 [1.39-1.98]*	Ref	1.31 [0.73-2.35]	1.16 [0.68-1.84]	0.68 [0.07-2.77]	1.02 [0.34-2.0]	1.97 [1.01-3.21]*	1.16 [0.0-5.51]	0.72 [0.0-1.92]
Metastatic Prostate Cancer	4.06 [3.74-4.44]***	1.53 [1.39-1.68]**	Ref	1.03 [0.76-1.39]	1.25 [0.96-1.6]	0.95 [0.35-2.26]	1.21 [0.83-1.68]	1.41 [1.07-1.86]	0.85 [0.16-1.95]	1.35 [0.66-2.13]
Any Prostate Cancer	4.67 [4.55-4.79]***	1.78 [1.73-1.84]***	Ref	1.15 [1.04-1.27]*	1.08 [1.0-1.17]	1.28 [0.98-1.7]	1.02 [0.9-1.13]	1.11 [1.02-1.21]	0.89 [0.64-1.23]	0.86 [0.69-1.03]
Clinical Endpoint	PC1	PC2	PC3	PC4	PC5	PC6	PC7	PC8	PC9	PC10
Fatal Prostate Cancer	0.11 [0.0-2.57]	0.26 [0.02-2.62]	0.56 [0.01-24.6]	0.08 [0.01-0.35]*	0.19 [0.01-2.88]	3.62 [0.1-119.93]	85.52 [0.18-167982.22]	0.01 [0.0-5.19]	3.14 [0.02-604.17]	3.42 [0.06-247.61]
Metastatic Prostate Cancer	0.01 [0.0-0.06]*	0.67 [0.18-2.34]	0.79 [0.09-6.75]	0.06 [0.02-0.14]*	1.01 [0.2-4.91]	5.77 [0.64-48.68]	0.42 [0.02-21.55]	0.47 [0.02-9.1]	5.73 [0.37-121.3]	0.18 [0.02-2.04]
Any Prostate Cancer	0.07 [0.04-0.13]***	0.5 [0.34-0.74]*	1.01 [0.53-1.92]	0.72 [0.55-0.96]	2.42 [1.52-3.87]*	2.51 [1.46-4.61]*	1.5 [0.58-4.5]	0.37 [0.16-0.89]	1.11 [0.45-2.71]	0.84 [0.41-1.8]

Supplementary Table 6: Multivariable models combining self-reported Race/Ethnicity, Family History, Top 10 Population Principal Components, and PHS290 for three prostate cancer clinical endpoints. Cox proportional hazards results for association with age at death from prostate cancer, age at diagnosis of metastatic prostate cancer, and age at diagnosis with prostate cancer. *P*-values reported are from multivariable models using race/ethnicity, family history, the top 10 population principal components, and PHS290. For PHS290, effect size was illustrated via the hazard ratio (HR_{80/20}) for the highest 20% vs. lowest 20% of genetic risk. Hazard ratios for racial/ethnic groups were estimated using Non-Hispanic White as the reference. Hazard ratios for family history were for one or more first-degree relatives diagnosed with prostate cancer. This analysis was limited to the 378,366 participants who provided family history information in baseline survey data. Numbers in brackets are 95% confidence intervals. Significant predictors in the multivariable model are indicated by * ($p < 0.01$), ** ($p < 10^{-10}$) and *** ($p < 10^{-16}$). PHS290 and family history remained significant after addition of the top 10 population principal components, but Black race was no longer independently associated with fatal or metastatic prostate cancer, implying this racial information was accounted for in the top 10 principal genetic components.



Supplementary Figure 1. (A) PHS290 score density plot in prostate cancer controls for White Non-Hispanic, Black or African American, Hispanic White and Asian self-reported race/ethnicity groups. **(B)** PHS290 score density plot in prostate cancer cases for White Non-Hispanic, Black or African American, Hispanic White and Asian self-reported race/ethnicity groups. These plots do not account for age.