

Table S4: Chi-square analysis describing associations between all experimental arms and proportion of hesitant participants who cited reasons of vaccine safety or side effect concerns after message exposure.

Hesitancy to:	Intention to vaccinate		Recommend healthy adults		Recommend elderly		Recommend people with health conditions	
	No (%)	Yes (%)	No (%)	Yes (%)	No (%)	Yes (%)	No (%)	Yes (%)
Worried about the safety or side effects of the vaccine.								
DN(70%)	22.78	77.22	30.00	70.00	20.00	80.00	18.95	81.05
DN	27.74	72.26	28.81	71.19	20.43	79.57	20.92	79.08
HCW	22.86	77.14	33.33	66.67	14.85	85.15	14.55	85.45
NF	32.67	67.33	34.48	65.52	22.86	77.14	25.53	74.47
PF	21.92	78.08	31.15	68.85	14.15	85.85	20.24	79.76
RC(S)	26.67	73.33	29.51	70.49	18.18	81.82	19.78	80.22
RC(SE)	20.53	79.47	19.3	80.7	25.24	74.76	19.34	80.66
Control	18.71	81.29	38.46	61.54	23.08	76.92	20.32	79.68
DN(70%) + DN	27.04	72.96	26.56	73.44	17.12	82.88	23.56	76.44
DN(70%) + HCW	22.73	77.27	27.27	72.73	15.6	84.4	19.66	80.34
DN(70%) + NF	29.33	70.67	28.07	71.93	20.41	79.59	23.3	76.7
DN(70%) + PF	30.67	69.33	40.00	60.00	28.57	71.43	21.76	78.24
DN(70%) + RC(S)	24.32	75.68	30.77	69.23	18.89	81.11	21.69	78.31
DN(70%) + RC(SE)	27.86	72.14	24.44	75.56	18.45	81.55	19.35	80.65
Pearson chi-square:	16.7902		9.7281		14.3616		9.3312	
P-Value:	0.209		0.716		0.349		0.747	