

Supplementary material




Figure S1. Flowchart of data selection




Figure S2. Proportion of SGTF among positive SARS-CoV-2 cases by age group, the Netherlands, 22 November 2021 - 31 March 2022, n = 941,692.

Table S1 WGS variant found in SGTF (S result, not detected) and non-SGTF (S result, detected) cases in each cohort. First column indicates the cohort, followed by the SGTF result. The number of WGS typed Omicron BA.1, Delta and BA.2 cases. Row-wise calculations on positive predictive value (PPV) and column-wise calculations on sensitivity.

Cohort	S result	WGS result Omicron BA.1	WGS result Delta	WGS result Omicron BA.2	PPV	Sensitivity
Cohort Delta-Omicron BA.1	Not detected	132	0	0	1.00	1.00
Cohort Delta-Omicron BA.1	Detected	0	353	0	1.00	1.00
Cohort Omicron BA.1-BA.2	Not detected	158	0	0	1.00	0.99
Cohort Omicron BA.1-BA.2	Detected	2	0	128	0.98	1.00

Table S2 Relative reduction in Delta and Omicron BA.1 infections after previous infection, primary vaccination, booster vaccination, or combinations of previous infection and vaccination (column 'Vaccination_and_previous_infection_status'), compared with naïve status $((1-OR) * 100)$, by time since last event and overall (column 'Time_since_event') in persons aged 18 and by variant (column 'variant'), for cohort Delta-Omicron BA.1 and cohort Omicron BA.1-BA.2 (column 'Cohort'). Estimates and 95%-CI (in brackets) are calculated for all cases (column 'VE_CI') and cases with reported symptoms (column 'VE_CI_symptomatic_infection').

Table S3 Protection estimates for booster vaccination and booster vaccination with a previous infection with 'Primary vaccinated' as reference.

Cohort	Immuunstatus	Variant	VE_CI
Delta-Omicron BA.1	Booster	Omicron BA.1	42% (38-46)
Delta-Omicron BA.1	Booster	Delta	73% (70-77)
Delta-Omicron BA.1	Previous infection, booster	Omicron BA.1	53% (45-61)
Delta-Omicron BA.1	Previous infection, booster	Delta	92% (83-96)
Omicron BA.1-BA.2	Booster	Omicron BA.1	47% (45-48)
Omicron BA.1-BA.2	Booster	Omicron BA.2	40% (38-42)
Omicron BA.1-BA.2	Previous infection, booster	Omicron BA.1	68% (66-70)
Omicron BA.1-BA.2	Previous infection, booster	Omicron BA.2	67% (65-69)

Table S4 Relative reduction in Delta and Omicron BA.1 infections after previous infection, primary vaccination, booster vaccination, or combinations of previous infection and vaccination (column Vaccination_and_previous_infection_status), compared with naïve status $((1-OR) * 100)$, by time since last event and overall (column Time_since_event), by age group (column Age_group) and by variant (column 'variant'), for cohort Delta-Omicron BA.1 and cohort Omicron BA.1-BA.2 (column Cohort).

Table S5 Interaction terms between age and vaccination and previous infection status per cohort and variant with age group 18-29 years old as reference group.