

Supplementary material 1 to:

Comparing health care use after hospital visit for SARS-CoV-2, respiratory syncytial virus, and other respiratory tract infections in children, Larsen et al. 2022

Table of contents

Table S1-1: Selected International Classification of Diseases (ICD-10) codes and diagnosis categories	1
Table S1-2: Descriptive statistics	1
Table S1-3: Crude percentages of primary and specialist health care use by age and diagnosis group ..	2
Table S1-4: Impacts of being hospitalized with COVID-19 on outpatient and inpatient specialist health care use in children, using children hospitalized with RSV infection and other RTI as comparison group	2
Table S1-5: Crude percentages of outpatient and inpatient specialist care use by age and diagnosis group	2

Table S1-1: Selected International Classification of Diseases (ICD-10) codes and diagnosis categories.

Category	ICD-10 codes
COVID-19	U07.1 U07.2
RSV infection	J12.1 J20.5 J21.0
Other RTI	J00 J02.0 J02.8 J02.9 J03.0 J03.8 J03.9 J04.0 J04.1 J04.2 J05.0 J05.1 J06.0 J06.8 J06.9 J12.0 J12.2 J12.3 J12.8 J12.9 J13 J14 J15.0 J15.1 J15.2 J15.3 J15.4 J15.5 J15.6 J15.7 J15.8 J15.9 J16.0 J16.8 J17.0 J17.1 J17.2 J17.3 J17.8 J18.0 J18.1 J18.2 J18.8 J18.9 J22 J20.0 J20.1 J20.2 J20.3 J20.4 J20.6 J20.7 J20.8 J20.9 J09 J10.0 J10.1 J10.8 J11.0 J11.1 J11.8

Table S1-2: Descriptive statistics.

	1-12 months			1-5 years		
	COVID-19	RSV	Other RTI	COVID-19	RSV	Other RTI
Children, N			34,214			
Children, N		12,058			23,682	
Children, N	53	2,438	9,600	74	1,540	22,362
Contacts, N	53	2,438	9,695	75	1,571	24,762
Person-weeks, N	1,141	53,346	220,617	1,742	37,824	607,009
Females, N (%)	23 (43.4)	1,072 (44.0)	3,975 (41.0)	33 (44.0)	670 (42.6)	10,313 (41.6)
Age, median [25%-75%]	2 [2-4]	2 [1-5]	3 [2-7]	2.4 [1.55-3.5]	1.6 [1.2-2.3]	2.1 [1.4-3.2]
Avg. length of stay, days (SD)	1.43 (3.98)	3.40 (3.62)	1.05 (5.02)	1.15 (1.67)	3.11 (4.31)	0.93 (2.53)

Note: Age presented as median and interquartile range (IQR) age in months/years at admission. The calculation of age and sex percentages are done using the number of unique hospital visits in the denominator. Due to the 6-month washout period, some children appear in more than one of the age and/or diagnosis groups.

Table S1-3: Crude percentages of primary and specialist health care use by age and diagnosis group.

Week	1-12 months						1-5 years					
	COVID-19		RSV		Other RTI		COVID-19		RSV		Other RTI	
	P	S	P	S	P	S	P	S	P	S	P	S
Pre	11.76	9.68	12.21	9.26	10.83	10.98	6.00	6.67	12.95	5.64	10.33	5.82
1-4	16.04	10.85	9.44	8.25	10.39	10.3	11.67	12.33	10.26	9.55	10.85	9.45
5-8	8.38	5.24	7.52	4.69	8.65	7.16	6.02	7.23	8.29	7.57	7.74	7.41
9-12	6.96	5.70	7.62	3.45	8.26	6.10	3.21	5.50	7.42	6.76	7.19	6.97

Note: Weekly percentages of children utilizing primary (P) and specialist (S) health care in the 12-week pre-admission period (pre) and 1-4-, 5-8-, and 9-12-week post-discharge period.

Table S1-4: Impacts of hospital visit for COVID-19 on outpatient and inpatient specialist health care use in children, using children hospitalized with RSV infection and other RTI as comparison group.

	Week 1-4		Week 5-8		Week 9-12	
	β	95 % CI	β	95 % CI	β	95 % CI
Outpatient care use						
1-11 months						
RSV infection	0.025	-0.023-0.072	-0.008	-0.076-0.06	0.012	-0.062-0.086
Other RTI	0.017	-0.025-0.06	-0.015	-0.073-0.044	0.002	-0.062-0.065
1-5 years						
RSV infection	0	-0.035-0.035	-0.019	-0.061-0.023	-0.018	-0.062-0.026
Other RTI	0.006	-0.028-0.04	-0.021	-0.064-0.022	-0.028	-0.066-0.011
Inpatient care use						
1-12 months						
RSV infection	0.044*	0.012-0.076	0.037*	0.004-0.069	0.038*	0.001-0.076
Other RTI	0.021	-0.005-0.048	0.019	-0.007-0.044	0.03	-0.002-0.062
1-5 years						
RSV infection	0.002	-0.029-0.033	-0.001	-0.033-0.031	-0.006	-0.035-0.023
Other RTI	0.002	-0.028-0.033	0.001	-0.033-0.035	-0.007	-0.033-0.02

Notes: Differences-in-differences estimates (β , in percentage points) for the change in the percentage of children utilizing outpatient and inpatient specialist health care per week after the day of discharge. The 12-week pre-admission period for the comparison group (RSV infection or other RTI) is used as reference category. Average outpatient care in the 12-week pre-admission period by age and diagnosis group is given in Table S1-5.

*Statistically significant on 5 % level.

Table S1-5: Crude percentages of outpatient and inpatient specialist care use by age and diagnosis group.

Week	1-12 months						1-5 years					
	COVID-19		RSV		Other RTI		COVID-19		RSV		Other RTI	
	O	I	O	I	O	I	O	I	O	I	O	I
Pre	6.83	4.36	5.51	4.62	6.69	5.50	5.89	1.67	4.67	1.17	4.85	1.34
1-4	9.43	4.25	6.51	2.03	7.85	3.36	9.67	3.33	7.77	2.31	7.48	2.62
5-8	4.19	2.62	3.66	1.16	5.62	2.09	5.62	2.41	5.86	2.03	6.10	1.83
9-12	5.06	3.16	2.66	0.88	4.87	1.62	4.13	1.38	5.44	1.56	5.82	1.59

Note: Weekly percentages of children utilizing outpatient (O) and inpatient (I) specialist health care in the 12-week pre-admission period (pre) and 1-4-, 5-8-, and 9-12-week post-discharge period.