Supplementary Materials

Supplementary Table 1. Search strategy used for identification of articles on Embase and PubMed Central.

Database	Search Strategy
Embase	('alcoholism'/exp OR 'alcohol consumption'/exp OR 'alcohol use disorders identification test'/exp OR 'alcohol psychosis'/exp OR 'alcohol liver disease'/exp OR 'alcohol withdrawal syndrome'/exp OR 'alcoholic cardiomyopathy'/exp OR 'alcohol abuse'/exp OR 'alcoholic pancreatitis'/exp OR 'fetal alcohol syndrome'/exp OR 'alcohol intoxication'/exp OR 'alcohol abstinence'/exp OR 'acamprosate' OR 'disulfiram' OR 'naltrexone' OR 'varenicline' OR 'topiramate' OR 'gabapentin') AND ('machine learning'/exp OR 'artificial intelligence'/exp OR 'decision support system'/exp)
PubMed Central	("alcohol related disorders"[Mesh] OR "alcoholics"[Mesh] OR "alcohol withdrawal delirium"[Mesh] OR "alcohol withdrawal seizures"[Mesh] OR "drinking behavior"[Mesh] OR "alcohol deterrents"[Mesh]) AND ("algorithms"[Mesh] OR "machine learning"[Mesh] OR "natural language processing"[Mesh] OR "neural networks, computer"[Mesh] OR "decision support systems, clinical"[Mesh])

ľ.	Supplementary	Table 2.	Overall so	coring f	for risk	of bias	assessment.
Г							

Risk of Bias Score	Rationale
"LOW"	Data source: multiple sources, or a national database. Missing data: none. External validation: performed.
"MODERATE"	Missing data: moderate amount of data missing. All categories: not specified.
"HIGH"	Data source: data collection from only one clinical site. Missing data: high amount of data missing. External validation: no external validation performed.
Overall Risk of Bias Score: "LOW"	No categories with "HIGH" risk of bias.
Overall Risk of Bias Score: "MODERATE"	One category with "HIGH" risk of bias, or one or two categories with "MODERATE" risk of bias.
Overall Risk of Bias Score: "HIGH"	Two or more categories with "HIGH" risk of bias, or three or more categories with "MODERATE" risk of bias.

First Author, Year	Type of ML algo	Sample Size	Data Type Ethnicity/race		Age	Gender	Comparator or competitor				
	Screening										
Afshar, 2019	NLP; LR, PAC, SVM, SGD	1422	EHR, including clinical notes	54% White; 18% Hispanic	Range ≥18	71% male	Rules based pre-selected keyword approach (keywords chosen by content experts); AUDIT				
To, 2020	LR with LASSO; NLP	2422	EHR, including clinical notes	58% White	Range ≥18	N/S	None				
Bonnell, 2020	LR, SVM; NN, k-nearest neighbors, DT, RF (RF used for final model)	43545	Demographic, clinical and laboratory information	N/S	Range ≥18	49% male	AUDIT-C				
Hassanpour, 2019	CNN and LSTM	2287	Instagram pictures and captions	White, Black, Asian, Hispanic	Range 18+	Males and females	None				
Crocamo, 2020	SVM and RF	180	Tweet content	N/S	N/S	N/S	None				
		Di	iagnosis/identification/discrimi	nation of AUD							
Maurelli, 1998	Genetic algorithm BEAGLE; 7 ANNs; Metasystem taking outputs from ANNs to produce new results	113	Blood markers: MCV, ALKPHOS, ALT, AST, GGT	N/S	N/S	N/S	None				
Dinevski, 2011	DT	482	Blood markers: MCV, GGT, GLDH, AST, ALT	N/S	Range 18-65	82-84% male	Individual biomarkers				
Pirro, 2013	UNEQ model	423	Blood markers: AST, ALT, GGT, MCV, CDT	N/S	Range 19-74	~80% male	Individual biomarkers				
Lewenstein, 2020	ANN	172	Polysomnography (EEG, EOG, EMG)	N/S	Range 30-46	50% male	None				
Guggenmos, 2018	Weighted robust distance-to-centroi d classifier	216	MRI	N/S	Range 20-65	84% male	None				
Kamarajan, 2020	RF	60	fMRI	42% Black; 15% White; 35% Asian; 2% American Indian	Mean 41 (AUD); 27 (control)	100% male	None				
Gowin, 2020	RF	486	fMRI and self-reported psychological / substance use history	AUD: 83% White; 9% Black; 7% Hispanic; Control: 69% White; 20% Black; 9% Hispanic	Mean 28±3 (AUD); 29±4 (control)	72% (binge-drinke rs); 33% (non-bingers)	None				
Hahn, 2020	LR and SVM	2034	MRI	N/S	Mean 45 (test-AUD); 42	57% male	None				

Supplementary Table 3. Study design details of selected articles.

					(test-control)		
Wang, 2019	AlexNet transfer learning model	379	MRI	N/S	N/S	N/S	None
Zhu, 2018	RF	92	fMRI	AUD: 54% White; 41% Black; 4% Asian; Control: 63% White; 28% Black; 9% Asian	Mean 40±10 (AUD); 32 ±9 (control)	72% male (AUD), 57% male (control)	None
Wang, 2017	CNN	235	fMRI	N/S	N/S	N/S	None
Adeli, 2019	LR	549	fMRI	7% Asian; 21% Black; 61% White; 11% Other	Mean 48 ±10 (AUD); 47±14 (control); 51±9 (HIV); 52±7 (AUD+HIV)	54% male (control), 66% male (AUD), 69% male (HIV), 63% male (AUD+HIV)	None
Bae, 2017	SVM	60	EEG	N/S	N/S	N/S	None
Acharya, 2012	SVM	122	EEG	N/S	N/S	N/S	None
Mehla, 2020	SVM	122	EEG	N/S	N/S	N/S	None
Hussain, 2018	k-D tree	58	EEG	N/S	Mean 36±5 (AUD); 26±3 (control)	N/S	None
Padma Shri, 2017	k-NN	1200	EEG	N/S	N/S	N/S	None
Mumtaz, 2017	LR	45	EEG	N/S	Mean 55±13 (AUD); 43±16 (control)	N/S	None
Mumtaz, 2018	SVM	60	EEG	N/S	N/S	N/S	None
Kumar, 2016	SVM and FCM	40	EEG	N/S	Range 32-38	100% male	None
Kumar, 2015	SVM and FCM	40	EEG	N/S	Mean 35	100% male	None
Zhang, 2020	CNN and SVM	125	EEG	N/S	Mean 36±5 (AUD); N/A (controls)	100% male	None
Anuragi, 2019	LS-SVM	122	EEG	N/S	N/S	N/S	None
Anuragi, 2020	LS-SVM	122	EEG	N/S	N/S	N/S	None
Faust, 2013	FSC	122	EEG	N/S	N/S	N/S	None
Zhu, 2014	HVGE	122	EEG	N/S	N/S	N/S	None
Chen, 2020	GS-SVM, CNN, CNN-LSTM	297	SNPs, demographic information	Chinese	Range Adults	100% male	None
Yu, 2005	SVM	1204	Microsatellite markers	N/S	N/S	N/S	None
Rosato, 2019	RF	120	microRNA	53% European American; 47%	Range 18+	47% male	None

				African American					
Falk, 2005	ANN	650	Risk factors, drinking habits and phenotypic measurements	N/S	N/S	N/S	None		
Li, 1999	ANN	699	Phenotypic ERP300 data; genotypic data; biomarker: MAO; gender	N/S	N/S	N/S	None		
Predicting AUD severity									
Fede, 2019	RF	59	fMRI	N/S	Range 22-60	68% male (primary); 50% male (validation)	None		
			Risk factors for alcoho	l use					
Winham, 2016	RF	3829	SNPs	69% European American; 31% African American	Range 18-77	46% male	None		
	Predicting future alcohol use								
King, 2011	LR	8655	Demographic, medical and psychological history, living conditions, AUDIT questionnaire; behavioral questions; alcohol, drugs and cigarette use; history of abuse; religious beliefs; family history of mental health; stress, anxiety and panic syndrome; List of Threatening Life Experiences Questionnaire; experiences in discrimnation; social support	European set: 97% White European Chilean set: 100% Non-white	Range 18-75	European set: 31% male; Chilean set: 24% male	None		
Kinreich, 2019	SVM	656	EEG, SNPs	European American and African American	Range 12–30	57% male	None		
			Prediction of Treatment O	utcomes					
Symons, 2020	RF, BayesNet, NB, SPegasos, RBFN, SMO, REPTree, DTNB, FURIA, Decision table	2236	Demographic, medical history, alcohol dependence measures, and psychological measures	N/S	Range 18–76	65-71% male	10 psychologists and traditional linear regression		
Symons, 2019	FURIA, ADT, LWL, JRip, RIDOR, SMO, REPTree, SPegasos, Decision Table, RBFNM, BayesNet, DTNB, BFTree	830	Demographic and psychometric assessment data, medical history and drinking measures	N/S	Range 18-76	N/S	10 addiction therapists		

Lindner, 2020	RF	607	Drinking measures	N/S	N/S	Males and females	None
Sekutowicz, 2019	linear SVM	188	fMRI	N/S	Mean 45	84% male	None
Seo, 2015	NB, SVM, LVQ	46	sMRI and fMRI	N/S	Average 40	65% male	ML models that used clinical data including demographics, medical history, and socioeconomic status.
Winterer, 1998 A	ANN	78	Quantitative EEG	N/S	Mean 45-48	58% male	None
Winterer, 1998 B	ANN	78	Quantitative EEG	N/S	Mean 45-48	58% male	ML models that used clinical data including demographic data, laboratory, psychiatric and neurological measurements
Satapathy, 2020	LR	241	Demographic, medical and previous rehabilitation history	73% Caucasian; 17% Black; 8% Hispanic; 0.4% Other	Mean 53	81% male	None
Chih, 2014	Bayesian network model	152	Behavioral and psychological measures, substance use, and lapse history	83% Caucasians	Range 20-64	62% male	None
Connor, 2007	Bayesian network model and DTs	139	Demographic, medical history, dependence severity, and drinking-related and psychological measures.	N/S	N/S	Males and females	Discriminant analysis
Kurth, 2001	LR and ANN	49	Demographic, laboratory values, and alcohol withdrawal assessment data	N/S	Range 27-66	86% male	None
Hillemacher, 2012	RF	200	Demographic, medical history, and laboratory variables	N/S	Mean 45-48	81% male	None
Burkhardt, 2020	SVM	1144	Laboratory variables, sociodemographic measures, and medical history	N/S	Mean 43	73-79% male	None
Lapuerta, 1997	ANN and LR	144	Laboratory values and clinical variables	66% Hispanic; 25% White; 5% African-American	Mean 40	83% male	Maddrey score
Choi, 2020	LR, DT, and MLP	8937	Demographic, psychological and behavioral data	East Asian	<65; 65+	Males and females	None
Hou, 2015	IT, TRM, and VT	251	SNPs	Hispanic and others	Range 20-78	N/S	None
Wei, 2020	IT-LT	251	SNPs	Hispanic and others	Range 20-78	N/S	None
Laska, 2020	RF	338	Demographic, substance use indicators, and psychiatric characteristics	65-77% White; 16-23% Black	Mean 50	66% male	None

Lin, 2020	RF	93	Demographic, OPRM1 promoter region DNA methylation	44% African American; 36% European American	Range 50-51	100% male	None				
	Predicting treatment seeking behavior										
Lee, 2019	ADT, RF, RT	1114	Behavioral and psychological data	39-73% African American; 20-52% Caucasian; 7-13% Other	Range 21-69	64-77% male	None				
	AUD in Adolescents										
Squeglia, 2017	RF	137	sMRI, fMRI, demographic, neuropsychological data	68% Caucasian; 32% N/S	Range 12-16	56% male	None				
Whelan, 2014	Voxelwise LR, Regularized LR	2000	sMRI, fMRI, demographic, neuropsychological data	Caucasian	Range 14-18	48% male	None				
Afzali, 2019	LR, SVM, RF, NN, Lasso Reg, Ridge Reg, Elastic-net	6016	Demographic, psychopathological, personality, risk behaviors, cognitive	N/S	Range 12-16	N/S	None				
Ruan, 2019	SVM	2000	rsMRI, fMRI, rsFC, SNPs, personality, cognitive	Caucasian	Range 14-19	49% male	None				
Vázquez, 2020	Elastic net, k-nearest neighbors, NN, RF	191880	Demographic, socioecological, personality,	Mexican	Range 10-12	51% male	None				
García, 2009	ANN, Decision trees, Naive bayes	7030	Personality variables from survey	Mexican	Range 14-18	N/S	None				
Weidacker, 2020	RVR	3064	sMRI, fMRI, demographic, behavioural data	N/S	Range 15-26	67% male	None				
O'Halloran, 2020	Elastic net	79	EEG, demographic, personality, drinking behavior, task-based tests	N/S	Range 18+	49% male	None				
O'Halloran, 2018	Elastic net	106	Demographic, behavioural, personality	N/S	Range 18-21	66% male	None				
Martinez-Loredo, 2018	Binary logistic regression	1239	Demographic, behavioural, personality	Spaniards	Range <15	54% male	None				

<u>Abbreviations</u>: Alanine aminotransferase (ALT); Alcohol use disorder (AUD); Alcohol Use Disorder Identification test (AUDIT); Alkaline phosphates (ALKPHOS); Alternating decision tree (ADT); Artificial neural network (ANN); Aspartate aminotransferase (AST); Biologic Evolutionary Algorithm Generating Logical Expressions (BEAGLE); Best-first decision tree (BFTree); Carbohydrate-deficient transferrin (CDT); Convolutional neural network (CNN); Decision table naive bayes (DTNB); Decision tree (DT); Electroencephalographic (EEG); Electromyogram (EMG); Electronic health record (EHR); Electrooculograms (EOG); Event-related potential (ERP); Functional MRI (fMRI); Fuzzy C-mean (FCM); Fuzzy sugeno classifier (FSC); Fuzzy unordered rule induction algorithm (FURIA); Gamma-glutamil transpeptidasis (GGT); Glutamate dehydrogenase (GLDH); Horizontal visibility graph entropy (HVGE); Interaction tree (IT); Interaction tree for longitudinal trajectories (IT-LT); Learning vector quantization (LVQ); Least Absolute Shrinkage and Selection Operator (LASSO); Locally weighted learning (LWL); Logistic regression (LR); Long short-term memory (LSTM); Magnetic resonance imaging (MRI); Mean corpuscular volume (MCV); Monoamine oxidase (MAO); Multilayer perceptron model (MLP); Multivariate unequal dispersed classes (UNEQ); Naïve Bayes (NB); Natural language processing (NLP); Neural network (NN); Passive Aggressive Classifier (PAC); Radial basis function neural networks (RBFN); Random forest (RF); Random tree (RT); Reduced-error pruning tree (REPTree); Relevant Vector Regression (RVR); Resting-state functional connectivity (rsFC); Ripple down rule learner (RIDOR); Sequential minimal optimization (SMO); Single nucleotide polymorphism (SNP); Support vector machine (SVM); Traditional regression method (TRM); Virtual twins (VT)

First Author, Year	Sample source	Missing data	External Validation	Other sources	Overall risk					
	Screening									
Afshar, 2019	HIGH; 1 source	LOW	HIGH: no	MODERATE: based on clinical notes analyzed from 1 center, and language variation in clinical notes in other locations may vary	HIGH					
To, 2020	HIGH; 1 source	LOW	HIGH: no	MODERATE: based on clinical notes analyzed from 1 center, and language variation in clinical notes in other locations may vary	HIGH					
Bonnell, 2020	LOW; Large dataset	LOW	HIGH: no	HIGH: did not use EHR to build algorithm, but target is to use tool on EHR, which has a much higher %age of missing data	HIGH					
Hassanpour, 2019	HIGH; Social media platform	MODERATE: N/S	HIGH: no	LOW: N/S	HIGH					
Crocamo, 2020	MODERATE; Social media platform	MODERATE: N/S	HIGH: no	HIGH: ambiguities in assessing tweets	HIGH					
		Diagnosis/identification	/discrimination of AUD	•						
Maurelli, 1998	HIGH; 1 source	MODERATE: N/S	HIGH: no	LOW: N/S	HIGH					
Dinevski, 2011	HIGH; 1 site	LOW	HIGH: no	MODERATE: Control patients and patients with alcohol dependence syndrome from different sources	HIGH					
Pirro, 2013	MODERATE; 2 sites	LOW	HIGH: no	LOW: N/S	MODERATE					
Lewenstein, 2020	HIGH; 1 source	MODERATE: N/S	HIGH: no	LOW: N/S	HIGH					
Guggenmos, 2018	LOW; Multisite study	MODERATE: N/S	LOW: yes	LOW: N/S	MODERATE					
Kamarajan, 2020	MODERATE; 1 source but from larger study	MODERATE: N/S	HIGH: no	LOW: N/S	MODERATE					
Gowin, 2020	MODERATE; 1 source but sampled from larger study	MODERATE: N/S	HIGH: no	LOW: N/S	MODERATE					

Supplementary Table 4. Risk of bias assessment of selected studies

Hahn, 2020LOW; MWang, 2019HIGHZhu, 2018LOW; NWang, 2017LOW; NWang, 2017LOW; RAdeli, 2019LOW; RBae, 2017LOW; RAcharya, 2012LOW; RInargeLOW; RMehla, 2020LOW; RHussain, 2018LOW; RPadma Shri, 2017HIGHMumtaz, 2018MODEKumar, 2016HIGH	Aultisite study H; 1 source NIH recruited W; 3 sites Repository for e datasets Repository for e datasets Repository for e datasets	MODERATE: N/S MODERATE: N/S MODERATE: N/S MODERATE: N/S MODERATE: N/S MODERATE: N/S	LOW: yes LOW: yes HIGH: no HIGH: no HIGH: no	LOW: N/S LOW: N/S LOW: N/S LOW: N/S LOW: N/S LOW: N/S	LOW MODERATE MODERATE MODERATE MODERATE
Wang, 2019HIGHZhu, 2018LOW; NWang, 2017LOW; NAdeli, 2019LOW; RBae, 2017LOW; RAcharya, 2012LOW; RMehla, 2020LOW; RHussain, 2018LOW; RPadma Shri, 2017HIGHMumtaz, 2018MODEKumar, 2016HIGH	H; 1 source NIH recruited W; 3 sites Repository for e datasets Repository for e datasets Repository for e datasets	MODERATE: N/S MODERATE: N/S MODERATE: N/S MODERATE: N/S MODERATE: N/S MODERATE	LOW: yes HIGH: no HIGH: no HIGH: no	LOW: N/S LOW: N/S LOW: N/S LOW: N/S LOW: N/S	MODERATE MODERATE MODERATE MODERATE
Zhu, 2018LOW; NWang, 2017LOW; NAdeli, 2019Bae, 2017LOW; R largAcharya, 2012LOW; R largMehla, 2020LOW; R largHussain, 2018LOW; R largPadma Shri, 2017HIC Mumtaz, 2018Mumtaz, 2018MODE Kumar, 2015	NIH recruited W; 3 sites Repository for e datasets Repository for e datasets Repository for e datasets	MODERATE: N/S MODERATE: N/S MODERATE: N/S MODERATE: N/S MODERATE	HIGH: no HIGH: no HIGH: no HIGH: no	LOW: N/S LOW: N/S LOW: N/S LOW: N/S	MODERATE MODERATE MODERATE
Wang, 2017LOWAdeli, 2019	W; 3 sites Repository for e datasets Repository for e datasets Repository for e datasets	MODERATE: N/S MODERATE: N/S MODERATE: N/S MODERATE	HIGH: no HIGH: no HIGH: no	LOW: N/S LOW: N/S LOW: N/S	MODERATE MODERATE
Adeli, 2019Bae, 2017LOW; R largAcharya, 2012LOW; R largMehla, 2020LOW; R largHussain, 2018LOW; R largPadma Shri, 2017HIC Mumtaz, 2017Mumtaz, 2018MODE Kumar, 2015	Repository for e datasets Repository for e datasets Repository for e datasets	MODERATE: N/S MODERATE: N/S MODERATE	HIGH: no HIGH: no	LOW: N/S LOW: N/S	MODERATE
Bae, 2017LOW; R largAcharya, 2012LOW; R largMehla, 2020LOW; R largHussain, 2018LOW; R largPadma Shri, 2017HIC Mumtaz, 2017Mumtaz, 2018MODE Kumar, 2015	Repository for e datasets Repository for e datasets Repository for e datasets	MODERATE: N/S MODERATE	HIGH: no	LOW: N/S	
Acharya, 2012LOW; R largMehla, 2020LOW; R largHussain, 2018LOW; R largPadma Shri, 2017HICMumtaz, 2017HICMumtaz, 2018MODEKumar, 2016HIGH	Repository for e datasets Repository for e datasets	MODERATE		1	MODERATE
Mehla, 2020LOW; R largHussain, 2018LOW; R largPadma Shri, 2017HICMumtaz, 2017HICMumtaz, 2018MODEKumar, 2016HIGHKumar, 2015HIGH	Repository for e datasets		HIGH: no	LOW: N/S	MODERATE
Hussain, 2018LOW; R largPadma Shri, 2017HICMumtaz, 2017HICMumtaz, 2018MODEKumar, 2016HIGHKumar, 2015HIGH		MODERATE	HIGH: no	LOW: N/S	MODERATE
Padma Shri, 2017HICMumtaz, 2017HICMumtaz, 2018MODEKumar, 2016HIGHKumar, 2015HIGH	Repository for e datasets	MODERATE: N/S	HIGH: no	LOW: N/S	MODERATE
Mumtaz, 2017 HIC Mumtaz, 2018 MODE Kumar, 2016 HIGE Kumar, 2015 HIGE	GH; 1 site	MODERATE: N/S	HIGH: no	LOW: N/S	HIGH
Mumtaz, 2018 MODE Kumar, 2016 HIGE Kumar, 2015 HIGE	GH; 1 site	MODERATE: N/S	HIGH: no	LOW: N/S	HIGH
Kumar, 2016 HIGH Kumar, 2015 HIGH	ERATE; N/S	MODERATE: N/S	HIGH: no	LOW: N/S	HIGH
Kumar, 2015 HIGH	H; 1 source	MODERATE: N/S	HIGH: no	LOW: N/S	HIGH
	H; 1 source	MODERATE: N/S	HIGH: no	LOW: N/S	HIGH
Zhang, 2020 LOW; R large	Repository for e datasets	MODERATE: N/S	HIGH: no	LOW: N/S	MODERATE
Anuragi, 2019 LOW; R large	Repository for e datasets	MODERATE	HIGH: no	LOW: N/S	MODERATE
Anuragi, 2020 LOW; R large	Repository for e datasets	MODERATE	HIGH: no	LOW: N/S	MODERATE
Faust, 2013 LOW; R large	Repository for e datasets	MODERATE	HIGH: no	LOW: N/S	MODERATE
Zhu, 2014 LOW; R large	Repository for e datasets	MODERATE	HIGH: no	LOW: N/S	MODERATE
Chen, 2020 HIC	GH; 1 site	MODERATE: N/S	HIGH: no	LOW: N/S	HIGH
Yu, 2005 LOW; 9	9 sites across he US	MODERATE: N/S	HIGH: no	LOW: N/S	MODERATE
Rosato, 2019 HIC	GH; 1 site	MODERATE: N/S	HIGH: no	LOW: N/S	HIGH
Falk, 2005	9 sites across he US	MODERATE: N/S	HIGH: no	LOW: N/S	MODERATE
Li, 1999 LOW; 9	9 sites across he US	MODERATE: N/S	HIGH: no	LOW: N/S	MODERATE
		Predicting A	UD severity		
Fede, 2019 MOD cor					

Risk factors for alcohol use									
Winham, 2016	LOW; Multisite study	MODERATE: N/S	HIGH: no	LOW: N/S	MODERATE				
Predicting future alcohol use									
King, 2011	LOW; Large dataset across 6 countries	LOW	LOW: yes	LOW: N/S	LOW				
Kinreich, 2019	LOW; 9 sites across the US	MODERATE: N/S	HIGH: no	LOW: N/S	MODERATE				
Prediction of Treatment Outcomes									
Symons, 2020	HIGH; 1 site	HIGH	HIGH: no	LOW: N/S	HIGH				
Symons, 2019	HIGH; 1 site	HIGH	HIGH: no	LOW: N/S	HIGH				
Lindner, 2020	HIGH; 1 site	HIGH	HIGH: no	LOW: N/S	HIGH				
Sekutowicz, 2019	LOW; Multisite study	MODERATE: N/S	HIGH: no	LOW: N/S	MODERATE				
Seo, 2015	MODERATE; N/S	MODERATE: N/S	HIGH: no	LOW: N/S	HIGH				
Winterer, 1998 A	HIGH; 1 site	MODERATE: N/S	MODERATE: N/S	LOW: N/S	HIGH				
Winterer, 1998 B	HIGH; 1 site	MODERATE: N/S	MODERATE: N/S	LOW: N/S	MODERATE				
Satapathy, 2020	HIGH; 1 site	HIGH	HIGH: no	MODERATE: Patients with extensive history of prior rehabilitation did receive transplants	HIGH				
Chih, 2014	LOW; 2 sites	LOW	LOW: yes	LOW: N/S	LOW				
Connor, 2007	MODERATE; N/S	LOW	MODERATE: N/S	LOW: N/S	MODERATE				
Kurth, 2001	MODERATE; N/S	MODERATE: N/S	MODERATE: N/S	LOW: N/S	MODERATE				
Hillemacher, 2012	HIGH; 1 source	LOW	HIGH: no	LOW: N/S	HIGH				
Burkhardt, 2020	LOW; 2 sites	MODERATE	LOW: yes	LOW: N/S	MODERATE				
Lapuerta, 1997	MODERATE; N/S	MODERATE: N/S	LOW: yes	LOW: N/S	MODERATE				
Choi, 2020	LOW; Nationwide study Korea	MODERATE	LOW: yes	LOW: N/S	MODERATE				
Hou, 2015	LOW; 2 sites	LOW	HIGH: no	LOW: N/S	MODERATE				
Wei, 2020	LOW; 2 sites	LOW	MODERATE: N/S	LOW: N/S	MODERATE				
Laska, 2020	LOW; Multisite study	HIGH	MODERATE: N/S	LOW: N/S	MODERATE				
Lin, 2020	LOW; Multisite study	MODERATE	HIGH: no	HIGH: Cell heterogeneity from blood samples; Single population	HIGH				
		Predicting treatment	nt seeking behavior						
Lee, 2019	LOW; Multisite study	MODERATE	LOW: yes	LOW: N/S	MODERATE				

AUD in Adolescents								
Squeglia, 2017	LOW; Multisource study	LOW	HIGH: no	LOW: N/S	MODERATE			
Whelan, 2014	LOW; Multisite study	LOW	LOW: yes	LOW: N/S	LOW			
Afzali, 2019	LOW; 2 sites	LOW	HIGH: no	LOW: N/S	MODERATE			
Ruan, 2019	LOW; Multisite study	LOW	HIGH: no	LOW: N/S	MODERATE			
Vázquez, 2020	LOW; Multisource study	HIGH	MODERATE: N/S	LOW: N/S	MODERATE			
García, 2009	MODERATE; N/S	MODERATE: N/S	HIGH: no	LOW: N/S	HIGH			
Weidacker, 2020	LOW; Multisite study	MODERATE: N/S	HIGH: no	LOW: N/S	MODERATE			
O'Halloran, 2020	HIGH; 1 source	MODERATE: N/S	HIGH: no	LOW: N/S	HIGH			
O'Halloran, 2018	LOW; 2 sites	MODERATE: N/S	HIGH: no	LOW: N/S	MODERATE			
Martinez-Loredo, 2018	LOW; 22 sites	LOW	HIGH: no	LOW: N/S	MODERATE			

Abbreviations: Electroencephalographic (EEG); Electronic health record (EHR); Not specified (NS).