

**Supplementary tables**

Table S1 **Recommended infant vaccination schedule in Manitoba**

<b>2 months</b>	<b>4 months</b>	<b>6 months</b>	<b>12 months</b>	<b>18 months</b>
DTaP	DTaP	DTaP	MMR	DTaP
Polio	Polio	HiB	Varicella	Polio
HiB	HiB	Rotavirus <sup>1</sup>	PCV	Hib
PCV	PCV		MenC-C	
Rotavirus	Rotavirus			

DTaP = Diphtheria, tetanus, and acellular pertussis vaccine

HiB = Haemophilus influenzae type b vaccine

MenC-C = Meningococcal conjugate serogroup C vaccine

MMR = Measles, mumps, and rubella vaccine

PCV = 13-valent pneumococcal conjugate vaccine

Note that some of these are administered in combination as one product, e.g., MMR and varicella.

<sup>1</sup>Only for kids born March 2018 and later. This vaccine is excluded from the analysis, because the 6-month dose was introduced shortly before the start of the pandemic.

Table S2 **Changes in vaccine uptake (%) of routine childhood vaccines during the pandemic (data ending November 30, 2021).**

<b>Polio</b>	<b>Dose 2 uptake at 12 months<sup>a</sup></b>			
	<b>Pre-pandemic</b>	<b>Pandemic</b>	<b>Relative change</b>	<b>Fold change</b>
Overall	88	83	-6%	
Gender				
Male	88	82	-7%	ref.
Female	88	83	-6%	0.8
Income quintile				
Q1 (lowest)	85	77	-9%	2.9
Q2	89	83	-7%	2.2
Q3	90	85	-5%	1.5
Q4	88	83	-6%	1.7
Q5 (highest)	91	88	-3%	ref.
Unknown	89	86	-3%	1.0
Residence				
Rural	81	71	-12%	6.0
Urban	94	92	-2%	ref.
Unknown	88	100	14%	-7.0
Winnipeg region of residence				
Northern suburbs	95	92	-3%	2.1
Inner city	91	88	-3%	2.3
Southern suburbs	95	94	-1%	ref.
Regional health authority of residence				
Winnipeg	94	92	-2%	ref.
Interlake-Eastern	86	76	-13%	6.1
Northern	84	72	-15%	7.2
Southern	73	65	-12%	5.9
Prairie Mountain	89	85	-5%	2.3
Public Trustee / In CFS care	88	100	14%	-6.8
	<b>Dose 3 uptake at 18 months<sup>b</sup></b>			
<b>PCV</b>	<b>Pre-pandemic</b>	<b>Pandemic</b>	<b>Relative change</b>	<b>Fold change</b>
Overall	78	72	-8%	
Gender				
Male	79	72	-8%	ref.

Female	78	73	-7%	0.9
<b>Income quintile</b>				
Q1 (lowest)	71	64	-11%	2.9
Q2	80	72	-10%	2.7
Q3	81	76	-7%	1.8
Q4	80	74	-7%	1.9
Q5 (highest)	84	80	-4%	ref.
Unknown	81	78	-3%	0.7
<b>Residence</b>				
Rural	70	63	-10%	1.7
Urban	85	80	-6%	ref.
Unknown	78	55	-29%	4.9
<b>Winnipeg region of residence</b>				
Northern suburbs	86	81	-5%	0.9
Inner city	79	73	-8%	1.4
Southern suburbs	87	82	-6%	ref.
<b>Regional health authority of residence</b>				
Winnipeg	85	80	-6%	ref.
Interlake-Eastern	76	68	-10%	1.7
Northern	70	61	-13%	2.2
Southern	64	58	-9%	1.6
Prairie Mountain	80	73	-9%	1.5
Public Trustee / In CFS care	78	55	-29%	5.1

**Dose 1 uptake at 24 months<sup>c</sup>**

<b>MenC-C</b>	<b>Pre-pandemic Pandemic Relative change Fold change</b>			
Overall	86	80	-7%	
<b>Gender</b>				
Male	86	80	-7%	ref.
Female	86	80	-7%	1.1
<b>Income quintile</b>				
Q1 (lowest)	83	73	-11%	4.8
Q2	88	79	-9%	4.0
Q3	87	84	-3%	1.2
Q4	87	81	-7%	2.8
Q5 (highest)	89	87	-2%	ref.
Unknown	87	86	-1%	0.6
<b>Residence</b>				

Rural	80	70	-12%	4.1
Urban	92	89	-3%	ref.
Unknown	77	53	-31%	10.9
Winnipeg region of residence				
Northern suburbs	93	90	-3%	2.0
Inner city	87	83	-5%	3.6
Southern suburbs	93	92	-1%	ref.
Regional health authority of residence				
Winnipeg	92	89	-3%	ref.
Interlake-Eastern	83	78	-6%	2.2
Northern	85	71	-16%	6.3
Southern	72	63	-12%	4.6
Prairie Mountain	87	79	-10%	3.7
Public Trustee / In CFS care	77	53	-31%	12.0

#### Dose 1 uptake at 24 months<sup>c</sup>

Varicella	Dose 1 uptake at 24 months <sup>c</sup>			
	Pre-pandemic	Pandemic	Relative change	Fold change
Overall	86	80	-7%	
Gender				
Male	85	80	-6%	ref.
Female	86	80	-7%	1.1
Income quintile				
Q1 (lowest)	82	73	-11%	5.1
Q2	87	79	-9%	4.2
Q3	86	84	-3%	1.2
Q4	86	80	-6%	2.8
Q5 (highest)	88	86	-2%	ref.
Unknown	85	85	0%	-0.1
Residence				
Rural	79	70	-11%	4.1
Urban	91	88	-3%	ref.
Unknown	81	53	-34%	12.0
Winnipeg region of residence				
Northern suburbs	92	89	-3%	3.0
Inner city	87	82	-6%	5.5
Southern suburbs	92	91	-1%	ref.
Regional health authority of residence				
Winnipeg	91	89	-3%	ref.

Interlake-Eastern	83	77	-7%	2.6
Northern	85	71	-16%	6.3
Southern	71	63	-11%	4.4
Prairie Mountain	88	80	-9%	3.4
Public Trustee / In CFS care	81	53	-34%	13.1

#### Dose 3 uptake at 24 months<sup>d</sup>

Polio	Dose 3 uptake at 24 months <sup>d</sup>			
	Pre-pandemic	Pandemic	Relative change	Fold change
Overall	87	84	-3%	
Gender				
Male	87	84	-3%	ref.
Female	87	84	-4%	1.2
Income quintile				
Q1 (lowest)	83	78	-6%	6.7
Q2	88	85	-4%	4.6
Q3	88	87	-2%	1.9
Q4	87	85	-2%	2.2
Q5 (highest)	90	89	-1%	ref.
Unknown	89	88	-1%	1.2
Residence				
Rural	80	76	-5%	3.4
Urban	93	91	-2%	ref.
Unknown	84	85	1%	-0.6
Winnipeg region of residence				
Northern suburbs	94	92	-2%	1.7
Inner city	90	86	-4%	4.1
Southern suburbs	94	93	-1%	ref.
Regional health authority of residence				
Winnipeg	93	92	-2%	ref.
Interlake-Eastern	85	82	-3%	1.9
Northern	83	75	-10%	5.5
Southern	73	70	-5%	3.0
Prairie Mountain	88	86	-2%	1.0
Public Trustee / In CFS care	84	85	1%	-0.5

#### Dose 4 uptake at 24 months<sup>d</sup>

HiB	Dose 4 uptake at 24 months <sup>d</sup>			
	Pre-pandemic	Pandemic	Relative change	Fold change
Overall	71	63	-10%	
Gender				

Male	70	63	-10%	ref.
Female	71	63	-10%	1.1
Income quintile				
Q1 (lowest)	60	49	-18%	3.9
Q2	71	60	-14%	3.1
Q3	75	70	-6%	1.3
Q4	75	69	-8%	1.7
Q5 (highest)	78	75	-5%	ref.
Unknown	70	71	2%	-0.4
Residence				
Rural	61	52	-15%	2.2
Urban	78	73	-7%	ref.
Unknown	55	42	-23%	3.4
Winnipeg region of residence				
Northern suburbs	81	75	-7%	1.5
Inner city	69	60	-13%	2.9
Southern suburbs	81	77	-5%	ref.
Regional health authority of residence				
Winnipeg	78	73	-7%	ref.
Interlake-Eastern	68	60	-12%	1.7
Northern	51	38	-25%	3.7
Southern	59	50	-16%	2.4
Prairie Mountain	74	67	-9%	1.4
Public Trustee / In CFS care	55	42	-23%	3.4

PCV = 13-valent pneumococcal conjugate vaccine

MenC-C = Meningococcal conjugate serogroup C vaccine

HiB = Haemophilus influenzae type b vaccine

<sup>a</sup> Recommended age is 4 months; Pre-pandemic cohort, born January 1, 2017 - March 15, 2019; Pandemic cohort, born November 15, 2019 and later

<sup>b</sup> Recommended age is 12 months; Pre-pandemic cohort, born January 1, 2017 - September 15, 2018; Pandemic cohort, born March 15, 2019 and later

<sup>c</sup> Recommended age is 12 months; Pre-pandemic cohort, born January 1, 2017 - March 15, 2018; Pandemic cohort, born March 15, 2019 and later

<sup>d</sup> Recommended age is 18 months; Pre-pandemic cohort, born January 1, 2017 - March 15, 2018; Pandemic cohort, born September 15, 2018 and later

Table S3 **Vaccine uptake (%) of routine childhood vaccines during the pandemic (data ending November 30, 2021).**

<b>Polio</b>	<b>Dose 2 uptake at 12 months<sup>a</sup></b>
	<b>Intermediate</b>
Overall	87
Gender	
Male	86
Female	87
Income quintile	
Q1 (lowest)	82
Q2	87
Q3	89
Q4	85
Q5 (highest)	92
Unknown	90
Residence	
Rural	79
Urban	93
Unknown	76
Winnipeg region of residence	
Northern suburbs	93
Inner city	88
Southern suburbs	94
Regional health authority of residence	
Winnipeg	93
Interlake-Eastern	84
Northern	83
Southern	72
Prairie Mountain	88
Public Trustee / In CFS care	76
<b>DTaP</b>	<b>Dose 3 uptake at 12 months<sup>b</sup></b>
	<b>Intermediate</b>
Overall	77
Gender	
Male	77

Female	78
Income quintile	
Q1 (lowest)	67
Q2	77
Q3	81
Q4	79
Q5 (highest)	86
Unknown	85
Residence	
Rural	69
Urban	84
Unknown	58
Winnipeg region of residence	
Northern suburbs	86
Inner city	78
Southern suburbs	87
Regional health authority of residence	
Winnipeg	85
Interlake-Eastern	75
Northern	62
Southern	66
Prairie Mountain	78
Public Trustee / In CFS care	58

**Dose 1 uptake at 18 months<sup>c</sup>**

MMR	Intermediate
	Overall
Gender	
Male	81
Female	81
Income quintile	
Q1 (lowest)	73
Q2	82
Q3	84
Q4	84
Q5 (highest)	86
Unknown	81
Residence	

Rural	73
Urban	88
Unknown	82
Winnipeg region of residence	
Northern suburbs	90
Inner city	80
Southern suburbs	89
Regional health authority of residence	
Winnipeg	88
Interlake-Eastern	80
Northern	71
Southern	66
Prairie Mountain	84
Public Trustee / In CFS care	82
<b>Dose 3 uptake at 18 months<sup>c</sup></b>	
<b>PCV</b>	<b>Intermediate</b>
Overall	77
Gender	
Male	78
Female	76
Income quintile	
Q1 (lowest)	70
Q2	77
Q3	80
Q4	81
Q5 (highest)	81
Unknown	76
Residence	
Rural	69
Urban	83
Unknown	82
Winnipeg region of residence	
Northern suburbs	85
Inner city	77
Southern suburbs	85
Regional health authority of residence	
Winnipeg	84

Interlake-Eastern	77
Northern	65
Southern	63
Prairie Mountain	80
Public Trustee / In CFS care	82

<b>MenC-C</b>	<b>Dose 1 uptake at 24 months<sup>d</sup></b>
	<b>Intermediate</b>
Overall	85
Gender	
Male	85
Female	84
Income quintile	
Q1 (lowest)	80
Q2	85
Q3	87
Q4	85
Q5 (highest)	88
Unknown	84
Residence	
Rural	77
Urban	91
Unknown	87
Winnipeg region of residence	
Northern suburbs	92
Inner city	86
Southern suburbs	92
Regional health authority of residence	
Winnipeg	91
Interlake-Eastern	82
Northern	80
Southern	68
Prairie Mountain	87
Public Trustee / In CFS care	87

<b>MMR</b>	<b>Dose 1 uptake at 24 months<sup>d</sup></b>
	<b>Intermediate</b>
Overall	85
Gender	

Male	85
Female	85
Income quintile	
Q1 (lowest)	80
Q2	85
Q3	87
Q4	85
Q5 (highest)	89
Unknown	85
Residence	
Rural	77
Urban	91
Unknown	87
Winnipeg region of residence	
Northern suburbs	92
Inner city	86
Southern suburbs	92
Regional health authority of residence	
Winnipeg	91
Interlake-Eastern	83
Northern	80
Southern	69
Prairie Mountain	88
Public Trustee / In CFS care	87

**Dose 1 uptake at 24 months<sup>d</sup>**

<b>Varicella</b>	
	<b>Intermediate</b>
Overall	84
Gender	
Male	84
Female	84
Income quintile	
Q1 (lowest)	80
Q2	85
Q3	86
Q4	84
Q5 (highest)	88
Unknown	85

Residence	
Rural	77
Urban	90
Unknown	87
Winnipeg region of residence	
Northern suburbs	92
Inner city	86
Southern suburbs	91
Regional health authority of residence	
Winnipeg	90
Interlake-Eastern	82
Northern	80
Southern	68
Prairie Mountain	87
Public Trustee / In CFS care	87

**Dose 3 uptake at 24 months<sup>e</sup>**

<b>Polio</b>	<b>Intermediate</b>
Overall	86
Gender	
Male	87
Female	86
Income quintile	
Q1 (lowest)	82
Q2	87
Q3	88
Q4	86
Q5 (highest)	91
Unknown	90
Residence	
Rural	78
Urban	93
Unknown	83
Winnipeg region of residence	
Northern suburbs	94
Inner city	89
Southern suburbs	94
Regional health authority of residence	

Winnipeg	93
Interlake-Eastern	84
Northern	81
Southern	70
Prairie Mountain	88
Public Trustee / In CFS care	83

<b>DTaP</b>	<b>Dose 4 uptake at 24 months<sup>e</sup></b>
	<b>Intermediate</b>
Overall	69
Gender	
Male	70
Female	68
Income quintile	
Q1 (lowest)	56
Q2	70
Q3	74
Q4	74
Q5 (highest)	78
Unknown	72
Residence	
Rural	59
Urban	77
Unknown	50
Winnipeg region of residence	
Northern suburbs	79
Inner city	68
Southern suburbs	80
Regional health authority of residence	
Winnipeg	77
Interlake-Eastern	64
Northern	51
Southern	55
Prairie Mountain	75
Public Trustee / In CFS care	50

<b>HiB</b>	<b>Dose 4 uptake at 24 months<sup>e</sup></b>
	<b>Intermediate</b>
Overall	69

Gender	
Male	69
Female	68
Income quintile	
Q1 (lowest)	56
Q2	69
Q3	74
Q4	73
Q5 (highest)	77
Unknown	71
Residence	
Rural	59
Urban	77
Unknown	50
Winnipeg region of residence	
Northern suburbs	79
Inner city	68
Southern suburbs	79
Regional health authority of residence	
Winnipeg	77
Interlake-Eastern	63
Northern	51
Southern	54
Prairie Mountain	74
Public Trustee / In CFS care	50

DTaP = Diphtheria, tetanus, and acellular pertussis vaccine

MMR = Measles, mumps, and rubella vaccine

PCV = 13-valent pneumococcal conjugate vaccine

MenC-C = Meningococcal conjugate serogroup C vaccine

HiB = Haemophilus influenzae type b vaccine

<sup>a</sup> Recommended age is 4 months; Intermediate cohort, born March 15, 2019 - November 15, 2019

<sup>b</sup> Recommended age is 6 months; Intermediate cohort, born March 15, 2019 - September 15, 2019

<sup>c</sup> Recommended age is 12 months; Intermediate cohort, born September 15, 2018 - March 15, 2019

<sup>d</sup> Recommended age is 12 months; Intermediate cohort, born March 15, 2018 - March 15, 2019

<sup>e</sup> Recommended age is 18 months; Intermediate cohort, born March 15, 2018 - September 15, 2018

Table S4 **Changes in vaccine uptake (%) of routine childhood vaccines during the pandemic according to region of residence and income quintile (data ending November 30, 2021).**

DTaP	Dose 3 uptake at 12 months <sup>a</sup>				Dose 4 uptake at 24 months <sup>b</sup>			
	Pre-pandemic	Pandemic	Relative change	Fold change	Pre-pandemic	Pandemic	Relative change	Fold change
Overall	80	74	-7%		71	64	-10%	
Winnipeg								
Q1 (lowest)	81	77	-5%	-47.6	70	62	-12%	4.7
Q2	87	85	-3%	-25.2	79	72	-9%	3.5
Q3	89	87	-3%	-25.4	81	79	-4%	1.4
Q4	91	88	-3%	-33.7	84	80	-4%	1.8
Q5 (highest)	91	91	0%	ref.	85	83	-2%	ref.
Interlake-Eastern								
Q1 (lowest)	70	45	-36%	3.6	63	40	-36%	37.8
Q2	84	77	-8%	0.8	71	68	-4%	4.1
Q3	78	68	-13%	1.3	61	58	-5%	5.7
Q4	81	77	-4%	0.4	70	70	0%	-0.1
Q5 (highest)	84	75	-10%	ref.	77	77	-1%	ref.
Northern								
Q1 (lowest)	51	40	-21%	4.1	39	30	-23%	13.6
Q2	63	41	-34%	6.7	51	30	-41%	23.9
Q3	68	56	-18%	3.5	58	40	-31%	18.1
Q4	80	71	-12%	2.3	67	54	-20%	11.9
Q5 (highest)	88	84	-5%	ref.	75	73	-2%	ref.
Southern								
Q1 (lowest)	55	43	-23%	5.7	44	33	-25%	2.2
Q2	66	55	-17%	4.3	61	49	-20%	1.8
Q3	67	62	-8%	1.9	61	54	-12%	1.0
Q4	66	56	-15%	3.6	61	51	-17%	1.6
Q5 (highest)	73	70	-4%	ref.	68	61	-11%	ref.

Prairie Mountain								
Q1 (lowest)	73	65	-11%	3.5	64	53	-17%	1.1
Q2	83	77	-8%	2.4	74	67	-9%	0.6
Q3	88	84	-5%	1.5	80	76	-5%	0.3
Q4	88	84	-5%	1.4	80	76	-5%	0.3
Q5 (highest)	85	82	-3%	ref.	84	71	-16%	ref.
<b>Dose 1 uptake at 18 months<sup>c</sup></b>				<b>Dose 1 uptake at 24 months<sup>d</sup></b>				
<b>MMR</b>	<b>Pre-pandemic</b>	<b>Pandemic</b>	<b>Relative change</b>	<b>Fold change</b>	<b>Pre-pandemic</b>	<b>Pandemic</b>	<b>Relative change</b>	<b>Fold change</b>
Overall	82	75	-9%		86	80	-7%	
Winnipeg								
Q1 (lowest)	83	77	-7%	2.2	88	83	-6%	4.8
Q2	89	85	-5%	1.5	92	89	-4%	3.5
Q3	90	88	-2%	0.6	92	92	-1%	0.6
Q4	92	89	-3%	0.9	93	91	-2%	1.8
Q5 (highest)	93	90	-3%	ref.	94	93	-1%	ref.
Interlake-Eastern								
Q1 (lowest)	76	57	-25%	7.7	82	67	-19%	-13.4
Q2	81	74	-9%	2.6	82	77	-6%	-4.4
Q3	77	70	-8%	2.5	81	80	-1%	-0.9
Q4	81	77	-5%	1.5	84	84	0%	0.3
Q5 (highest)	85	82	-3%	ref.	86	87	1%	ref.
Northern								
Q1 (lowest)	68	53	-22%	3.9	79	65	-18%	16.0
Q2	77	52	-33%	5.8	89	71	-19%	17.3
Q3	78	55	-30%	5.3	86	74	-13%	11.9
Q4	82	61	-26%	4.5	92	70	-24%	21.4
Q5 (highest)	88	83	-6%	ref.	88	87	-1%	ref.
Southern								
Q1 (lowest)	59	45	-23%	3.8	66	51	-22%	3.0
Q2	69	58	-16%	2.5	74	61	-17%	2.2
Q3	68	62	-9%	1.4	73	68	-7%	0.9

Q4	67	57	-15%	2.4	72	60	-16%	2.2
Q5 (highest)	76	71	-6%	ref.	78	72	-8%	ref.
Prairie Mountain								
Q1 (lowest)	78	67	-15%	1.7	85	72	-15%	2.5
Q2	84	73	-12%	1.4	86	76	-12%	2.0
Q3	87	81	-7%	0.7	90	85	-5%	0.8
Q4	88	83	-6%	0.6	92	87	-5%	0.9
Q5 (highest)	90	82	-9%	ref.	93	88	-6%	ref.

DTaP = Diphtheria, tetanus, and acellular pertussis vaccine

MMR = Measles, mumps, and rubella vaccine

<sup>a</sup> Recommended age is 6 months; Pre-pandemic cohort, born January 1, 2017 - March 15, 2019; Pandemic cohort, born September 15, 2019 and later

<sup>b</sup> Recommended age is 18 months; Pre-pandemic cohort, born January 1, 2017 - March 15, 2018; Pandemic cohort, born September 15, 2018 and later

<sup>c</sup> Recommended age is 12 months; Pre-pandemic cohort, born January 1, 2017 - September 15, 2018; Pandemic cohort, born March 15, 2019 and later

<sup>d</sup> Recommended age is 12 months; Pre-pandemic cohort, born January 1, 2017 - March 15, 2018; Pandemic cohort, born March 15, 2019 and later