

Supplemental Table 3. Kidney and Liver Cystic Phenotypes of ALG8 P/LP PTV heterozygotes with CT or MRI imaging, by Variant

	p.Arg364Ter Age ≥30: n=34 Age <30: n=4	p.Arg179Ter Age ≥30: n=2 Age <30: n=1	p.Arg41Ter Age ≥30: n=5 Age <30: n=0	c.368+T2>G Age ≥30: 1 Age <30: 0
Consequence	Stop-gained	Stop-gained	Stop-gained	Splice donor
Clinvar classification	Pathogenic 8 submission 2 stars	Pathogenic 1 submission 0 stars	Pathogenic 3 submissions 2 stars	Likely pathogenic 1 submission 1 star
Conditions listed in Clinvar	Polycystic liver disease 3 with kidney cysts (AD); Congenital disorder of glycosylation type 1H (AR)	Polycystic liver disease 3 with kidney cysts (AD)	Congenital disorder of glycosylation type 1H (AR)	Congenital disorder of glycosylation type 1H (AR)
≥ 4 kidney cysts	Age ≥30: 18/34 (52.9%) Age <30: 1/4 (25%)	Age ≥30: 0/2 Age <30: 0/1	Age ≥30: 2/5 (60%)	Age ≥30: 1/1 (100%)
≥ 4 kidney cysts or TSTCs	Age ≥30: 25/34 (73.5%) Age <30: 2/4 (50%)	Age ≥30: 2/2 (100%) Age <30: 0/1 (0%)	Age ≥30: 3/5 (60%)	Age ≥30: 1/1 (100%)
Bilateral kidney cysts	Age ≥30: 23/34 (67.7%) Age <30: 2/4 (50%)	Age ≥30: 1/2 (50%) Age <30: 0/1	Age ≥30: 3/5 (60%)	Age ≥30: 1/1 (100%)
Bilateral kidney cysts or TSTCs	Age ≥30: 27/34 (79.4%) Age <30: 2/4 (50%)	Age ≥30: 2/2 (100%) Age <30: 1/1 (100%)	Age ≥30: 4/5 (80%)	Age ≥30: 1/1 (100%)
Liver cysts	Age ≥30: 3/34 (8.8%) Age <30: 0/4	Age ≥30: 0/2 Age <30: 0/1	Age ≥30: 1/5 (20%)	Age ≥30: 1/1 (100%)

Liver cysts or TSTCs	Age ≥30: 10/34 (29.4%) Age <30: 1/4 (25%)	Age ≥30: 0/2 Age <30: 0/1	Age ≥30: 1/5 (20%)	Age ≥30: 1/1 (100%)
Nephrolithiasis	Age ≥30: 18/34 (52.9%) Age <30: 1/4 (25%)	Age ≥30: 0/2 Age <30: 0/1	Age ≥30: 3/5 (60%)	Age ≥30: 1/1 (100%)

This table only includes families where a proband had imaging (complete US, CT, or MRI) above the age of 20.