

Supplementary Table 3 – *CNGA1* gene variant identified in this study (HGVS nomenclature and variant classification applying ACMG Criteria)

Chromosome	Position (hg38)	Ref Seq	Alt Seq	Gene	RS ID	HGVS Nomenclature
chr4	47936957	C	T	CNGA1	rs544588016	NM_001379270.1:c.1525G>A (p.Gly509Arg)
ACMG Criteria	Justification					
PS3 Strong	Well established in vitro functional studies (Jin et al., 2016- PMID: 26802146).					
PP3 Supporting	Pathogenic computational verdict based on 12 pathogenic predictions from BayesDel_addAF (Score - 0.4308), CADD (Score -26.1), DEOGEN2 (Score - 0.8299), EIGEN (Score -1.0079), FATHMM-MKL (Score -0.9805), LIST-S2 (Score -0.996), M-CAP (Score -0.7232), MVP (Score -0.9722), MutationAssessor (Score -4.1), MutationTaster (Score -1), PrimateAI (Score - 0.9237) and SIFT (Score -0) vs no benign predictions. – Meta predictors – All Meta predictors predicted this variant as pathogenic -REVEL (Score- 0.93); MetaLR (Score-0.9481); MetaSVM (Score-1.1063); MetaRNN (Score- 0.9325).					
PM2 Moderate	Variant is present at extremely low frequency in the control population – gnomAD exomes - Allele frequency - 0.000008029 (2/249086) – No homozygotes – Present in only Asian population (East Asian – 1 in 17974; South Asian – 1 in 30588); gnomAD genomes (Allele frequency - 0.00001972 (3/152130)) {East Asians, 1 in 5194 alleles (0.0001925); European (Non Finnish, 1 in 68,040 (0.00001470); African/African American, 1 in 41,400 (0.00002415)}					
PP1 Supporting	We have found this variant to co-segregate in autosomal recessive pattern (homozygous) among individuals with RP in our family. Furthermore, the same variant reported to co-segregate in a Chinese family as compound heterozygous mutation along with another variation.					
Classification	Pathogenic (PS3 Strong + PM2 Moderate + PP1 Supporting + PP3 Supporting)					