

Abstract

Title: “Knowledge practice gap of nurses towards COVID-19 patients’ dead body care in a tertiary care hospital”.

Authors: Girraj Saini, Mahendra Singh, Prasan Kumar Panda, Manish Kumar Sharma, Pankaj Punjot, Raviprakash Meshram, Puneet Kumar Gupta.

Aim: To know the dead body care of COVID -19 patients.

Objective: 1.To determine health care professionals, knowledge, attitude, and practice towards Covid-19 dead body care.

2. To find the association of knowledge, attitude, practice with selected demographic variables.

Background: COVID -19 was a global pandemic and it was a serious note for health care professionals from many aspects. The virus was infective and causes serious infectionsto patients which were easily transmitted, hence specific dead body care is required for such kinds of patients. To keep this background in mind the study was conducted to identify the knowledge, practice and attitude towards COVID-19 dead body care among nurses.

Methodology: A cross sectional survey based study was done on 282 samples.Quantitative research design with purposive sampling technique data was collected for knowledge,attitude, and practice.

Result: Knowledge, attitude and practice were assessed and association was done with demographic profile. Hence the good knowledge, attitude and practicewere observed in experienced and trained nurses (p value<0.005. Whereas no significant changes were observed with age, gender and education qualification.

Conclusion: Overall knowledge, attitude and practice regarding COVID -19 dead body care were moderate to good. But it was important to identify the gap as it was a global pandemic and higher chances of spreading of infection.

Keywords: Contact transmission, Corona virus disease 2019, Dead body, Healthcare workers, and Questionnaire study.

Title: “Knowledge practice gap of nurses towards COVID-19 patients’ dead body care in a tertiary care hospital”.

Short title: COVID dead body care among nursing professionals.

Authors: Girraj Saini, Mahendra Singh, Prasan Kumar Panda, Manish Kumar Sharma, Pankaj Punjot, RaviprakashMeshram, Puneet Kumar Gupta

Affiliations:All India Institute of Medical Sciences (AIIMS), Rishikesh, 249203, India

Introduction:

The COVID pandemic is an ongoing global pandemic caused by SARS-COV-19, which is first identified during an outbreak in Wuhan, China. The 1st case in India is reported in Thrissur district in Kerala followed by arising number of cases, as of the 28th of May 2022, there had been 43147530 confirmed cases and 524539 deaths (1-2). There are still some uncertainties in the natural history of COVID including source, transmissibility mechanisms, viral shedding and persistency of the virus in the environment. In health care settings, aerosol generating procedures, as well as the surfaces, fomites, and contaminated hands of healthcare workers (HCWs) have a role in the spread of disease. Since it is unclear how long the virus stays in the victim’s body, and the absence of data on infection from dead bodies, the WHO recommends the safe handling of bodies to prevent the possible spread of infection(ref). Therefore, the safety and well- being of persons handling the dead bodies should be the priority: it includes the use of personal protective equipment (PPE) while safe handling the bodies, placement of dead body in an impermeable bag to prevent leakage of body fluids before being moved from the isolation area or patient care room. (3-5)

For aerosol generating procedures for all confirmed or suspected COVID patients, all HCWs should follow the standard, contact, droplet and airborne precautions. All the persons including nursing staff handling the victims’ bodies in the isolation area, mortuary, ambulance, during crematorium, and burial ground should be trained in infection prevention and control practices. During hospital role of nursing staff remains crucial for dead body handling. Thus, this study aims to describe nurses knowledge, attitude, and practice gaps concerning COVID victim’s body care and factors that affect the above- mentioned precautions. So that the proper strategies to minimize the risk of infection in individuals handling the bodies of deceased persons infected with COVID are identified as the need of

the hour. The information resulted from this study can serve as a vital tool to formulate relevant policies and guidelines during and after the outbreak and guide the nurses to prioritize self-protection and avoid occupational exposure.(6-8)

Material and Methods:

A single-centre cross-sectional survey-based study was conducted in the tertiary health care centre. To implement social distancing to avoid the spread of the COVID, it was not feasible to collect self-reported data-based surveys therefore the investigators used an online method of data collection. The sample size calculated by investigators as all health care workers posted in COVID units during data collection from July 2020 to September 2020. The survey commenced with 282 nurses posted in COVID units, and the required sample size was achieved on 20th September 2020. All Nursing officers working in COVID units were recruited as study participants in the present study. A questionnaire was designed on Google forms, and a link was shared with WhatsApp groups and personal messages of COVID unit staff.

Measurement and Data Collection:

The structured questionnaire was prepared after reviewing published literature based on WHO guidelines, institutional infection control protocols, and course material regarding emerging respiratory diseases including COVID(5–8). An initial draft of the questionnaire was designed and subsequently validated. The study instrument was sent to experts from the field of medicine, microbiology, nursing and toxicology, which was requested to give their expert opinion concerning its simplicity, relativity and importance. The questionnaire contains four parts socio-demographics, knowledge, attitude, and practice related items.

Table 1: Criteria of score range with cut-off level.

S.N.	Criteria	Score Range	Cut off level	
1	Knowledge	0-8	Good Knowledge	Poor knowledge
			≥5	□ 5
2	Practice	0-17	Good Practice	Poor Practice
			≥11	□ 11
3	Attitude	13-80	Positive Attitude	Negative Attitude
			≥48	□ 48

Ethical considerations:

Ethical clearance was obtained from the institute ethical committee vide letter no AIIMS/IEC/20/558 Dated 22/08/2020 before conducting the study. The study questionnaire contained a consent section that stated the purpose of the study, nature of the survey, study objectives, voluntary participation, declaration of confidentiality, and anonymity.

Result:

Response data of knowledge, attitude, and practices regarding COVID dead body care was collected from the 282 nurses. Demographic profile of study participants along with its association with knowledge is mentioned in Table 2. The majority of nurses were having work experience in COVID units. Whereas less the work experience shows significant low in knowledge ($P=0.0055$) for 0-30 days.

Table 2: Association between demographic profile with Knowledge of Nurses (n=282)							
Variable	Good	%	Poor	%	Total	OR (95% CI)	p-Value
Age							
21-30 yr	181	71.3	73	28.7	254	1	0.44
>30 yr	18	64.3	10	35.7	28	1.37(0.60-3.12)	
Gender							
Male	137	71.7	54	28.3	191	1	
Female	62	68.1	29	31.9	91	1.18(0.69-2.04)	0.53
Years of experience in current profession							
<2 yr	84	68.9	38	31.1	122	1.18(0.70-1.99)	0.51
2-5 yr	113	72.4	43	27.6	156	1	
>5 yr	2	50	2	50	04	2.62(0.35-19.2)	0.34
Work experience in COVID designated area							
0-30 days	24	52.2	22	47.8	46	2.65(1.33-5.27)	0.0055
31-60 days	48	73.8	17	26.2	65	1.02(0.52-1.99)	0.94
61-80 days	20	74.1	7	25.9	27	1.01(0.39-2.58)	0.97
>80 days	107	74.3	37	25.7	144	1	
Educational qualification							
General Nursing Midwifery	28	58.3	20	41.7	48	1.93(1.01-3.68)	0.43
BSc/Msc Nursing	171		63		234	1	
Experience of handling dead body of COVID patient							
Yes	151	83	31	17	182	1	
No	48	48	52	52	100	5.27(3.04-9.15)	<0.001
Experience of handling dead body of SARS, MERSA, Swine Flu, NIPAH, EBOLA							
Yes	104	86.7	16	13.3	120	1	
No	95	58.6	67	41.4	162	4.58(2.48-8.45)	<0.001
Any training							
Yes	119	85	21	15	140	1	

No	80	56.3	62	43.7	142	4.39(2.48-7.76)	<0.001
----	----	------	----	------	-----	-----------------	--------

The attitude gap was more significant in work experience in COVID designated area, educational qualification, the experience of handling dead bodies of COVID patients, and handling of other infections including the training (P<0.05) (Table 3). While age, gender, and experience of the current profession were not showing any differences in attitude.

Variable	Positive	%	Negative	%	Total	OR (95% CI)	p-Value
Age							
21-30 yr	80	31.5	174	68.5	254	1.88(0.85-4.14)	0.115
>30 yr	13	46.4	15	53.6	28	1	
Gender							
Male	70	36.6	121	63.4	191	1	
Female	23	25.3	68	74.7	91	1.71(0.98-2.98)	0.058
Years of experience in current profession							
<2 yr	33	27	89	73	122	8.09(0.81-80.5)	0.074
2-5 yr	57	36.5	99	63.5	156	5.21(0.52-51.2)	0.157
>5 yr	3	75	1	4	4	1	
Work experience in COVID designated area							
0-30 days	18	39.1	28	60.9	46	1	
31-60 days	19	29.2	46	70.8	65	1.55(0.70-3.45)	0.277
61-80 days	10	37	17	63	27	1.09(0.41-2.91)	0.859
>80 days	46	31.9	98	68.1	144	0.067(0.03-0.14)	<0.0001
Educational qualification							
General Nursing Midwifery	23	47.9	25	52.1	48	1	
BSc Nursing	70		164		234	2.155(1.14-4.05)	0.017
Experience of handling dead body of COVID patient							
Yes	46	25.3	136	74.7	182	2.62(1.56-4.39)	0.0002
No	47	47	53	53	100	1	
Experience of handling dead body of SARS, MERSA, Swine Flu, NIPAH, EBOLA							
Yes	22	18.3	98	81.7	120	3.47(1.99-6.06)	<0.0001

No	71	43.8	91	56.2	162	1	
Any training							
Yes	36	25.7	104	74.3	140	1.93(1.16-3.21)	0.010
No	57	40.1	85	59.9	142	1	

The practice of male nurses were having good practice compared to females (P=0.009) (Table 4). Age, education, years of nursing experience, dedicated COVID training, and experience of handling other infections had no significant practice gaps in dead body care. Work experiences in the COVID unit and experience of handling the COVID dead body had significant practice gaps like more experiences had good practices (P<0.005).

Table 4: Association between demographic profile with Practice of Nurses (n=282)							
Variable	Good	%	Poor	%	Total	OR (95% CI)	p-Value
Age							
21-30 yr	230	90.6	24	9.4	254	2.81(0.366-21.6)	0.31
>30 yr	27	96.4	1	3.6	28	1	
Gender							
Male	182	95.3	9	4.7	191	1	
Female	75	82.4	16	17.6	91	4.31(1.82-10.19)	0.0009
Years of experience in current profession							
<2 yr	111	91	11	9	122	0.92(0.04-18.35)	0.96
2-5 yr	142	91	14	9	156	0.91(0.04-17.87)	0.95
>5 yr	4	100	0	0	4	1	
Work experience in COVID designated area							
0-30 days	43	93.5	3	6.5	46	1.93(0.44-8.44)	0.377
31-60 days	54	83.1	11	16.9	65	5.66(1.87-17.06)	0.0021
61-80 days	21	77.8	6	22.2	27	7.94(2.22-28.35)	0.0014
>80 days	139	96.5	5	3.5	144	1	
Educational qualification							
General Nursing Midwifery	40	83.3	8	16.7	48	2.55(1.03-6.31)	0.42
BSc Nursing	217		17		234	1	
Experience of handling dead body of COVID patient							
Yes	176	96.7	6	3.3	182	1	
No	81	81	19	19	100	6.88(2.64-17.87)	0.0001
Experience of handling dead body of SARS, MERSA, Swine Flu, NIPAH, EBOLA							
Yes	114	95	6	5	120	1	
No	143	88.3	19	11.7	162	2.52(0.97-6.52)	0.56
Any training							
Yes	131	93.6	9	6.4	140	1	
No	126	88.7	16	11.3	142	1.84(0.78-4.33)	0.157

Discussion:

The current study examined the knowledge, practice, and attitude of nurses regarding COVID dead body care. Dead body care is an essential part of end-of-life care. The care after death especially during the major pandemic of COVID highlights many aspects for health care professionals to take care. In this study it is found that work experience in the COVID unit has a significant impact on knowledge-practice regarding COVID dead body care. Similarly, it is observed that training has improved the knowledge-practice of nursing personnel regarding dead body care. It is the first study that describes these observational findings on caring the COVID dead body by nurses.

The findings are strongly suggestive of good knowledge in those who worked in COVID areas. Many studies also suggested that effective and appropriate health education and training programme improve COVID knowledge and safe practices (8,9). COVID infection leads to many super-added infections and other health problems. Those who practice how to handle other infection like MERSA, SARS, NIPAH, etc are supposed to be well trained in handling a variety of infections (10). Similar findings were observed in this study that those who handle other infection had higher aspects of knowledge, attitude, and practice.

It is well established fact that training improves knowledge, practice, and attitude (11,12). The same observations were observed among nurses while handling COVID dead body care except for practices. Nurses were more in exposure as they were direct in care providing to this global pandemic. Preparation for the response is a vital step to cope effectively and then deal with actual health emergencies in the case of developing countries like India. Readiness for combating infectious diseases such as COVID begins with good understanding, positive thinking and safe and better practices. Similarly, in the study also attitude, and practice were assessed regarding handling of dead body care which gives direct exposure to the same (13,14). The dead body was considered infected and caring for a dead body with COVID infection for nurses requires a positive attitude with good knowledge and practical skills (15). The care during this war and specific protection with preventive aspects for the health care providers has a primary role in the reduction of the cases.

Limitation:

- a) The study was conducted in one institute and can be conducted with different institutions.
- b) There was no specific training provided regarding COVID- specific dead body care.
- c) Study can be performed on a larger sample size.

Conclusion

Overall knowledge, attitude, and practice regarding COVID dead body care among nurses were moderate to good. As it is pandemic and has direct exposure to health care provider it was necessary to identify the gaps in the knowledge-practice of nurses regarding this dead body care.

Reference:

1. Haque T, Hossain KM, Bhuiyan MM, Ananna SA, Chowdhury SH, Islam MR, Ahmed A, Rahman MM. Knowledge, attitude and practices (KAP) towards COVID-19 and assessment of risks of infection by SARS-CoV-2 among the Bangladeshi population: An online cross sectional survey.
2. Maheshwari S, Gupta PK, Sinha R, Rawat P. Knowledge, attitude, and practice towards coronavirus disease 2019 (COVID-19) among medical students: A cross-sectional study. *Journal of Acute Disease*. 2020 May 1;9(3):100.
3. Hussain A, Garima T, Singh BM, Ram R, Tripti RP. Knowledge, attitudes, and practices towards COVID-19 among Nepalese Residents: A quick online cross-sectional survey. *Asian Journal of Medical Sciences*. 2020 May 1;11(3):6-11.
4. Sharma M, Sharma D, Sharma AK, Mohanty A, Khapre M, Kalyani CV. Barriers faced by health-care workers in use of personal protective equipment during COVID pandemic at tertiary care hospital Uttarakhand, India: A qualitative study. *J Educ Health Promot*. 2022;11:74.
5. Patidar K, Sharma M, Gautam A, Sharma DK, Jain J. COVID-19 knowledge and perception among budding nurses: A questionnaire-based survey. *Int J Nurs Res*. 2020;6(2):59–65.
6. Padmanaban S, Rajendran P, Davis P, Velayutham P. Knowledge, attitude and practices towards COVID-19 among higher education students in India: a cross sectional study. *Z GesundhWiss*. 2021;1–13.

7. WH. Infection prevention and control for the safe management of a dead body in the context of COVID-19: interim guidance [Internet]. Who.int. World Health Organization; 2020 [cited 2022 Jun 6]. Available from: <https://www.who.int/publications/i/item/infection-prevention-and-control-for-the-safe-management-of-a-dead-body-in-the-context-of-covid-19-interim-guidance>.
8. Wang C, Horby PW, Hayden FG, Gao GF. A novel coronavirus outbreak of global health concern. *Lancet* [Internet]. 2020;395(10223):470–3. Available from: [http://dx.doi.org/10.1016/S0140-6736\(20\)30185-9](http://dx.doi.org/10.1016/S0140-6736(20)30185-9).
9. Saha AK, Mitra CR, Khatun RA, Reza HM. Nurses’ knowledge and practices regarding prevention and control of COVID-19 infection in a tertiary level hospital. *Banglad J Infect Dis* [Internet]. 2020;S27–33. Available from: <http://dx.doi.org/10.3329/bjid.v7i00.50159>.
10. WHO. A Guide to Developing Knowledge, Attitude and Practice Surveys 2008
11. Reda AA, Frisseha S, Mengistie B, Vanderweerd J. Standard precautions: occupational exposure and behavior of health care workers in Ethiopia. *Plos One* 2010;5 (12):e14420.
12. Kafle A, Pandit Pahari S, Khanal S, Baral K, Pathak K, Baral S, et al. Knowledge regarding COVID-19 among registered nurses of Pokhara, Nepal. *Europasian J Med Sci* [Internet]. 2020;2(2):6–12. Available from: <http://dx.doi.org/10.46405/ejms.v2i2.88>
13. Bhagavathula AS, Aldhalei WA, Rahmani J, Mahabadi MA, Bandari DK. Knowledge and perceptions of COVID-19 among health care workers: Cross-sectional study. *JMIR Public Health Surveill* [Internet]. 2020;6(2):e19160. Available from: <http://dx.doi.org/10.2196/19160>.
14. Lee VJ, Ho M, Kai CW, Aguilera X, Heymann D, Wilder-Smith A. Epidemic preparedness in urban settings: new challenges and opportunities. *Lancet Infect Dis* [Internet]. 2020;20(5):527–9. Available from: [http://dx.doi.org/10.1016/s1473-3099\(20\)30249-8](http://dx.doi.org/10.1016/s1473-3099(20)30249-8).
15. Gulilat K. Assessment of knowledge, attitude and practice of health care workers on infection prevention in health institution Bahir Dar city administration. *Sci J Publ Health* [Internet]. 2014;2(5):384. Available from: <http://dx.doi.org/10.11648/j.sjph.20140205.13>