

**Supplementary Table S1. The demographic and clinical characteristics of patients after propensity score-matching**

Characteristics	Fish oil user	Fish oil non-user	Overall	SMD
<b>Number of participants, n(%)</b>	29 424	29 424	58 848	
<b>Age, mean(SD), years</b>	58.1 (7.0)	58.4 (7.4)	58.2 (7.2)	0.035
<b>Sex, female, n(%)</b>	16 589 (56.4)	16 109 (54.7)	32 698 (55.6)	0.033
<b>Ethnicity, White, n(%)</b>	28 710 (97.6)	28 751 (97.7)	57 461 (97.6)	0.009
<b>Household income (£)</b>				0.072
<18 000, n(%)	4 389 (14.9)	4 553 (15.5)	8 942 (15.2)	
18 000-30 999, n(%)	7 061 (24.0)	6 659 (22.6)	13 720 (23.3)	
31 000-51 999, n(%)	7 479 (25.4)	7 107 (24.2)	14 586 (24.8)	
52 000-100 000, n(%)	5 902 (20.1)	5 796 (19.7)	11 698 (19.9)	
>100 000, n(%)	1 478 (5.0)	1 695 (5.8)	3 173 (5.4)	
Missing, n(%)	3 115 (10.6)	3 614 (12.3)	6 729 (11.4)	
<b>Deprivation index, mean(SD)</b>	-1.93 (2.7)	-1.96 (2.7)	-1.94 (2.7)	0.013
<b>BMI, mean(SD), kg/m<sup>2</sup></b>	26.58 (4.3)	26.58 (4.3)	26.58 (4.3)	0.002
<b>Alcohol consumption</b>				0.049
Daily or almost daily, n(%)	7 033 (23.9)	7 501 (25.5)	14 534 (24.7)	
Three or four times a week, n(%)	7 935 (27.0)	7 716 (26.2)	15 651 (26.6)	
Once or twice a week, n(%)	7 406 (25.2)	7 171 (24.4)	14 577 (24.8)	
One to three times a month, n(%)	2 986 (10.1)	2 814 (9.6)	5 800 (9.9)	
Special occasions only or never, n(%)	2 544 (8.6)	2 543 (8.6)	5 087 (8.6)	
Never, n(%)	1 520 (5.2)	1 679 (5.7)	3 199 (5.4)	
<b>Smoking status</b>				0.036
Never smoker, n(%)	16 289 (55.4)	16 401 (55.7)	32 690 (55.5)	
Previous smoker, n(%)	11 311 (38.4)	10 974 (37.3)	22 285 (37.9)	
Current smoker, n(%)	1 824 (6.2)	2 049 (7.0)	3 873 (6.6)	
<b>Physical activity, mean(SD), MET minutes/week</b>	2 732 (2 561)	2 773 (2 750)	2 753 (2 657)	0.016
<b>Fresh fruit, mean(SD), pieces/day</b>	2.3 (2.3)	2.3 (2.2)	2.3 (2.2)	0.009
<b>Raw vegetable, mean(SD), tablespoons/day</b>	1.89 (3.2)	1.86 (3.2)	1.87 (3.2)	0.011
<b>Oily fish consumption (times/week)</b>				0.004
<2, n(%)	23 090 (78.5)	23 144 (78.7)	46 234 (78.6)	
≥2, n(%)	6 334 (21.5)	6 280 (21.3)	12 614 (21.4)	
<b>Non-oily consumption fish (times/week)</b>				0.003
<2, n(%)	24 473 (83.2)	24 504 (83.3)	48 977 (83.2)	
≥2, n(%)	4 951 (16.8)	4 920 (16.7)	9 871 (16.8)	
<b>Vitamin supplementation, n(%)</b>	11 031 (37.5)	7 853 (26.7)	18 884 (32.1)	0.233
<b>Mineral and other dietary supplementation, n(%)</b>	5 686 (19.3)	4 177 (14.2)	9 863 (16.8)	0.138
<b>Comorbidities</b>				
Hypertension, n(%)	7 162 (24.3)	7 448 (25.3)	14 610 (24.8)	0.023

Type 2 diabetes, n(%)	137 (0.5)	138 (0.5)	275 (0.5)	<0.001
Renal failure, n(%)	34 (0.1)	29 (0.1)	63 (0.1)	0.005
Myocardial infarction, n(%)	539 (1.8)	634 (2.2)	1 173 (2.0)	0.023
Stroke, n(%)	235 (0.8)	247 (0.8)	482 (0.8)	0.005
COPD, n(%)	54 (0.2)	56 (0.2)	110 (0.2)	0.002
Asthma, n(%)	3 148 (10.7)	3 162 (10.7)	6 310 (10.7)	0.002
Longstanding illness, n(%)	8 650 (29.4)	8 699 (29.6)	17 349 (29.5)	0.004

SMD: standard mean difference; BMI: body mass index; MET: metabolic equivalent of task;

COPD: chronic obstructive pulmonary disease; SD: standard deviation.

**Supplementary Table S2. Propensity score-matched analysis**

	Case/person-years	Unadjusted model		Multivariable-adjusted model*	
		HR (95%CI)	P	HR (95%CI)	P
<b>SARS-CoV-2 infection</b>					
Fish oil non-user	13 219/160 063	1.00(Reference)		1.00(Reference)	
Fish oil user					
Time <sup>#</sup> <12.1 years	6 111/69 718	0.96(0.93-0.99)	0.037	0.96(0.93-0.99)	<b>0.033</b>
Time≥12.1 years	6 851/87 844	0.99(0.96-1.02)	0.640	1.00(0.97-1.04)	0.856
<b>COVID-19 hospital admission</b>					
Fish oil non-user	743/8 687	1.00(Reference)		1.00(Reference)	
Fish oil user	590/6 904	0.76(0.68-0.85)	<b>&lt;0.001</b>	0.77(0.69-0.86)	<b>&lt;0.001</b>
<b>COVID-19 mortality</b>					
Fish oil non-user	242/2 830	1.00(Reference)		1.00(Reference)	
Fish oil user	172/2 030	0.68(0.56-0.83)	<b>&lt;0.001</b>	0.70(0.57-0.85)	<b>&lt;0.001</b>

\*Adjusted for vitamin supplementation, and mineral and other dietary supplementation

<sup>#</sup>follow-up time

HR: hazard ratio; CI: confidence interval; COVID-19: coronavirus disease 2019

**Supplementary Table S3. Odds ratios (95% CI) for the associations of use of fish oil with the risk of SARS-CoV-2 infection and COVID-19-related outcomes**

Case/person-years	Non-adjusted model		Age and sex-adjusted model		Fully adjusted model*	
	OR (95% CI)	P	OR (95% CI)	P	OR (95% CI)	P
<b>SARS-CoV-2 infection</b>						
Fish oil non-user	33 935/412 007	1.00 (Reference)		1.00 (Reference)		1.00 (Reference)
Fish oil user	12 962/157 562	1.09(1.06-1.12)	<0.001	0.97(0.93-0.99)	0.012	0.98(0.95-1.01)
<b>COVID-19 hospital admission</b>						
Fish oil non-user	1 743/20 422	1.00 (Reference)		1.00 (Reference)		1.00 (Reference)
Fish oil user	590/6 904	0.93(0.84-1.02)	0.137	0.77(0.70-0.85)	<0.001	0.81(0.73-0.90)
<b>COVID-19 mortality</b>						
Fish oil non-user	473/5 573	1.00 (Reference)		1.00 (Reference)		1.00 (Reference)
Fish oil user	172/2 030	1.00(0.84-1.20)	0.989	0.71(0.59-0.85)	<0.001	0.75(0.62-0.90)

\*Adjusted for age, sex, ethnicity (white, other), average total annual household income (<£18 000, £18 000-£30 999, £31 000-£51 999, £52 000-£100 000, >£100 000, and missing), Deprivation Index, body mass index, alcohol consumption(daily or almost daily, three or four times a week, once or twice a week, one to three times a month, special occasions only or never, never), smoking status(never, previous, current), physical activity, fresh fruit consumption, raw vegetable consumption, oily fish consumption (<2 or ≥2 times/week), non-oily fish consumption (<2 or ≥2 times/week), vitamin supplement use (yes or no), mineral and other dietary supplement use (yes or no), hypertension (yes or no), type 2 diabetes(yes or no), renal failure(yes or no), myocardial infarction (yes or no), stroke (yes or no), chronic obstructive pulmonary disease (yes or no), asthma (yes or no), longstanding illness (yes or no)

OR: Odds ratios; CI: confidence interval; COVID-19: coronavirus disease 2019