

Intraoperative phrenic stimulation offsets diaphragm fiber weakness during cardiothoracic surgery

Guilherme Bresciani ¹, Thomas Beaver ², A. Daniel Martin ³, Robbert van der Pijl ⁴, Robert Mankowski ⁵, Christiaan Leeuwenburgh ⁵, Coen A.C. Ottenheijm ⁴, Tomas Martin ², George Arnaoutakis ², Shakeel Ahmed ³, Vinicius Mariani,¹ Wei Xue,⁶ *Barbara K. Smith ³, *Leonardo F. Ferreira ¹

¹ Department of Applied Physiology and Kinesiology, University of Florida

² Department of Surgery, University of Florida

³ Department of Physical Therapy, University of Florida

⁴ Department of Cellular and Molecular Medicine, University of Arizona

⁵ Department of Aging and Geriatric Research, University of Florida

⁶ Department of Biostatistics, University of Florida

*equal contributors

Corresponding Author:

Leonardo F. Ferreira
University of Florida
1864 Stadium Rd, 100 FLG
Gainesville, FL 32611-8205
Phone: +1 (352) 294-1724
Email: ferreira@hhp.ufl.edu

Author Contributions:

Conception of the work – TB, CL, CACO, DM, LF, BS

Acquisition of data – GB, TB, RM, DM, TM, GA, SA, BS

Analysis and interpretation of the data – GB, TB, RV, CL, CACO, DM, LF, BS

Drafting the work – GB, TB, RV, CACO, LF, BS

Critical revisions of the work – RM, CL, DM, TM, GA, SA

Grant support: NIH R01 AR072328

ABSTRACT

Rationale:

Mechanical ventilation rapidly induces slow and fast fiber contractile dysfunction in the human diaphragm, which could be attenuated by phrenic nerve stimulation. Here, we present data from a controlled trial of intraoperative phrenic stimulation to offset slow and fast fiber contractile dysfunction and myofilament protein derangements.

Objectives:

In this study, we tested the hypothesis that intraoperative hemidiaphragm stimulation would mitigate slow and fast fiber loss of contractile function in the human diaphragm.

Methods:

Nineteen adults (9 females, age 59 ± 12 years) consented to participate. Unilateral phrenic twitch stimulation was applied for one minute, every 30 minutes during cardiothoracic surgery. Thirty minutes following the last stimulation bout, biopsies were obtained from the hemidiaphragms for single fiber force mechanics and quantitation of thin filament protein abundance. Effects of stimulation and fiber type on force mechanics were evaluated with linear mixed models with the subject treated as a random intercept effect.

Measurements and Main Results:

Subjects underwent 6 ± 2 hemidiaphragm stimulations at 17 ± 6 mA, during 278 ± 68 minutes of mechanical ventilation. In slow-twitch fibers, cross-sectional area ($p < 0.0001$) and specific force ($p < 0.0005$) were significantly greater on the stimulated side. Longer-duration surgeries were associated with lower slow-twitch specific force ($p < 0.001$). Stimulation did not alter contractile function of fast-twitch fibers or calcium-sensitivity in either fiber type. There were no differences in abundance or phosphorylation of myofilament proteins.

Conclusion:

Unilateral phrenic stimulation during open chest surgery preserved contractile function of slow-twitch diaphragm fibers, but had no effect on relative abundance of sarcomeric proteins.

Keywords: mechanical ventilation, cardiothoracic surgery, diaphragm, electrical stimulation

INTRODUCTION

Mechanical ventilation (MV) is life-saving treatment and required for open-chest surgeries. However, extended periods of MV can lead to ventilator-induced diaphragm dysfunction (VIDD), which increases chances of medical complications and has been recognized as a major clinical barrier to weaning among patients who require prolonged MV support (1). The signs of VIDD occur within hours of MV and include increased protein degradation and loss of force generation (2-6). The loss of force in VIDD occurs due to fiber atrophy and contractile dysfunction (4, 7, 8), with the latter being documented by a decrease in force divided per cross-sectional area (specific force). The exact mechanisms of VIDD, especially in humans, are unknown but appear to involve unloading of the mechanosensitive sarcomeric protein titin (9). Titin extends from the Z-disk to the M-band of the sarcomere and contributes to elastic properties of the sarcomere (10) . Unloading by MV induces passive shortening of the diaphragm (11), and changes in sarcomere length induce conformational changes in titin, which trigger signaling pathways involved in protein turnover (12, 13). Prolonged controlled MV leads to the degradation of multiple sarcomeric elements within diaphragm fibers, including the primary contractile protein, myosin heavy chain, as well as titin (14). Sarcomeric protein degradation is a putative mechanism for contractile dysfunction in VIDD (13).

Activity-based interventions, such as preoperative inspiratory muscle training, have been proposed as a method to maintain diaphragm loading and prevent VIDD (15). However, pre-MV interventions do not directly treat the inactivity that causes

diaphragm dysfunction *during* MV. Diaphragm activation during MV is possible via electrical stimulation of the phrenic nerve, which prevents diaphragm atrophy in animals (16, 17) and humans (18, 19), and it partially protects against contractile dysfunction in rodents during prolonged MV (16). We reported that intermittent intraoperative phrenic stimulation increased force of fast-twitch diaphragm fibers in a small pilot study with a cohort of patients undergoing open-chest surgery (20). Positive outcomes in pre-clinical studies and our earlier pilot findings led us to complete the current study to test the hypothesis that intermittent electrical phrenic stimulation during open-chest surgery protects from sarcomeric protein degradation and diaphragm fiber contractile dysfunction induced by MV.

METHODS

Patients

Eligibility criteria for intraoperative hemidiaphragm stimulation included adults (18-80 years) scheduled for non-emergent clinical cardiothoracic surgeries anticipated to last at least 3 hours, enabling 4 or more bouts of phrenic stimulation. Patients were ineligible to participate for any of the following: (1) history of surgery to the diaphragm or pleura, (2) severe obstructive lung disease (FEV1 <40% predicted), (3) restrictive lung disease, (4) severe chronic heart failure (NYHA class IV), (5) chronic kidney disease with serum creatinine >1.6 mg/dL, (6) body mass index <20 or >40 kg/m², (7) chronic uncontrolled metabolic disease, (8) concurrent neoplastic or myopathic illness, or (9) use of immunosuppressants, corticosteroids, or aminoglycoside antibiotics within 28 days of surgery. The protocol was approved by the University of Florida Institutional Review

Board and prospectively registered (NCT03303040). Patients were recruited from the University of Florida cardiothoracic surgery clinic, and each patient provided their written informed consent in advance of participation.

Pulmonary Function Testing

Subjects completed forced vital capacity (FVC) maneuvers to screen for significant obstructive disease. Seated spirometry (Discovery 2, FutureMed) was conducted in accordance with recommended guidelines (21). Additionally, maximal inspiratory pressure (P_Imax) was measured with a commercial manometer (MicroRPM, Vyair Medical) using recommended procedures (22). For both tests, trials were repeated 3-5 times, and the highest effort was reported.

Intraoperative Phrenic Stimulation

All subjects underwent a midline sternotomy. During their surgery, the right or left phrenic nerve was stimulated using an external cardiac pacemaker (Model #5388, Medtronic) and temporary pacing wires. Selection of the stimulated side was determined by the surgeon, based on ease of access to the corresponding phrenic nerve and the type of procedure being performed. The surgeon placed electrodes adjacent (~5-10 mm) to the phrenic nerve, either near the pulmonary artery or at the right pericardium, and confirmed the stimulating electrodes did not directly contact the phrenic nerve. Twitch stimulation parameters included a 1.5 ms pulse width, frequency and 1-minute duration. Atrial output was 0 mA, and ventricular output was set at twice the twitch threshold, which ensured a vigorous hemidiaphragm contraction. Stimulations

continued every 30 min from the time the phrenic nerves were initially exposed until completion of the procedure.

Diaphragm Biopsies

Approximately 30 minutes following the final stimulation bout, diaphragm biopsies (~150 mg) were obtained from the stimulated and unstimulated costal hemidiaphragms and processed as described in Supplemental Methods.

Statistical analyses

Demographic characteristics of the patients were summarized using n (%) and Mean \pm SD. Single fiber outcomes of interest (area, abs force, passive tension, maximal specific force, Ktr, H and KD) were summarized with Mean, SD, and range. Then we summarized the average measures by side and fiber type. The individual fiber data were used to compare effects of stimulation on contractile function via a linear mixed model (SAS 9.4). Relationships between fiber CSA, specific force, and duration of MV were evaluated with linear regression. The abundance of proteins was tested for normality and compared with a Wilcoxon test.

RESULTS

Patient Characteristics

Figure 1 outlines the subject enrollment information. Of 1,018 surgical clinic patients reviewed for eligibility, 25 subjects were enrolled. Three subjects were enrolled but did not receive intraoperative stimulation, due to preoperative changes in their clinical care

plan. One subject initiated but did not complete intraoperative stimulation due to the requirement of high-dose neuromuscular blockade as part of surgical care. Of 21 patients who completed intraoperative hemidiaphragm stimulation, single fiber mechanics were not evaluated in two patients, due to unavailability of study personnel (n=1), and secondary to non-cardiac surgery (n=1). **Table 1** illustrates the demographic characteristics of the sample (n=19, 9 female). Patients underwent an average of 6 ± 2 intraoperative stimulations with stimulation amplitude of 17 ± 6 mA, and they were mechanically ventilated for 284 ± 62 minutes prior to biopsy. Of the 38 biopsies obtained for this study, two specimens (subject 15: unstimulated hemidiaphragm; subject 17: stimulated hemidiaphragm) contained primarily fat and connective tissue.

Diaphragm Single Fiber

In total, 745 diaphragm fibers were activated. Of those, 382 were classified as slow (186 stimulated and 196 unstimulated) and 363 fibers were classified as fast (186 stimulated and 177 unstimulated). **Figures 2, and S2 – S4** show the effects of stimulation on single fiber size and contractile function. In slow fibers, absolute force (**Figure 2A**), cross-sectional area (**Figure 2B**), and specific force (**Figure 2C**) were significantly higher (10-25%) for the stimulated hemidiaphragm (all $p < 0.005$). **Figure 2D** shows specific force vs pCa relationship for data averaged per subject. **Figure S2** shows that there were no significant differences in calcium sensitivity (pCa50), rate-constant of force redevelopment (k_{tr} , cross-bridge kinetics), and Hill coefficient (cooperativity). In fast fibers, we did not detect significant differences between stimulated and non-stimulated hemidiaphragms for any of the variables examined

(**Figure 2E-H and S1**). We also examined the relationship between fiber area \times force (**Figure S3**) and MV duration \times specific force (**Figure S4**) in stimulated and non-stimulated hemidiaphragms. Fiber force was directly related to estimated cross-sectional area in slow ($p < 0.001$) and fast fibers ($p < 0.001$) as expected, with a steeper slope for stimulated slow fibers ($p = 0.001$) that is consistent with higher specific force (**Figure S3**). Averaged specific force was inversely related to MV duration in unstimulated slow fibers (unstimulated fibers: $p < 0.005$, stimulated fibers: $p = 0.153$), further supporting protection of VIDD with stimulation (**Figure S4**). There was no significant relationship between force and MV duration in fast fibers (stimulated fibers: $p = 0.239$, unstimulated fibers: $p = 0.649$).

Protein abundance and post-translational modifications

There were no significant differences between sides in the relative abundance of the key structural, regulatory, and contractile proteins titin, nebulin, MyBP-C, MyHC, or actin. Similarly, no significant differences between stimulated and non-stimulated were evident in the abundance of titin or its degradation products (**Figure 3**), titin phosphorylation status (**Figure 4**), or abundance of titin-associated proteins (**Figure 3**). Total phosphorylation levels of nebulin and myosin binding protein C were also not different between sides (**Figure 4**). The protein calpain (CAPN3), which interacts with titin and is involved in sarcomere and cytoskeletal remodeling, has been implicated in VIDD. We measured total CAPN3 and CAPN3 degradation products (markers of activation) and found no difference between sides (**Figure 3B**). Similarly, the levels of the stress response proteins vinculin and CSRP3 (**Figure 3C**) were similar for each

side. Altogether the analyses of myofibrillar proteins, phosphorylation, and titin-linked or stress-responsive proteins revealed no differences between stimulated and unstimulated diaphragm.

DISCUSSION

The primary novel findings of our study were that intermittent phrenic stimulation during 4-6 hours of cardiothoracic surgery resulted in higher hemidiaphragm fiber forces and estimated cross-sectional area in slow fibers. We also found an inverse relationship between non-stimulated slow fiber contractile function and the duration of MV. Fast fibers were not affected by stimulation. The higher force in slow fibers was not accompanied by changes in abundance and phosphorylation status of myofibrillar proteins or titin-linked proteins measured in whole-tissue homogenates.

Clinical implications and current therapies for VIDD.

Findings from this study may have implications for clinical management of patients recovering from cardiothoracic surgery. Sonographic evidence of poor pre-operative diaphragm function is independently associated with post-operative pulmonary complications and increased ICU length of stay (23). Additionally, serial ultrasounds identified post-operative diaphragm dysfunction in 38 of 100 patients, following elective cardiac surgeries (24). While no preoperative patient attributes distinguished those who eventually developed diaphragm dysfunction, average cardiopulmonary bypass time was 87% longer in patients with postoperative diaphragm dysfunction. Post-operative diaphragm dysfunction resulted in more frequent pulmonary complications, leading to slower weaning from MV and longer ICU recovery time (24, 25). A recent large

multinational study found that 55% of patients who underwent cardiac surgery with cardiopulmonary bypass experienced at least one post-operative pulmonary complication (26). While several risk factors were identified, included among them were intraoperative PEEP and neuromuscular blockade, which can worsen existing diaphragm dysfunction (27, 28). While our findings suggest potential to attenuate diaphragm fiber contractile dysfunction with intraoperative phrenic stimulation during controlled MV, further studies are needed on its postoperative functional impact.

Impact of mechanical ventilation on diaphragm fiber size and contractile function. The consensus from animals and human studies is that MV causes diaphragm atrophy and impairs contractile function (4, 7, 29, 30). However, a previous study showed that MV, in the absence of critical illness, did not impair contractile function of permeabilized single fibers (31). While we must be careful with direct comparisons between studies, these findings could suggest that the loss of pressure generation capacity in humans (32, 33) and contractile function in intact diaphragm bundles (4, 34, 35) after MV result from impairments in neuromuscular transmission, membrane depolarization, and calcium handling. Our study was not designed to establish the effects of MV alone on contractile function. Nonetheless, we found an inverse relationship between slow fiber specific force and time from onset of MV to biopsy. This novel finding suggests that MV causes a progressive decline in sarcomeric protein function in diaphragm fibers of patients who are not critically ill. The contrasting findings between the current and previous study (31) may arise from different modes of ventilation, or exacerbation of VIDD during open-chest surgery (36). During open-chest surgery, the diaphragm is not subjected to the

same mechanical forces as when MV occurs with a closed chest (e.g. shortening due to PEEP and cyclic stretches). The surgical procedures performed in our study also required cardiopulmonary bypass, which can further impair diaphragm contractile function (37). Further, protective ventilation during surgery has gained recent consideration (38), but MV modes used during surgery often differ from those used to protect against VIDD in critical illness (39). Overall, studies in humans and animal models support the notion that the diaphragm undergoes rapid (within hours) atrophy and loss of contractile function and, based on our data, open-chest surgery with cardiopulmonary bypass appears to exacerbate these processes. Thus, VIDD may delay weaning of MV and prolong hospital stay not only in patients with critical illness but also those recovering from cardiothoracic surgeries (24).

Phrenic stimulation during mechanical ventilation. Recent studies suggest stimulation is a feasible, safe, and effective approach to enhance inspiratory pressure production and promote ventilator weaning in patients unable to complete voluntary inspiratory strengthening exercises (40). Studies in animal models and humans have shown that phrenic stimulation during MV prevented or attenuated the decrease in diaphragm thickness, fiber cross-sectional area, and contractile function (16, 17, 19, 41, 42). In comparison to our trial (1Hz stimulation for 60 seconds, every 30 minutes), these previous studies examined longer periods of MV (rats: 18 hours (16), pigs: 19-35 hours (17), humans: 48 to 60 hours (19, 41), sheep: 72 hours (42)) and employed protocols with higher stimulus frequency (20-30 Hz) and train rates (every 20 seconds in rats (16), from every breath to every fourth breath in humans (19, 41), and every breath in sheep

(42)). Rat diaphragm specific force assessed in 'intact' bundles ex vivo was preserved when MV included phrenic stimulation, but fiber cross-sectional area was not examined (16). In comparison, we observed a higher slow fiber maximal specific force and estimated fiber area in permeabilized fibers in the current study.

The relationship between MV duration and slow fiber specific force suggests that phrenic stimulation did not elicit full protection of sarcomeric protein function. Thus, the greater protection of contractile function in intact rodent diaphragm with phrenic stimulation is most likely due to combined improvements in excitation-contraction coupling and sarcomeric protein function.

Phrenic stimulation during MV in sheep attenuated the decrease in fast fiber cross-sectional area (42). In the current study, slow, but not fast fiber cross-sectional area was greater with stimulation. This discrepancy may be related to differences in stimulation protocols, duration of MV, or species. In pigs, higher diaphragm thickness and fiber cross-sectional area were not accompanied by improved inspiratory pressure (17), and there was no functional assessment reported in humans with higher diaphragm thickness after stimulation during MV (19). Since higher diaphragm thickness did not translate into better function in pigs (17), it is possible that, in larger mammals, phrenic stimulation during MV increases diaphragm thickness partially due to edema, and the higher cross-sectional area may occur through mechanisms that do not preserve contractile function. However, we found that higher slow fiber cross-sectional area in the stimulated hemidiaphragm accompanied proportional increases in maximal fiber force (**Figure S4**). The steeper slope for slow fiber area-to-maximal force relationship in the stimulated hemidiaphragm also reflects the improvement in maximal

specific force (**Figure 2**). The slope for fast fibers was also slightly steeper, but not statistically significant ($P = 0.054$), in the stimulated diaphragm. In general, our study reinforces the safety and feasibility of phrenic stimulation during MV and advances the field by showing intraoperative feasibility and direct benefits to diaphragm fibers.

Potential mechanisms of VIDD and (partial) protection with phrenic stimulation. MV results in diminished or abolished drive to breathe with a potential for phrenic and neuromuscular transmission dysfunction (43). Unloading the diaphragm triggers disuse atrophy and contractile dysfunction via abnormal proteostasis (increased protein degradation and decreased synthesis), impaired calcium release by the ryanodine receptor 1 (6, 44), and post-translational modifications of sarcomeric proteins (45). Titin-based mechanosensing has been proposed as a signaling hub for diaphragm dysfunction induced by unloading (9). Titin is a critical determinant of myofiber tension at optimal sarcomere length, and phrenic stimulation resulted in higher passive tension. However, there was no detectable effect of phrenic stimulation on titin abundance, degradation products, or titin-binding proteins that modulate stiffness (MARF1 and 2) in whole-tissue homogenates. The greater increase in slow fiber absolute force than specific force suggests that improvements elicited by phrenic stimulation was in proteostasis. Calpains have been implicated in protein degradation, atrophy, and contractile dysfunction induced by MV (46). Yet, we saw no effect of phrenic stimulation on calpain 3 abundance or degradation products that serve as markers of activation. The exact mechanisms underlying sarcomeric protein dysfunction and loss of specific force in VIDD remain unclear (47). The increase in maximal specific force without

changes in calcium sensitivity suggests improvements in function of thick filament proteins. We assessed key thick-filament proteins and found no difference in their abundance or phosphorylation status.

Skeletal muscle unloading causes cross-bridges to enter a super-relaxed state that prevents interaction with actin and force generation (48). One plausible scenario is that phrenic stimulation, by imposing intermittent load on the diaphragm during MV, mitigates the loss of specific force by minimizing the transition of cross-bridges to a super-relaxed state. We did not prepare the samples to assess cross-bridge states but this topic is worth investigating in future studies. Overall, we were unable to define potential molecular mechanisms associated with higher fiber area and force induced by phrenic stimulation in whole tissue homogenates. Fiber type-specific post-translational modification proteomics and synchrotron radiation x-ray diffraction will be necessary to resolve potential biochemical and biophysical events underlying the improvements in fiber size and force with phrenic stimulation.

Limitations and methodological considerations. We stimulated at twice the minimal threshold (up to 25 mA twitches) to achieve supramaximal stimulation. The clinical environment did not permit us to make quantification of force generation, but we could clearly observe vigorous contractions of the stimulated side. The optimal parameters and timing of phrenic stimulation remains unresolved. Our goal was to test a protocol that was considered ‘minimally disruptive’ for the surgical procedures. However, the benefits we report here serve as impetus to optimize phrenic stimulation protocols and examine impact on duration of MV and post-operative pulmonary complications. We

acknowledge that selecting the stimulated hemidiaphragm side based on surgeon discretion is not a statistical randomization. In this initial effort, we had to compromise the ideal experimental design to meet the surgical care of the patient. Moreover, supramaximal stimulation and contraction of a hemidiaphragm applies slight lengthening forces upon the contra-lateral unstimulated side. Diaphragm stretches might be enough to blunt the detrimental effects of MV and inactivity (49), and the unstimulated side may not represent a pure “inactivity” control. In this case, our results would underestimate the benefits of intermittent phrenic stimulation. Finally, we measured protein abundance in whole-tissue homogenates to gain insight into potential mechanisms of protection. However, the effects with our protocol seem fiber type-specific, which would require single fiber analysis that are difficult to complete with the amount of tissue available for our study.

Conclusion. Intermittent, unilateral phrenic stimulation during cardiothoracic surgery led to greater maximal specific force and fiber cross-sectional area in slow fibers of the stimulated hemidiaphragm, without significantly affecting fast diaphragm fiber function. Maximal specific force was positively associated with fiber cross sectional area and inversely related to MV duration. Abundance and post-translational modifications of myofibrillar proteins and titin-linked proteins from whole-tissue homogenates remained unchanged. These findings suggest that protection of maximal specific force occurred by a mechanism independent of effects on sarcomeric protein degradation. Our study supports further examination of intraoperative phrenic stimulation to optimize protocols

for increases in fiber size or force and investigation of its potential to decrease postoperative complications.

ACKNOWLEDGEMENTS

We thank the Division of Cardiothoracic Surgery research coordinators with their regulatory assistance. We are grateful for the participation of the patients.

REFERENCES

1. Funk GC, Anders S, Breyer MK, Burghuber OC, Edelmann G, Heindl W, Hinterholzer G, Kohansal R, Schuster R, Schwarzmaier-D'Assie A, Valentin A, Hartl S. Incidence and outcome of weaning from mechanical ventilation according to new categories. *Eur Respir J* 2010; 35: 88-94.
2. Welvaart WN, Paul MA, Kuster DW, van Wieringen W, Rustenburg F, Stienen GJ, Vonk-Noordegraaf A, Ottenheijm CA. Gene expression profile in the diaphragm following contractile inactivity during thoracic surgery. *Int J Physiol Pathophysiol Pharmacol* 2011; 3: 167-175.
3. Huang TT, Deoghare HV, Smith BK, Beaver TM, Baker HV, Mehinto AC, Martin AD. Gene expression changes in the human diaphragm after cardiothoracic surgery. *J Thorac Cardiovasc Surg* 2011.
4. Hooijman PE, Beishuizen A, Witt CC, de Waard MC, Girbes AR, Spoelstra-de Man AM, Niessen HW, Manders E, van Hees HW, van den Brom CE, Silderhuis V, Lawlor MW, Labeit S, Stienen GJ, Hartemink KJ, Paul MA, Heunks LM, Ottenheijm CA. Diaphragm muscle fiber weakness and ubiquitin-proteasome activation in critically ill patients. *Am J Respir Crit Care Med* 2015; 191: 1126-1138.
5. Levine S, Biswas C, Dierov J, Barsotti R, Shrager JB, Nguyen T, Sonnad S, Kucharchzuk JC, Kaiser LR, Singhal S, Budak MT. Increased proteolysis, myosin depletion, and atrophic AKT-FOXO signaling in human diaphragm disuse. *Am J Respir Crit Care Med* 2011; 183: 483-490.
6. Matecki S, Dridi H, Jung B, Saint N, Reiken SR, Scheuermann V, Mrozek S, Santulli G, Umanskaya A, Petrof BJ, Jaber S, Marks AR, Lacampagne A. Leaky ryanodine receptors contribute to diaphragmatic weakness during mechanical ventilation. *Proc Natl Acad Sci U S A* 2016; 113: 9069-9074.
7. Levine S, Nguyen T, Taylor N, Friscia ME, Budak MT, Rothenberg P, Zhu J, Sachdeva R, Sonnad S, Kaiser LR, Rubinstein NA, Powers SK, Shrager JB. Rapid disuse atrophy of diaphragm fibers in mechanically ventilated humans. *N Engl J Med* 2008; 358: 1327-1335.
8. van den Berg M, Hooijman PE, Beishuizen A, de Waard MC, Paul MA, Hartemink KJ, van Hees HWH, Lawlor MW, Brocca L, Bottinelli R, Pellegrino MA, Stienen GJM, Heunks LMA, Wust RCI, Ottenheijm CAC. Diaphragm Atrophy and Weakness in the Absence of Mitochondrial Dysfunction in the Critically Ill. *Am J Respir Crit Care Med* 2017; 196: 1544-1558.
9. van der Pijl RJ, Granzier HL, Ottenheijm CAC. Diaphragm contractile weakness due to reduced mechanical loading: role of titin. *Am J Physiol Cell Physiol* 2019; 317: C167-C176.
10. Granzier HL, Labeit S. Titin and its associated proteins: the third myofilament system of the sarcomere. *Adv Protein Chem* 2005; 71: 89-119.
11. Lindqvist J, van den Berg M, van der Pijl R, Hooijman PE, Beishuizen A, Elshof J, de Waard M, Girbes A, Spoelstra-de Man A, Shi ZH, van den Brom C, Bogaards S, Shen S, Strom J, Granzier H, Kole J, Musters RJP, Paul MA, Heunks LMA, Ottenheijm CAC. Positive End-Expiratory Pressure Ventilation Induces Longitudinal Atrophy in Diaphragm Fibers. *Am J Respir Crit Care Med* 2018; 198: 472-485.

12. Lange S, Xiang F, Yakovenko A, Vihola A, Hackman P, Rostkova E, Kristensen J, Brandmeier B, Franzen G, Hedberg B, Gunnarsson LG, Hughes SM, Marchand S, Sejersen T, Richard I, Edstrom L, Ehler E, Udd B, Gautel M. The kinase domain of titin controls muscle gene expression and protein turnover. *Science* 2005; 308: 1599-1603.
13. Ottenheijm CA, van Hees HW, Heunks LM, Granzier H. Titin-based mechanosensing and signaling: role in diaphragm atrophy during unloading? *Am J Physiol Lung Cell Mol Physiol* 2011; 300: L161-166.
14. Hussain SN, Cornachione AS, Guichon C, Al Khunaizi A, Leite Fde S, Petrof BJ, Mofarrahi M, Moroz N, de Varennes B, Goldberg P, Rassier DE. Prolonged controlled mechanical ventilation in humans triggers myofibrillar contractile dysfunction and myofilament protein loss in the diaphragm. *Thorax* 2016; 71: 436-445.
15. Hulzebos EH, Helders PJ, Favie NJ, De Bie RA, Brutel de la Riviere A, Van Meeteren NL. Preoperative intensive inspiratory muscle training to prevent postoperative pulmonary complications in high-risk patients undergoing CABG surgery: a randomized clinical trial. *Jama* 2006; 296: 1851-1857.
16. Yang M, Wang H, Han G, Chen L, Huang L, Jiang J, Li S. Phrenic nerve stimulation protects against mechanical ventilation-induced diaphragm dysfunction in rats. *Muscle Nerve* 2013; 48: 958-962.
17. Reynolds SC, Meyyappan R, Thakkar V, Tran BD, Nolette MA, Sadarangani G, Sandoval RA, Bruulsema L, Hannigan B, Li JW, Rohrs E, Zurba J, Hoffer JA. Mitigation of Ventilator-induced Diaphragm Atrophy by Transvenous Phrenic Nerve Stimulation. *Am J Respir Crit Care Med* 2017; 195: 339-348.
18. Masmoudi H, Persichini R, Cecchini J, Delemazure J, Dres M, Mayaux J, Demoule A, Assouad J, Similowski T. Corrective effect of diaphragm pacing on the decrease in cardiac output induced by positive pressure mechanical ventilation in anesthetized sheep. *Respir Physiol Neurobiol* 2017; 236: 23-28.
19. Sotak M, Roubik K, Henlin T, Tyll T. Phrenic nerve stimulation prevents diaphragm atrophy in patients with respiratory failure on mechanical ventilation. *BMC Pulm Med* 2021; 21: 314.
20. Ahn B, Beaver T, Martin T, Hess P, Brumback BA, Ahmed S, Smith BK, Leeuwenburgh C, Martin AD, Ferreira LF. Phrenic nerve stimulation increases human diaphragm fiber force after cardiothoracic surgery. *Am J Respir Crit Care Med* 2014; 190: 837-839.
21. Graham BL, Steenbruggen I, Miller MR, Barjaktarevic IZ, Cooper BG, Hall GL, Hallstrand TS, Kaminsky DA, McCarthy K, McCormack MC, Oropez CE, Rosenfeld M, Stanojevic S, Swanney MP, Thompson BR. Standardization of Spirometry 2019 Update. An Official American Thoracic Society and European Respiratory Society Technical Statement. *Am J Respir Crit Care Med* 2019; 200: e70-e88.
22. Laveneziana P, Albuquerque A, Aliverti A, Babb T, Barreiro E, Dres M, Dube BP, Fauroux B, Gea J, Guenette JA, Hudson AL, Kabitz HJ, Laghi F, Langer D, Luo YM, Neder JA, O'Donnell D, Polkey MI, Rabinovich RA, Rossi A, Series F, Similowski T, Spengler CM, Vogiatzis I, Verges S. ERS statement on respiratory muscle testing at rest and during exercise. *Eur Respir J* 2019; 53.

23. Cavayas YA, Eljaiek R, Rodrigue E, Lamarche Y, Girard M, Wang HT, Levesque S, Denault AY. Preoperative Diaphragm Function Is Associated With Postoperative Pulmonary Complications After Cardiac Surgery. *Crit Care Med* 2019; 47: e966-e974.
24. Bruni A, Garofalo E, Pasin L, Serraino GF, Cammarota G, Longhini F, Landoni G, Lembo R, Mastroberto P, Navalesi P, Ma GICG. Diaphragmatic Dysfunction After Elective Cardiac Surgery: A Prospective Observational Study. *J Cardiothorac Vasc Anesth* 2020; 34: 3336-3344.
25. Laghlam D, Le MP, Srour A, Monsonogo R, Estagnasie P, Brusset A, Squara P. Diaphragm Dysfunction After Cardiac Surgery: Reappraisal. *J Cardiothorac Vasc Anesth* 2021; 35: 3241-3247.
26. Fischer MO, Brotons F, Briant AR, Suehiro K, Gozdzik W, Sponholz C, Kirkeby-Garstad I, Joosten A, Nigro Neto C, Kunstyr J, Parienti JJ, Abou-Arab O, Ouattara A, group Vs. Postoperative Pulmonary Complications After Cardiac Surgery: The VENICE International Cohort Study. *J Cardiothorac Vasc Anesth* 2021.
27. Jansen D, Jonkman AH, Vries HJ, Wennen M, Elshof J, Hoofs MA, van den Berg M, Man AME, Keijzer C, Scheffer GJ, van der Hoeven JG, Girbes A, Tuinman PR, Marcus JT, Ottenheijm CAC, Heunks L. Positive end-expiratory pressure affects geometry and function of the human diaphragm. *J Appl Physiol (1985)* 2021; 131: 1328-1339.
28. Dianti J, Angriman F, Ferreyro BL, Sklar MC, Brochard L, Ferguson ND, Goligher EC. Association of Mortality with Neuromuscular Blockade Differs according to Baseline Diaphragm Thickness. *Am J Respir Crit Care Med* 2020; 202: 1717-1720.
29. Tang H, C LK, Lee M, Gao Y, Xia H, Olguin F, Fraga DA, Ayers K, Choi S, Kim M, Tehrani A, Sowb YA, Rando TA, Shrager JB. Smad3 initiates oxidative stress and proteolysis that underlies diaphragm dysfunction during mechanical ventilation. *Sci Rep* 2017; 7: 14530.
30. Jaber S, Petrof BJ, Jung B, Chanques G, Berthet JP, Rabuel C, Bouyabrine H, Courouble P, Koechlin-Ramonatxo C, Sebbane M, Similowski T, Scheuermann V, Mebazaa A, Capdevila X, Mornet D, Mercier J, Lacampagne A, Philips A, Matecki S. Rapidly progressive diaphragmatic weakness and injury during mechanical ventilation in humans. *Am J Respir Crit Care Med* 2011; 183: 364-371.
31. Hooijman PE, Paul MA, Stienen GJ, Beishuizen A, Van Hees HW, Singhal S, Bashir M, Budak MT, Morgen J, Barsotti RJ, Levine S, Ottenheijm CA. Unaffected contractility of diaphragm muscle fibers in humans on mechanical ventilation. *Am J Physiol Lung Cell Mol Physiol* 2014; 307: L460-470.
32. Supinski GS, Westgate P, Callahan LA. Correlation of maximal inspiratory pressure to transdiaphragmatic twitch pressure in intensive care unit patients. *Crit Care* 2016; 20: 77.
33. Hermans G, Agten A, Testelmans D, Decramer M, Gayan-Ramirez G. Increased duration of mechanical ventilation is associated with decreased diaphragmatic force: a prospective observational study. *Crit Care* 2010; 14: R127.
34. Corpeno R, Dworkin B, Cacciani N, Salah H, Bergman HM, Ravara B, Vitadello M, Gorza L, Gustafson AM, Hedstrom Y, Petersson J, Feng HZ, Jin JP, Iwamoto H, Yagi N, Artemenko K, Bergquist J, Larsson L. Time course analysis of mechanical ventilation-induced diaphragm contractile muscle dysfunction in the rat. *J Physiol* 2014; 592: 3859-3880.

35. Thomas D, Maes K, Agten A, Heunks L, Dekhuijzen R, Decramer M, Van Hees H, Gayan-Ramirez G. Time course of diaphragm function recovery after controlled mechanical ventilation in rats. *J Appl Physiol (1985)* 2013; 115: 775-784.
36. Welvaart WN, Paul MA, Stienen GJ, van Hees HW, Loer SA, Bouwman R, Niessen H, de Man FS, Witt CC, Granzier H, Vonk-Noordegraaf A, Ottenheijm CA. Selective diaphragm muscle weakness after contractile inactivity during thoracic surgery. *Ann Surg* 2011; 254: 1044-1049.
37. Ermilov LG, Pulido JN, Atchison FW, Zhan WZ, Ereth MH, Sieck GC, Mantilla CB. Impairment of diaphragm muscle force and neuromuscular transmission after normothermic cardiopulmonary bypass: effect of low-dose inhaled CO. *Am J Physiol Regul Integr Comp Physiol* 2010; 298: R784-789.
38. Lagier D, Velly LJ, Guinard B, Bruder N, Guidon C, Vidal Melo MF, Alessi MC. Perioperative Open-lung Approach, Regional Ventilation, and Lung Injury in Cardiac Surgery. *Anesthesiology* 2020; 133: 1029-1045.
39. Goligher EC, Dres M, Patel BK, Sahetya SK, Beitler JR, Telias I, Yoshida T, Vaporidi K, Grieco DL, Schepens T, Grasselli G, Spadaro S, Dianti J, Amato M, Bellani G, Demoule A, Fan E, Ferguson ND, Georgopoulos D, Guerin C, Khemani RG, Laghi F, Mercat A, Mojoli F, Ottenheijm CAC, Jaber S, Heunks L, Mancebo J, Mauri T, Pesenti A, Brochard L. Lung- and Diaphragm-Protective Ventilation. *Am J Respir Crit Care Med* 2020; 202: 950-961.
40. Ataya A, Silverman EP, Bagchi A, Sarwal A, Criner GJ, McDonagh DL. Temporary Transvenous Diaphragmatic Neurostimulation in Prolonged Mechanically Ventilated Patients: A Feasibility Trial (RESCUE 1). *Crit Care Explor* 2020; 2: e0106.
41. Reynolds S, Ebner A, Meffen T, Thakkar V, Gani M, Taylor K, Clark L, Sadarangani G, Meyyappan R, Sandoval R, Rohrs E, Hoffer JA. Diaphragm Activation in Ventilated Patients Using a Novel Transvenous Phrenic Nerve Pacing Catheter. *Crit Care Med* 2017; 45: e691-e694.
42. Masmoudi H, Coirault C, Demoule A, Mayaux J, Beuvin M, Romero N, Assouad J, Similowski T. Can phrenic stimulation protect the diaphragm from mechanical ventilation-induced damage? *Eur Respir J* 2013; 42: 280-283.
43. Kramer C, Zoubaa S, Kretschmer A, Jordan D, Blobner M, Fink H. Denervation versus pre- and postsynaptic muscle immobilization: Effects On acetylcholine- and muscle-specific tyrosine kinase receptors. *Muscle Nerve* 2017; 55: 101-108.
44. Dridi H, Jung B, Yehya M, Daurat A, Reiken S, Moreau J, Marks AR, Matecki S, Lacampagne A, Jaber S. Late Ventilator-Induced Diaphragmatic Dysfunction After Extubation. *Crit Care Med* 2020; 48: e1300-e1305.
45. Salah H, Li M, Cacciani N, Gastaldello S, Ogilvie H, Akkad H, Namuduri AV, Morbidoni V, Artemenko KA, Balogh G, Martinez-Redondo V, Jannig P, Hedstrom Y, Dworkin B, Bergquist J, Ruas J, Vigh L, Salviati L, Larsson L. The chaperone co-inducer BGP-15 alleviates ventilation-induced diaphragm dysfunction. *Science translational medicine* 2016; 8: 350ra103.
46. Hyatt HW, Ozdemir M, Yoshihara T, Nguyen BL, Deminice R, Powers SK. Calpains play an essential role in mechanical ventilation-induced diaphragmatic weakness and mitochondrial dysfunction. *Redox Biol* 2021; 38: 101802.

47. Tang H, Shrager JB. The Signaling Network Resulting in Ventilator-induced Diaphragm Dysfunction. *American journal of respiratory cell and molecular biology* 2018; 59: 417-427.
48. Irving M. Regulation of Contraction by the Thick Filaments in Skeletal Muscle. *Biophys J* 2017; 113: 2579-2594.
49. Welvaart WN, Paul MA, van Hees HW, Stienen GJ, Niessen JW, de Man FS, Sieck GC, Vonk-Noordegraaf A, Ottenheijm CA. Diaphragm muscle fiber function and structure in humans with hemidiaphragm paralysis. *Am J Physiol Lung Cell Mol Physiol* 2011; 301: L228-235.


Table 1. Patient characteristics.

	Mean (SD)
Age (years)	59 (12)
gender	9F/10M
BMI (Kg/m ²)	30 (4)
%FVC	88 (12)
%FEV1	85 (13)
PI Max (cmH ₂ O)	90 (23)
MyHC1 (%)	64 (10)
Time (min) Intubation to biopsy	278 (68)

F: female; M: male; BMI: body mass index; %FVC: forced vital capacity; %FEV1: forced expiratory volume in the first second; PImax: Maximal inspiratory pressure; CABG: coronary artery bypass grafting; AVR: aortic valve replacement; MVR: mitral valve replacement; MyHC1: type 1 myosin heavy chain; v: number of vessels grafted

FIGURE 1 – CONSORT diagram.

CONSORT Flow Diagram






Figure 2. In slow fibers, maximal calcium activated force (**A**), fiber cross-sectional area (**B**), and maximal specific force (**C**) were significantly greater in the stimulated hemidiaphragm. Averaged specific force vs pCa relationship (**D**). In fast fibers, no significant differences in maximal force (**E**), fiber cross-sectional area (**F**), maximal specific force (**G**), or the specific force vs pCa relationship (**H**) were detected. Mixed effect model; mean and standard deviation shown. A-C and E-G dots are data from individual fibers. Statistical analysis by linear mixed model.




Figure 3. Content of titin fragments and titin-binding proteins. Western blot images and data of the N-terminal and C-terminal fragments of titin (**A**). (**B**) Calpain-3 (CAPN3), muscle specific protease at the N2A region of titin. Full-length, inactive calpain-3 (P94) and calpain fragments comprise the CAPN3 fraction. (**C**) Titin-binding proteins. Muscle Lim-Protein (MLP), telethonin-binding protein in the Z-disk region; muscle ankyrin repeat proteins 1 and 2 (MARP1 and MARP2), stress-responsive proteins at the N2A region of titin; and vinculin (VCL), membrane protein involved in sarcomere stability. Sample Western blot images are from the stimulated (STIM) and unstimulated (UNSTIM) sides of subjects 1 and 2. P-values shown from Wilcoxon tests; mean and standard deviation shown.




FIGURE 4 – Sarcomere protein phosphorylation. (A) Sample images of Pro-Q (phosphorylation) and SYPRO Ruby (total protein). Images are from the stimulated (STIM) and unstimulated (UNSTIM) sides of subjects 1 and 2. (B) Figures depict phosphorylation state of titin, nebulin, myosin heavy chain (MyHC), myosin binding protein C (Mybpc), and actin. P-values are from Wilcoxon tests; mean and standard deviation shown.