

Supplementary tables for “Covid-19 is a leading cause of death in children and young people ages 0-19 years in the United States”

Ages: < 28 days			
10 Leading Causes of Death (Infants)	Deaths	Crude Rate	Rank
#Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	3374	0.9	1
#Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3047	0.8	2
#Newborn affected by maternal complications of pregnancy (P01)	1240	0.3	3
#Newborn affected by complications of placenta, cord and membranes (P02)	731	0.2	4
#Bacterial sepsis of newborn (P36)	579	0.2	4
#Respiratory distress of newborn (P22)	409	0.1	6
#Neonatal hemorrhage (P50-P52,P54)	335	0.1	6
#Intrauterine hypoxia and birth asphyxia (P20-P21)	320	0.1	6
#Necrotizing enterocolitis of newborn (P77)	312	0.1	6
#Atelectasis (P28.0-P28.1)	230	0.1	6

Supplementary Table 1. Leading causes of death from the rankable causes on the NCHS 130 Selected Causes of Infant Death List, for neonates (< 28 days old) in 2019 in the US ranked. There were 36 Covid-19 deaths (April 1, 2021-March 31, 2022) for this age group. Deaths, crude rates per 100,000 live births, and ranks are shown.

Ages: 28-364 days			
10 Leading Causes of Death (Infants)	Deaths	Crude Rate	Rank
#Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1254	0.3	1
#Accidents (unintentional injuries) (V01-X59)	1125	0.3	1
#Sudden infant death syndrome (R95)	1110	0.3	1
#Diseases of the circulatory system (I00-I99)	326	0.1	4
#Assault (homicide) (*U01,X85-Y09)	247	0.1	4
#Diarrhea and gastroenteritis of infectious origin (A09)	180	0	5
#Influenza and pneumonia (J09-J18)	156	0	5
#Chronic respiratory disease originating in the perinatal period (P27)	148	0	5
#Septicemia (A40-A41)	131	0	5
#Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	71	0	5

Supplementary Table 2. Leading causes of death from the rankable causes on the NCHS 130 Selected Causes of Infant Death List, for infants 28-364 days old in 2019 in the US ranked. There were 103 Covid-19 deaths (April 1, 2021-March 31, 2022) for this age group. Deaths, crude rates per 100,000 live births, and ranks are shown.

Ages: 0-364 days			
10 Leading Causes of Death (Infants)	Deaths	Crude Rate	Rank
#Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4301	1.1	1
#Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	3445	0.9	2
#Accidents (unintentional injuries) (V01-X59)	1266	0.3	3
#Sudden infant death syndrome (R95)	1248	0.3	3

#Newborn affected by maternal complications of pregnancy (P01)	1245	0.3	3
#Newborn affected by complications of placenta, cord and membranes (P02)	742	0.2	6
#Bacterial sepsis of newborn (P36)	603	0.2	6
#Respiratory distress of newborn (P22)	424	0.1	8
#Diseases of the circulatory system (I00-I99)	406	0.1	8
#Necrotizing enterocolitis of newborn (P77)	354	0.1	8

Supplementary Table 3. Leading causes of death from the rankable causes on the NCHS 130 Selected Causes of Infant Death List, for infants 0-364 days old in 2019 in the US ranked. There were 139 Covid-19 deaths (April 1, 2021-March 31, 2022) for this age group. Deaths, crude rates per 100,000 live births, and ranks are shown.