

Projecting the seasonality of endemic COVID-19

Supplementary materials

Contents

Investigators	02
Supplemental Figure S1	03
Supplemental Figure S2	04
Supplemental Figure S3	05
Supplemental Figure S4	06
Supplemental Figure S5	07
Supplemental Figure S6	08
Supplemental Figure S7	09
Supplemental Figure S8	10

Investigators

Jeffrey P. Townsend^{1,2,3,4*},

April D. Lamb⁵,

Hayley B. Hassler¹,

Pratha Sah⁶,

Aia Alvarez Nishio⁷,

Cameron Nguyen⁵,

Alexandra Tew⁵,

Alison P. Galvani⁶,

Alex Dornburg⁵

Affiliations

¹*Department of Biostatistics, Yale School of Public Health; New Haven, Connecticut, USA*

²*Department of Ecology and Evolutionary Biology, Yale University; New Haven, Connecticut, USA*



³*Program in Computational Biology and Bioinformatics, Yale University; New Haven, Connecticut, USA*

⁴*Program in Microbiology, Yale University; New Haven, Connecticut, USA*

⁵*Department of Bioinformatics and Genomics, University of North Carolina at Charlotte; Charlotte, NC*

⁶*Center for Infectious Disease Modeling and Analysis, Yale University; New Haven, Connecticut 06510, USA*


⁷*Yale College; New Haven, Connecticut 06510, USA*



Supplementary Figure S1

HCoV-NL63


Proportions of HCoV-NL63 infections by month across locations in (A) North America, (B) Europe, and (C) Asia. Vertical gray bars indicate 95% confidence interval of raw monthly infections.



Supplementary Figure S2


Proportions of HCoV-229E infections by month across locations in (A) North America, (B) Europe, and (C) Asia. Vertical gray bars indicate 95% confidence interval of raw monthly infections.

HCoV-229E




Supplementary Figure S3

Proportions of HCoV-HKU1 infections by month across locations in (A) North America, (B) Europe, and (C) Asia. Vertical gray bars indicate 95% confidence interval of raw monthly infections.




Supplementary Figure S4

Proportions of HCoV-OC43 infections by month across locations in (A) North America, (B) Europe, and (C) Asia. Vertical gray bars indicate 95% confidence interval of raw monthly infections.




Supplementary Figure S5

Sensitivity of proportions of SARS-CoV-2 infections by month to method of time tree inference in North America (tan) and Europe (blue).




Supplementary Figure S6

Sensitivity of proportions of SARS-CoV-2 infections by month to method of time tree inference in Asia.



Supplementary Figure S7

Sensitivity of proportions of SARS-CoV-2 infections by month to use of only viral sequences deemed non-recombinating by location, in North America (tan) and Europe (blue).



Supplementary Figure S8

Sensitivity of proportions of SARS-CoV-2 infections by month to use of only viral sequences deemed non-recombinant by location, in Asia.