

Directed functional brain connectivity is altered in sub-threshold amyloid- β accumulators

Supplementary Information

Mite Mijalkov^{1,*}, Dániel Veréb¹, Anna Canal-Garcia¹, Giovanni Volpe², Joana B. Pereira^{1,3,*}, Alzheimer's Disease Neuroimaging Initiative

¹ Department of Neurobiology, Care Sciences and Society, Karolinska Institutet, Stockholm, Sweden.

² Department of Physics, Goteborg University, Goteborg, Sweden.


³ Memory Research Unit, Department of Clinical Sciences Malmö, Lund University, Lund, Sweden.

* Corresponding authors: Mite Mijalkov and Joana B. Pereira,


Address: KI, Dept. NVS, division of clinical geriatrics, Neo 7th floor, Blickagången 16, 141 83 Huddinge, Sweden.

\emails: mite.mijalkov@ki.se // joana.pereira@ki.se


A) Lag 2: Clustering coefficient



B) Lag 4: Clustering coefficient



C) Lag 2: Global efficiency



D) Lag 4: Global efficiency




Fig S1. Differences between different groups in global directed functional network topology. Between-group differences (calculated as group 2 - group 1 in the figure) in clustering coefficient at A) lag 2 and B) lag 4 as well as in global efficiency at C) lag 2 and D) lag 4. Orange circles represent the differences in the corresponding network measures as a function of network density; the upper and lower bounds of the 95% confidence intervals (CI) are plotted in blue. Differences are considered statistically significant if they fall outside the CIs.