

Supplemental Materials: Details of statistical regression models

(1) Models addressing relationship of insomnia as measured by the ISI to various outcomes, as described in text:

a) Model relating sleep to occupational functioning:

Generalized least squares fit by REML

Model: work ~ isi_total + age + gender

AIC	BIC	logLik
6160.261	6192.799	-3074.131

Correlation Structure: AR(1)

Formula: ~1 | PtID

Parameter estimate(s):

Phi
0.6263705

Coefficients:

	Value	Std. Error	t-value	p-value
(Intercept)	2.2535563	0.12275909	28.359910	0.0000
ISI Total	0.0909617	0.00867815	10.481693	0.0000
age	-0.0961472	0.07137116	-1.347144	0.1781
gender	0.1531653	0.10542437	1.452846	0.1465

Residual Standard Error	1.819436	
Degrees of Freedom	1678 total	1674 residual

b) Model relating sleep to self-reported likelihood of leaving current field:

Generalized least squares fit by REML

Model: infield ~ isi_total + age + gender

AIC	BIC	logLik
5527.824	5560.351	-2757.912

Correlation Structure: AR(1)

Formula: ~1 | PtID

Parameter estimate(s):

Phi
0.7662417

Coefficients:

	Value	Std. Error	t-value	p-value
(Intercept)	3.253587	0.11472488	28.359910	0.0000
ISI Total	0.050418	0.00734686	6.862578	0.0000
age	0.211740	0.07631342	2.774610	0.0056
gender	0.085483	0.11137435	0.767531	0.4429

Residual Standard Error	1.750745	
Degrees of Freedom	1675 total	1671 residual

a) Model relating sleep to thoughts of suicide, self-harm, or being better off dead:

Generalized least squares fit by REML

Model: phq9i9 ~ isi_total + age + gender

AIC	BIC	logLik
1060.662	1088.358	-524.331

Correlation Structure: AR(1)

Formula: ~1 | PtID

Parameter estimate(s):

Phi
0.5447143

Coefficients:

	Value	Std. Error	t-value	p-value
(Intercept)	-0.10044693	0.04722915	-2.126799	0.0338
ISI Total	0.02447273	0.00349304	7.006141	0.0000
age	-0.02131648	0.02412003	-0.883767	0.3771
gender	0.03231543	0.03481727	0.928144	0.3536

Residual Standard Error	0.5206405	
Degrees of Freedom	751 total	747 residual

(2) Core model addressing predictors of ISI from text:

Generalized least squares fit by REML

Model: ISI total score ~ Personal history of COVID-19 infection + Family member with COVID-19 infection + Past 2 weeks COVID-related occupational stressors + Peak COVID-related occupational stressors + age + gender

AIC	BIC	logLik
9392.458	9441.313	-4687.229

Correlation Structure: AR(1)

Formula: ~1 | PtID

Parameter estimate(s):

Phi
0.7159753

Coefficients:

	Value	Std. Error	t-value	p-value
(Intercept)	9.726505	0.5120368	18.995716	0.0000
Personal history of COVID-19 infection	0.692786	0.2236221	3.098021	0.0020
Family member with COVID-19 infection	-0.096676	0.2210195	-0.437409	0.6619
Past 2 weeks COVID-related occupational stressors	1.704757	0.1402930	12.151402	0.0000
Peak COVID-related occupational stressors	0.082858	0.0256810	3.226414	0.0013
Age	0.014379	0.2163650	0.066459	0.9470
Gender	0.559322	0.3131433	1.786152	0.0743

Correlation:

(Intr)	PERSONAL HISTORY OF COVID-19 INFECTION	FAMILY MEMBER WITH COVID-19 INFECTION	Past 2 weeks COVID-related	PEAK COVID-19 RELATED OCCUPATIONAL STRESSORS	age
--------	--	---------------------------------------	----------------------------	--	-----

				occupational stressors		
Personal history of COVID-19 infection	0.170					
Family member with COVID-19 infection	0.041	-0.312				
Past 2 weeks COVID-related occupational stressors	0.328	-0.014	-0.036			
Peak COVID-related occupational stressors	-0.902	-0.186	-0.027	-0.381		
age	-0.108	0.064	-0.103	0.041	0.120	
gender	-0.069	0.025	0.000	0.015	-0.096	-0.007

Standardized residuals:

Min	Q1	Med	Q3	Max
-2.88856050	-0.67696599	0.03740746	0.67985578	2.97731249

Residual Standard Error	5.090767		
Degrees of Freedom	1690 total	1683 residual	

(3) Models for Fig 1A top

> summary(fitisi)

Generalized least squares fit by REML

Model: ISI total score ~ Personal history of COVID-19 infection + Family member with COVID-19 infection + Past 2 weeks COVID-related occupational stressors + Peak COVID-related occupational stressors + age + gender

AIC	BIC	logLik
9392.458	9441.313	-4687.229

Correlation Structure: AR(1)

Formula: ~1 | PtID

Parameter estimate(s):

Phi
0.7159753

Coefficients:

	Value	Std.Error	t-value	p-value
(Intercept)	9.726505	0.5120368	18.995716	0.0000
Personal history of COVID-19 infection	0.692786	0.2236221	3.098021	0.0020
Family member with COVID-19 infection	-0.096676	0.2210195	-0.437409	0.6619
Past 2 weeks COVID-related occupational stressors	1.704757	0.1402930	12.151402	0.0000
Peak COVID-related occupational stressors	0.082858	0.0256810	3.226414	0.0013
age	0.014379	0.2163650	0.066459	0.9470
gender	0.559322	0.3131433	1.786152	0.0743

Correlation:

	(Intr)	PERSONAL HISTORY OF COVID-19 INFECTION	FAMILY MEMBER WITH COVID-19 INFECTION	Past 2 weeks COVID-related occupational stressors	PEAK COVID-19 RELATED OCCUPATIONAL STRESSORS	age
Personal history of COVID-19 infection	0.170					
Family member with COVID-19 infection	0.041	-0.312				
Past 2 weeks COVID-related occupational stressors	0.328	-0.014	-0.036			
Peak COVID-related occupational stressors	-0.902	-0.186	-0.027	-0.381		

age	-0.108	0.064	-0.103	0.041	0.120	
gender	-0.069	0.025	0.000	0.015	-0.096	-0.007

Standardized residuals:

Min	Q1	Med	Q3	Max
-2.88856050	-0.67696599	0.03740746	0.67985578	2.97731249

Residual standard error	5.090767	
Degrees of freedom	1690 total	1683 residual

> summary(fitpcl)

Generalized least squares fit by REML

Model: pcl5total ~ Personal history of COVID-19 infection + Family member with COVID-19 infection + Past 2 weeks COVID-related occupational stressors + Peak COVID-related occupational stressors + age + gender

Data: data2use

AIC	BIC	logLik
12382.31	12430.92	-6182.157

Correlation Structure: AR(1)

Formula: ~1 | PtID

Parameter estimate(s):

Phi
0.738682

Coefficients:

	Value	Std. Error	t-value	p-value
Intercept	17.113036	1.4329872	11.942211	0.0000
Personal history of COVID-19 infection	1.307692	0.6287211	2.079924	0.0377
Family member with COVID-19 infection	0.379777	0.6227244	0.609864	0.5420
Past 2 weeks COVID-related occupational stressors	6.857538	0.3801570	18.038703	0.0000

Peak COVID-related occupational stressors	0.407807	0.0716705	5.690021	0.0000
age	-1.779545	0.6111902	-2.911606	0.0036
gender	0.192778	0.8780037	0.219563	0.8262

Correlation:

	(Intr)	PERSONAL HISTORY OF COVID-19 INFECTION	FAMILY MEMBER WITH COVID-19 INFECTION	Past 2 weeks COVID-related occupational stressors	PEAK COVID-19 RELATED OCCUPATIONAL STRESSORS	age
Personal history of COVID-19 infection	0.169					
Family member with COVID-19 infection	0.046	-0.308				
Past 2 weeks COVID-related occupational stressors	0.315	-0.012	-0.031			
Peak COVID-related occupational stressors	-0.900	-0.186	-0.031	-0.370		
age	-0.106	0.061	-0.105	0.036	0.119	
gender	-0.068	0.024	0.001	0.014	-0.096	-0.012

Standardized residuals:

Min	Q1	Med	Q3	Max
-2.4462126	-0.6925527	-0.1903986	0.5415783	4.3297910

Residual Standard Error	14.05411		
Degrees of Freedom	1644 total	1637 residual	

(4) Models for Fig1A bottom

> summary(fitF1)

Generalized least squares fit by REML

Model: mhsmall3f_1 ~ Personal history of COVID-19 infection + Family member with COVID-19 infection + Past 2 weeks COVID-related occupational stressors + Peak COVID-related occupational stressors + age + gender

Data: data2use

AIC	BIC	logLik
4140.04	4188.602	-2061.02

Correlation Structure: AR(1)

Formula: ~1 | PtID

Parameter estimate(s):

Phi
0.6597996

Coefficients:

	Value	Std. Error	t-value	p-value
Intercept	0.8688129	0.10431685	8.328596	0.0000
Personal history of COVID-19 infection	0.0724225	0.04499485	1.609573	0.1077
Family member with COVID-19 infection	0.0134383	0.04430643	0.303303	0.7617
Past 2 weeks COVID-related occupational stressors	0.3346461	0.03159189	10.592786	0.0000
Peak COVID-related occupational stressors	0.0146770	0.00525709	2.791850	0.0053
age	-0.0992555	0.04316311	-2.299545	0.0216
gender	-0.0071763	0.06226016	-0.115263	0.9083

Correlation:

	(Intr)	PERSONAL HISTORY OF COVID-19 INFECTION	FAMILY MEMBER WITH COVID-19 INFECTION	Past 2 weeks COVID-related occupational stressors	PEAK COVID-19 RELATED OCCUPATIONAL STRESSORS	age
Personal history of COVID-19 infection	0.177					
Family member with COVID-19 infection	0.037	-0.306				
Past 2 weeks COVID-related occupational stressors	0.367	-0.014	-0.039			
Peak COVID-related occupational stressors	-0.906	-0.189	-0.020	-0.417		
age	-0.106	0.064	-0.112	0.043	0.111	
gender	-0.062	0.030	0.000	0.019	-0.099	0.004

Standardized residuals:

Min	Q1	Med	Q3	Max
-3.1839267	-0.7271638	-0.1886885	0.5096133	3.7781573

Residual Standard Error	1.049544		
Degrees of Freedom	1636 total	1629 residual	

> summary(fitF2)

Generalized least squares fit by REML

Model: mhsmall3f_2 ~ Personal history of COVID-19 infection + Family member with COVID-19 infection + Past 2 weeks COVID-related occupational stressors + Peak COVID-related occupational stressors + age + gender

Data: data2use

AIC	BIC	logLik
4051.473	4100.035	-2016.737

Correlation Structure: AR(1)

Formula: ~1 | PtID

Parameter estimate(s):

Phi
0.6101749

Coefficients:

	Value	Std. Error	t-value	p-value
Intercept	0.03333596	0.09584669	0.347805	0.7280
Personal history of COVID-19 infection	0.00219464	0.04090616	0.053651	0.9572
Family member with COVID-19 infection	0.04240595	0.04014654	1.056279	0.2910
Past 2 weeks COVID-related occupational stressors	0.26250679	0.03102805	8.460306	0.0000
Peak COVID-related occupational stressors	0.02437243	0.00484542	5.029998	0.0000
age	-0.10231473	0.03897661	-2.625029	0.0087
gender	-0.04428009	0.05635074	-0.785794	0.4321

Correlation:

	(Intr)	PERSONAL HISTORY OF COVID-19 INFECTION	FAMILY MEMBER WITH COVID-19 INFECTION	Past 2 weeks COVID-related occupational stressors	PEAK COVID-19 RELATED OCCUPATIONAL STRESSORS	age
Personal history of COVID-19 infection	0.181					
Family member with COVID-19 infection	0.032	-0.308				
Past 2 weeks COVID-related occupational stressors	0.396	-0.015	-0.044			
Peak COVID-related	-0.909	-0.191	-0.014	-0.443		

occupational stressors						
age	-0.105	0.065	-0.116	0.047	0.106	
gender	-0.058	0.034	-0.002	0.022	-0.101	0.013

Standardized residuals:

Min	Q1	Med	Q3	Max
-3.4511443	-0.6584667	-0.1186989	0.5706452	4.0646472

Residual standard error:	0.9810603		
Degrees of freedom:	1636 total	1629 residual	

> summary(fitF3)

Generalized least squares fit by REML

Model: mhsmall3f_3 ~ Personal history of COVID-19 infection + Family member with COVID-19 infection + Past 2 weeks COVID-related occupational stressors + Peak COVID-related occupational stressors + age + gender

Data: data2use

AIC	BIC	logLik
3812.102	3860.664	-1897.051

Correlation Structure: AR(1)

Formula: ~1 | PtID

Parameter estimate(s):

Phi
0.6789277

Coefficients:

	Value	Std.Error	t-value	p-value
(Intercept)	1.5813248	0.09662911	16.364891	0.0000
Personal history of COVID-19 infection	0.1075206	0.04185497	2.568885	0.0103
Family member with COVID-19 infection	-0.0359708	0.04126998	-0.871597	0.3836
Past 2 weeks COVID-related occupational stressors	0.2230959	0.02843435	7.846000	0.0000

Peak COVID-related occupational stressors	0.0065025	0.00486237	1.337318	0.1813
age	0.0528811	0.04026294	1.313394	0.1892
gender	0.1286761	0.05802223	2.217704	0.0267

Correlation:

	(Intr)	PERSONAL HISTORY OF COVID-19 INFECTION	FAMILY MEMBER WITH COVID-19 INFECTION	Past 2 weeks COVID-related occupational stressors	PEAK COVID-19 RELATED OCCUPATIONAL STRESSORS	age
Personal history of COVID-19 infection	0.175					
Family member with COVID-19 infection	0.039	-0.306				
Past 2 weeks COVID-related occupational stressors	0.355	-0.014	-0.037			
Peak COVID-related occupational stressors	-0.905	-0.189	-0.023	-0.407		
age	-0.106	0.063	-0.110	0.041	0.113	
gender	-0.063	0.029	0.000	0.018	-0.098	0.000

Standardized residuals:

Min	Q1	Med	Q3	Max
-3.26387164	-0.68447668	0.08933059	0.70115020	3.09688658

Residual standard error	0.9659589		
Degrees of freedom	1636 total	1629 residual	