
## Supplementary Table 1: N of complete data for relevant variables across time points

Variables		Timepoints			
		Four	Three	Two	One
Insight		765	941	1117	1328
Illness severity (patient)		777	949	1117	1335
Illness severity (clinician)		775	951	1127	1335
Depression		785	952	1124	1338
PANSS		786	953	1128	1339
Neurocognition*		-	539	824	1165
General physical health		757	934	1113	1325
General mental health		757	934	1113	1325
Quality of life		764	939	1112	1324
Competencies*	Understanding	-	628	964	1253
	Appreciation	-	628	964	1253
	Reasoning	-	628	964	1253
	Choice	-	628	963	1253

<sup>\*</sup>These variables only have 3 timepoints for consistent measurements for most participants.



Supplementary Figure 1: Correlogram of relevant variables from baseline. Measures for variables are included in parenthesis. CDRS—Calgary Depression Rating Scale; CGIS—Clinical Global Impression of Symptom Severity; ITAQ—Illness and Treatment Attitude Questionnaire; MacCAT-CR—MacArthur Competence Assessment Tool- Clinical Research; PANSS—Positive and Negative Syndrome Scale; SF-12—12-item Short Form Survey; QLS—Heinrich-Carpenter Quality of Life Scale.