

Supplement 4 - Percent of episodes where type of health care provider (HCP) was the last HCP seen by episode sequence and population Area Deprivation Index (ADI) and percent of the population that is non-Hispanic white (% NHW)

	Initial HCP - No 2nd						2nd HCP - No 3rd					
	ADI 75-100 % NHW 0-25		ADI 50-75 % NHW 50-75		ADI 0-25 % NHW 75-100		ADI 75-100 % NHW 0-25		ADI 50-75 % NHW 50-75		ADI 0-25 % NHW 75-100	
	Episodes	% No 2nd	Episodes	% No 2nd	Episodes	% No 2nd	Episodes	% No 3rd	Episodes	% No 3rd	Episodes	% No 3rd
PCP	5641	74.0%	17431	71.6%	14563	69.3%	1182	67.3%	3995	65.3%	3343	64.9%
Nurse	1267	64.4%	4286	64.0%	2350	56.2%	406	65.0%	1429	61.1%	844	59.4%
PA	793	60.8%	2743	57.3%	2301	56.9%	280	58.6%	991	57.0%	826	53.3%
DO	5	80.0%	46	73.9%	67	76.1%	1	100.0%	3	100.0%	9	55.6%
DC	1629	84.0%	11124	84.3%	13206	84.9%	269	73.6%	1646	72.4%	1762	73.7%
PT	62	71.0%	318	61.3%	824	59.5%	105	72.4%	511	72.4%	1127	69.5%
LAc	25	92.0%	88	78.4%	408	83.1%	2	50.0%	16	62.5%	70	64.3%
OS	504	64.5%	2326	62.4%	3576	61.7%	210	56.7%	925	57.0%	1284	54.2%
PMR	219	62.6%	903	63.9%	1807	62.1%	111	55.0%	422	59.0%	797	57.7%
PM	226	66.8%	988	61.3%	793	61.8%	117	53.0%	486	52.7%	394	54.1%
NS	72	51.4%	280	49.3%	335	54.0%	36	58.3%	214	52.3%	218	45.0%
Neuro	69	71.0%	361	55.1%	431	57.5%	43	72.1%	169	55.0%	164	57.3%
Rheum	81	58.0%	318	65.1%	311	61.7%	43	55.8%	144	52.1%	131	49.6%
MD Oth	333	54.4%	1345	54.5%	1361	54.1%	168	63.1%	630	57.8%	651	59.4%
EM	802	64.8%	1568	61.3%	1093	56.2%	221	67.4%	495	56.8%	358	50.0%
Rad	1055	54.1%	2851	50.2%	2837	46.6%	667	53.5%	2172	49.9%	2398	47.1%
UC	175	77.7%	391	61.9%	443	58.7%	30	73.3%	113	65.5%	147	63.9%
	3rd HCP - No 4th						4th HCP - No 5th					
	ADI 75-100 % NHW 0-25		ADI 50-75 % NHW 50-75		ADI 0-25 % NHW 75-100		ADI 75-100 % NHW 0-25		ADI 50-75 % NHW 50-75		ADI 0-25 % NHW 75-100	
	Episodes	% No 4th	Episodes	% No 4th	Episodes	% No 4th	Episodes	% No 5th	Episodes	% No 5th	Episodes	% No 5th
PCP	537	61.8%	1855	62.1%	1468	59.8%	234	62.0%	954	57.0%	704	53.3%
Nurse	191	62.3%	743	51.7%	412	54.6%	89	46.1%	416	49.8%	244	46.3%
PA	129	60.5%	524	49.8%	392	44.9%	63	61.9%	285	41.4%	239	48.5%
DO	0	N/A	8	62.5%	9	66.7%	0	N/A	2	100.0%	5	20.0%
DC	99	61.6%	799	67.8%	801	67.5%	58	60.3%	357	63.6%	364	63.2%
PT	76	65.8%	369	66.1%	776	63.0%	49	71.4%	239	63.2%	461	62.0%
LAc	1	0.0%	6	50.0%	43	72.1%	2	100.0%	5	60.0%	18	55.6%
OS	142	62.0%	590	50.5%	741	52.6%	93	52.7%	348	51.1%	428	51.2%
PMR	90	56.7%	336	60.4%	545	61.5%	64	56.2%	219	56.2%	360	59.2%
PM	101	61.4%	433	55.9%	345	56.2%	72	44.4%	335	59.4%	249	53.0%
NS	40	52.5%	164	47.0%	198	42.9%	29	34.5%	124	42.7%	127	42.5%
Neuro	18	66.7%	101	57.4%	114	50.0%	12	41.7%	77	49.4%	60	41.7%
Rheum	25	60.0%	84	52.4%	74	54.1%	13	46.2%	63	49.2%	48	60.4%
MD Oth	89	58.4%	398	57.0%	422	52.1%	67	58.2%	287	48.1%	274	54.0%
EM	71	52.1%	166	48.8%	108	44.4%	35	57.1%	84	56.0%	51	43.1%
Rad	245	42.0%	969	44.9%	1088	40.9%	118	37.3%	498	36.1%	586	40.3%
UC	8	25.0%	25	44.0%	29	62.1%	3	33.3%	8	37.5%	11	27.3%

Cells in red have risk ratio 95% confidence interval crossing the PCP reference of 1

Cells highlighted with bold font have risk ratio 95% plus confidence interval greater than the PCP reference of 1 indicating greater likelihood to be the last HCP seen

PCP=primary care provider, DO=doctor of osteopathy, DC=doctor of chiropractic, PT=physical therapist, LAc=licensed acupuncturist, OS=orthopedic surgeon, PMR=physical medicine & rehabilitation, PM=pain medicine, NS=neurosurgeon, Neuro=neurologist, Rheum=rheumatologist, MD Oth = other medical physician, EM=emergency medicine, Rad=radiologist, UC=urgent care