

1 **Independent and combined effects of nutrition and sanitation interventions on enteric**
2 **pathogen carriage and child growth in rural Cambodia: a factorial cluster-randomised**
3 **controlled trial**

4
5 Amanda Lai^{1,2}, Irene Velez³, Ramya Ambikapathi⁴, Krisna Seng³, Karen Levy⁵, Erin
6 Kowalsky^{1,2}, David Holcomb¹, Konstantinos T. Konstantinidis², Oliver Cumming⁶, Joe Brown^{1*}

7
8 ¹ Department of Environmental Sciences and Engineering, Gillings School of Global Public
9 Health, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, USA

10 ² School of Civil and Environmental Engineering, Georgia Institute of Technology, Atlanta,
11 Georgia, USA

12 ³ MSI, A Tetra Tech Company, Arlington, Virginia, USA

13 ⁴ Department of Public Health, Purdue University, West Lafayette, Indiana, USA

14 ⁵ Department of Environmental and Occupational Health Sciences, University of Washington,
15 Seattle, Washington, USA

16 ⁶ Department of Disease Control, Faculty of Infectious and Tropical Diseases, London School of
17 Hygiene and Tropical Medicine, London, UK

18
19 * Corresponding author: Joe Brown, joebrown@unc.edu

20 Summary

21 Background

22 Childhood exposure to enteric pathogens associated with poor sanitation contributes to
23 undernutrition, associated with adverse effects later in life. This trial assessed the independent
24 and combined effects of nutrition and sanitation interventions on child growth outcomes and
25 enteric pathogen infection in rural Cambodia, where the prevalence of childhood stunting
26 remains high.

27 Methods

28 We conducted a factorial cluster-randomised controlled trial of 4,015 households with 4,124
29 children (1-28 months of age at endline) across three rural provinces in Cambodia. Fifty-five
30 communes (clusters) were randomly assigned to a control arm or one of three treatments: a
31 nutrition-only arm, a sanitation-only arm, and a combined nutrition and sanitation arm receiving
32 both treatments. The primary outcome was length-for-age Z-score (LAZ); other outcomes
33 included weight-for-age Z-score (WAZ), weight-for-length Z-score (WLZ), stunting, wasting,
34 underweight, and caregiver-reported diarrhoea. We assayed stool specimens from a subset of all
35 children (n = 1,620) for 27 enteric pathogens (14 bacteria, 6 viruses, 3 protozoa, and 4 soil-
36 transmitted helminths) and estimated effects of interventions on enteric pathogen detection and
37 density. Analysis was by intention-to-treat. The trial was pre-registered with ISRCTN Registry
38 ([ISRCTN77820875](https://www.isrctn.com/ISRCTN77820875)).

39 Findings

40 Self-reported adherence was high for the nutrition intervention but uptake was low for sanitation.
41 Compared with a mean LAZ of -1.04 (SD 1.2) in the control arm, children in the nutrition-only
42 arm (LAZ +0.08, 95% CI -0.01-0.18) and combined nutrition and sanitation arm (LAZ +0.10,
43 95% CI 0.01-0.20) experienced greater linear growth; there were no measurable differences in
44 LAZ in the sanitation-only arm (LAZ -0.05, 95% CI -0.16-0.05). We found no effect of any
45 intervention (delivered independently or combined) on either enteric pathogen frequency or
46 pathogen load in stool. Compared with a mean WAZ of -1.05 (SD 1.1) in the control arm,
47 children in the nutrition-only arm (WAZ +0.10, 95% CI 0.00-0.19) and combined intervention
48 arm (WAZ +0.11, 95% CI 0.03-0.20) were heavier for their age; there was no difference in WAZ
49 in the sanitation-only arm. There were no differences between arms in prevalence of stunting,
50 wasting, underweight status, one-week period prevalence of diarrhoea, pathogen prevalence, or
51 pathogen density in stool.

52 Interpretation

53 Improvements in child growth in nutrition and combined nutrition and sanitation arms are
54 consistent with previous efficacy trials of combined nutrition and sanitation interventions. We
55 found no evidence that the sanitation intervention alone improved child growth or reduced
56 enteric pathogen detection, having achieved only modest changes in access and use.

57 Funding

58 United States Agency for International Development (USAID), contracts AID-OAA-M-13-
59 00017 and AID-OAA-TO-16-00016. The contents of this publication are the sole responsibility
60 of the authors and do not necessarily reflect the views of USAID or the United States
61 Government.

62 Introduction

63 Childhood undernutrition is associated with higher susceptibility to infectious disease, reduced
64 cognitive function, and various adverse outcomes later in life¹. Growth faltering is an effect of
65 chronic undernutrition and tends to manifest in a child's first two years². Many studies have
66 focused on improving infant and child nutrition to achieve better growth outcomes^{3,4}. However,
67 nutrition interventions alone have not been successful in eliminating stunting, suggesting that
68 broader interventions addressing other important factors are needed alongside exclusive
69 breastfeeding and improved nutritional intake⁵.

70
71 Reducing early childhood exposure to enteric pathogens through safe water, sanitation, and
72 hygiene (WASH) may complement other interventions by reducing diarrhoeal diseases and
73 environmental enteric dysfunction (EED)⁶—both of which can impact early childhood growth
74 and development⁷. Observational studies have found strong associations between child growth
75 faltering and poor access to sanitation⁸. However, recent randomised controlled trials (RCTs) in
76 Zimbabwe⁹, Bangladesh³, and Kenya⁴ that delivered standalone household-level sanitation
77 interventions (not coupled with other nutrition or hygiene interventions) were not found to
78 improve child growth.

79
80 The community-led total sanitation (CLTS) framework is an approach to ending open defecation
81 (OD) through behavioural change and collective action rather than through the provision of
82 hardware and materials. CLTS and other rural promotion-based interventions shift the focus
83 from individual and household sanitation practices to a community-level concern over OD by
84 triggering collective behaviour change through powerful emotional drivers such as shame and
85 disgust, as well as positive motivators such as improved health, dignity, and pride. Observational
86 studies in Cambodia¹⁰ and Ecuador¹¹ found higher community-level sanitation coverage to be
87 associated with reduced prevalence of stunting. Despite this, recent RCTs employing promotion-
88 based interventions have found mixed effects on child growth. One trial was found to be
89 successful in improving child growth in Mali¹², but this effect was not observed in other trials
90 elsewhere^{13,14}.

91
92 This study contributes to a growing body of literature on the impact of combined nutrition and
93 sanitation interventions on early child growth, caregiver-reported diarrhoea, and detection and
94 quantification of enteric pathogens in stool as a proxy for enteric infection. While diarrhoea has
95 been widely used as a primary outcome measure in WASH studies^{3,4,12,15}, recent studies have
96 used stool-based detection of enteric pathogens¹⁵ and anthropometry measurements^{3,4,9} as
97 primary outcomes that are more objectively measurable and may also broadly indicate health
98 status by capturing cumulative effects of exposures via EED⁶. We used a factorial cluster-
99 randomised controlled trial (cRCT) to assess the independent and combined effects of nutrition
100 and sanitation interventions delivered in the context of a large-scale, USAID-funded rural
101 nutrition and sanitation/hygiene program in Cambodia. We hypothesised that children receiving
102 both sanitation and nutrition interventions would have increased linear growth compared with
103 children from control areas lacking these interventions. We further hypothesised that combined
104 nutrition and sanitation interventions would lead to synergistic improvements in linear growth
105 beyond what was realised in either standalone intervention arm. The hypothesised pathway for

106 these effects, consistent with secondary outcome measures, was reduced enteric pathogen
107 frequency and enteric pathogen load in stools (Figure 1).

108 **Methods**

109 **Study design and participants**

110 We implemented a two-by-two factorial cRCT in rural communes in three provinces in
111 Cambodia: Battambang, Pursat, and Siem Reap. The communes targeted by the program were
112 selected based on two criteria: communes where at least 30% of the population was living below
113 the poverty line according to the 2011 Cambodia Ministry of Planning’s Commune Database;
114 and communes where latrine subsidies were not then in place. This study is reported per the
115 Consolidated Standards of Reporting Trials (CONSORT) guideline (see Supplementary Material
116 for CONSORT checklist).

117

118 **Randomisation and masking**

119 In 2015, prior to the start of project activities, we randomly assigned communes to one of three
120 treatment arms (nutrition only, sanitation/hygiene only, combined nutrition and
121 sanitation/hygiene) or control arm using a random number generator with reproducible seed in
122 Stata 13 (Stata, College Station, TX). Randomisation was conducted at the commune level to
123 limit the risk of contamination between study arms and all villages within each commune
124 received the assigned intervention. Following randomisation, three communes were dropped
125 from the trial due to objections from the local governments of overlap with other current
126 programming. This resulted in 55 communes with treatment arms of different sizes: 11
127 communes in nutrition-only arm; 13 in sanitation-only arm; 12 in combined-intervention arm;
128 and 19 in control arm (Figure 2). The trial enrolled primary caregivers with a child who was born
129 after intervention implementation began (up to 28 months prior) and who had lived in the
130 commune during the child’s entire life, resulting in a participant population of children 1-28
131 months old. Neither participants nor field staff were masked to treatment status due to the nature
132 of the interventions, but data collection teams were blinded to the arm assignment and number of
133 treatment arms.

134

135 **Procedures**

136 The interventions were delivered in the 36 intervention communes over the course of two years,
137 between 2015-2017, while the remaining 19 control communes were unexposed to the
138 programmes. Two international non-governmental organizations—Save the Children and SNV—
139 provided programmatic implementation and coordinated activities with local governments. The
140 nutrition interventions included complementary feeding activities and education through
141 community-based growth promotion sessions; caregiver groups; home visits; and conditional
142 cash transfers (CCTs) linked to the utilization of key health and nutrition services focusing on
143 first 1,000 days of life. The sanitation interventions consisted primarily of CLTS as it was
144 delivered here, latrine vouchers coupled with supply-side support for sanitation and hygiene
145 products, and social behaviours change communications (SBCC). Intervention activities and
146 frequency are summarized in Table 1, and additional details about the interventions are described
147 in the Supplementary Material.

148

149 The survey was communicated in the Khmer language to assess household and child-level risk
150 factors of children under 28 months of age. Enumerators completed in-home interviews with the

151 primary caregiver of children in the household about basic household member information;
152 breastfeeding and nutrition of children up to age 28 months; number of pregnancies and child
153 births of the caregiver; intervention exposure and participation; household WASH conditions and
154 practices; and household assets/characteristics to construct wealth scores (excluding WASH
155 variables). We also documented process evaluation (PE) indicators based on self-reported receipt
156 of, and participation in, intervention activities to assess intervention fidelity and adherence,
157 respectively. We attempted to collect a stool sample from each child and randomly selected a
158 subset of stools for analysis by reverse-transcription quantitative polymerase chain reaction (RT-
159 qPCR) of 30 enteric pathogen genes using a custom-developed TaqMan Array Card (TAC;
160 ThermoFisher Scientific, Waltham, VA), as described in the Supplementary Material.

161

162 Outcomes

163 The primary outcome was length-for-age Z-score (LAZ). For children 1-24 months in age, we
164 measured recumbent length; for children 24-28 months in age, we measured standing height.
165 Herein, “length” will be inclusive of both recumbent length and standing height. Secondary
166 outcomes included weight-for-age Z-score (WAZ); weight-for-length Z-score (WLZ); proportion
167 of children stunted (LAZ<-2), underweight (WAZ<-2), and wasted (WLZ<-2); caregiver
168 reported diarrhoea; all-cause mortality; and enteric pathogen detection and quantification in
169 stool. Child length and weight were measured by trained paired enumerators following
170 guidelines from the National Health and Nutrition Examination Survey (NHANES)¹⁶. Final
171 measurements took place in August 2019, 28 months after the end of the roll-out period. Data
172 collection was completed by KHANA Centre for Population Health Research, with oversight and
173 support from Management Systems International (MSI). Data collection details, measurement
174 protocols, and PE indicators are further described in Supplementary Material.

175

176 We assessed enteric pathogens as the prevalence of individual gene targets, the number of co-
177 detected pathogens, and enteric pathogen-associated gene copies per gram of stool based on PCR
178 quantification cycle (Cq) and standard curves. *E.coli* pathotypes were defined as: EAEC (*aaiC*,
179 or *aatA*, or both), atypical EPEC (*eae* without *bfpA*, *stx1*, and *stx2*), typical EPEC (*bfpA*), ETEC
180 (*STh*, *STp*, or *LT*), and STEC (*eae* without *bfpA* and with *stx1*, *stx2*, or both). Details on nucleic
181 acid extraction and molecular assaying are described in Supplementary Material.

182

183 Statistical analysis

184 We performed an intention-to-treat (ITT) analysis for all outcomes using generalised estimating
185 equations (GEE) with robust standard errors to account for clustering at the village level. We did
186 not consider pre-intervention covariate balance¹⁷ but present secondary analyses adjusted for pre-
187 specified pre-intervention covariates in the Supplementary Material. Outcomes in each treatment
188 arm were compared to the control arm and between standalone treatment arms and the combined
189 treatment arm. We used linear regression to estimate mean differences in LAZ, WAZ, WLZ, and
190 log₁₀-transformed pathogen gene target densities and used log-linear Poisson regression to
191 estimate the prevalence ratio (PR) between arms for nutritional status (stunting, wasting, and
192 underweight), diarrhoea, and overall mortality. Enteric pathogen gene outcomes were
193 dichotomised, with positive detections defined by a Cq <35¹⁸, and Poisson regression was used
194 to estimate PRs for individual pathogens detected in stool. We further estimated the incidence
195 rate ratio (IRR) of co-detected pathogens (total and in subgroups by bacteria, viruses, protozoa,
196 and STHs) using negative binomial regression. We did not adjust for multiple comparisons for

197 growth, diarrhoea, or mortality outcomes^{3,19}, but we did apply the Benjamini-Hochberg
198 procedure to control the false discovery rate within analyses of multiple enteric pathogen
199 outcomes²⁰. Details on power calculations are included in Supplementary Material.

200

201 Ethics

202 The study received approval from the National Ethics Committee for Health Research in the
203 Cambodian Ministry of Health, Georgia Institute of Technology, and New England Institutional
204 Review Board. Prior to any data collection, the trial was explained to participants in the Khmer
205 language. Written and verbal consent were obtained prior to administering the surveys and
206 anthropometry measurements. The trial was pre-registered with ISRCTN Registry
207 ([ISRCTN77820875](https://www.isrctn.com/ISRCTN77820875)).

208 Results

209 Among 82 presumptively eligible communes, the provincial governments in 27 declined to
210 participate. Ultimately, the evaluation included 55 communes randomly assigned to one of three
211 treatment arms (n=36 communes) or control arm (n=19 communes); the control arm was
212 relatively oversized to enhance statistical efficiency of multiple hypothesis testing²¹. Figure 2
213 shows the trial profile by intervention subgroups. 4,015 households participated in endline
214 surveys; 4,005 households were included in these analyses (10 were excluded due to incomplete
215 surveys), and 4,124 children had anthropometry measures taken.

216

217 Household and caregiver characteristics were mostly similar across treatment and control groups
218 (Table 2). Primary caregivers in the control group reported lower levels of primary school
219 attendance compared to the treatment groups, but paternal primary school attendance was
220 similar. Households in the nutrition-only and sanitation-only groups had higher wealth index
221 scores compared to households in the combined intervention and control groups. The control
222 group had a higher prevalence of improved water source as their main source of drinking water
223 compared to the treatment groups.

224

225 Nutrition intervention fidelity was high, with households in the nutrition-only and combined-
226 intervention arms reporting significantly higher participation in these activities compared to the
227 sanitation-only and control groups (Table 3). Approximately 60% of households in the nutrition-
228 only and combined-intervention arms reported participating in at least four of the eight nutrition
229 intervention activities, compared to 4% in the sanitation-only and control arms. Conversely,
230 sanitation intervention fidelity was very low, with only 6% of households in the sanitation-only
231 and control arms reporting participation in any CLTS activity, compared to 14% of households
232 in the nutrition-only arm and 25% in the combined-intervention arm.

233

234 More households in the control arm (70%) had an improved water source as their main source of
235 drinking water, an indicator of nutrition intervention adherence, compared to other arms
236 (approximately 60% in other arms; Table 4). The combined intervention arm had greater access
237 to improved sanitation facilities (61%) compared to the nutrition-only (55%), sanitation-only
238 (51%), and control (52%) arms. OD (self-reported) was practiced less in the combined
239 intervention arm (7%) compared to the nutrition-only (14%), sanitation-only (16%), and control
240 (16%) arms. Notably, the sanitation-only arm experienced a significantly larger increase in
241 sanitation coverage (+25 percentage points [pp]) compared to all other arms (+14pp in nutrition-

242 only arm, +19pp in combined and control arms), though sanitation gains across all arms were
243 evident in the intervention period, reflecting a strong secular trend of sanitation expansion that
244 has been widely documented in rural Cambodia^{8,10,22,23}. Additional intervention adherence
245 indicators related to environmental hygiene are reported in the Supplementary Material.
246

247 [Primary and secondary outcomes](#)

248 Mean LAZ in the control arm was -1.04 (SD 1.20). Compared with control, children in the
249 nutrition-only arm were longer by a mean of 0.08 LAZ (95% CI -0.01, 0.18), and children in the
250 combined-intervention arm were longer by 0.10 LAZ (95% CI 0.01, 0.20), although these
251 differences were not observed in the adjusted analyses (
252

253 Table 5, Figure 3, Supplementary Material). Children in the nutrition-only arm and combined-
254 intervention arm were heavier than children in the control arm by a mean of 0.10 WAZ (95% CI
255 0.00, 0.19) and 0.11 (95% CI 0.03, 0.20), respectively. These differences were slightly attenuated
256 in the adjusted analyses (
257

258 Table 5, Supplementary Material). No differences were observed between the control arm and
259 intervention arms in terms of WLZ. Children in the combined intervention arm were also longer
260 and heavier, on average, than children in the sanitation-only arm by 0.16 LAZ (95% CI 0.04,
261 0.27) and 0.10 WAZ (95% CI 0.01, 0.20), respectively. LAZ and WAZ were similar between
262 children in the nutrition-only and combined intervention arms.

263
264 Compared with the control arm, none of the intervention arms differed in the prevalence of
265 children who experienced stunting, wasting, diarrhoea (7-day recall), or mortality (Table 6).
266 However, the combined intervention reduced underweight prevalence by 18% (PR 0.82, 95% CI
267 0.68, 0.99) relative to the control arm. Although the combined intervention did not significantly
268 impact stunting prevalence compared with the control arm or the nutrition-only arm, the
269 sanitation-only arm was associated with a 20% increase (PR 1.2, 95% CI 1.0, 1.5) in the
270 prevalence of both stunting and underweight status when compared to the combined
271 intervention. All associations with stunting and underweight were attenuated in adjusted analyses
272 (Supplementary Material).

273

274 Enteric pathogen results

275 We assessed enteric pathogen-associated gene targets in 1,620 randomly selected stools that
276 demonstrated acceptable amplification (of 4,114 stools total, see Supplementary Material): 305
277 from the nutrition arm, 333 from the sanitation arm, 438 from the combined-intervention arm,
278 and 544 from the control arm. We detected at least one bacterial gene in 87% of all samples, at
279 least one viral gene in 49% of samples, at least one protozoan gene in 20% of samples, and at
280 least one STH gene in 2% of samples. Enteroaggregative *E. coli* (EAEC), enteric pathogenic *E.*
281 *coli* (EPEC), enterovirus, *Campylobacter* spp., and enterotoxigenic *E. coli* (ETEC) were the most
282 prevalent pathogens (Table 7). We detected a mean 2.2 bacterial genes (out of 9), 0.59 viral
283 genes (out of 6), 0.21 protozoan genes (out of 4), and 0.03 STH genes (out of 4) in each sample.
284 We found no differences in the rate of bacterial, viral, protozoan, or STH gene co-detection
285 between the control arm and any treatment arm or between the combined arm and the standalone
286 intervention arms (Table 8). Prevalence increased with age for many pathogens (aEPEC, ETEC,
287 *Shigella*/EIEC (*ipah*), STEC, adenovirus, *Giardia*), while prevalence peaked for children 9-17
288 months for other pathogens (*Campylobacter* spp., *C.diff*, EAEC, *Salmonella* spp.; Supplementary
289 Material).

290
291 Examining prevalence of specific targets compared to the control arm, the nutrition-only arm
292 demonstrated increased prevalence of any bacterial gene (PR 1.06, 95% CI 1.01, 1.11),
293 adenovirus (PR 1.88, 95% CI 1.41, 2.51) and heat-labile/heat-stable ETEC (PR 2.00, 95% CI
294 1.19, 3.36) and reduced prevalence of EIEC/*Shigella* spp. (PR 0.60, 95% CI 0.39, 0.94). Children
295 in the sanitation-only arm had less EPEC (PR 0.88, 95% CI 0.78, 1.00) compared to control. In
296 the combined-intervention arm, atypical-EPEC prevalence decreased (PR 0.85, 95% CI 0.73,
297 0.98) while the prevalence increased for heat-stable ETEC (PR 1.42, 95% CI 1.01, 2.00), heat-
298 labile/heat-stable ETEC (PR 1.74, 95% CI 1.06, 2.86), and any viral gene (PR 1.16, 95% CI
299 1.02, 1.31). We found similar mixed effects when comparing pathogen gene prevalence in
300 individual treatment arms compared to the combined arm; there was slightly lower combined
301 prevalence of any bacterial target (PR 0.96, 95% CI 0.91, 1.02) and enterovirus (PR 0.82, 95%
302 CI 0.67, 1.00) in the sanitation-only arm, and we found higher prevalence of adenovirus (PR
303 1.42, 95% CI 1.07, 1.87) in the nutrition-only arm (Table 9).

304

305 Generally, differences in mean gene quantities were consistent with prevalence differences

306 (Table 9;

307 Table 10). We detected lower concentrations of pathogen-associated genes in the nutrition-only
308 and sanitation-only arms compared with the control arm; children in the nutrition-only arm
309 carried lower quantities of STEC (-1.46 log₁₀-copies, 95% CI -2.97, 0.06) and *Giardia* (-1.73
310 log₁₀-copies, 95% CI -3.02, -0.44), and children in the sanitation-only arm carried lower
311 quantities of EPEC (-0.54 log₁₀-copies, 95% CI -1.17, 0.09) and STEC (-1.71 log₁₀-copies, 95%
312 CI -3.07, -0.34). There was no measurable difference in mean gene quantities between the
313 combined and control arm. There was no significant difference in quantity of pathogen genes
314 between treatment arms after adjusting for multiple comparisons²⁰.

315 Discussion

316 We found a modest effect on growth from the nutrition-only intervention and a greater effect in
317 the combined intervention arm that was likely attributable to the nutrition intervention alone. By
318 contrast, the sanitation intervention alone was not associated with growth improvements, relative
319 to control conditions, and demonstrated significantly poorer growth than the combined
320 intervention arm. The similar impacts on linear growth between the nutrition-only and combined
321 intervention arms were consistent with the observed linear growth improvements being
322 attributable primarily to the nutrition intervention alone, further suggesting that the addition of
323 this sanitation intervention to the nutrition intervention did not produce synergistic effects.
324 Intermediate outcomes of meal frequency and dietary diversity were similar between arms, so the
325 observed effects may have been attributable to other elements of the nutrition or combined
326 intervention not captured by these measures. We observed no meaningful differences between
327 arms with respect to secondary outcome measures of WAZ, WLZ, stunting, wasting,
328 underweight status, diarrhoea, mortality, pathogen prevalence, pathogen co-detection rate, or
329 pathogen gene copy quantity. Although molecular detection of a specific pathogen in stool does
330 not necessarily signal active enteric infection, potential for disease, or direct effects on the
331 individual, it does unambiguously indicate prior exposure to that pathogen. Our specific
332 pathogen targets were selected *a priori* based on a range of globally observed enteric pathogens
333 and may not fully capture the relevant enteric pathogens in rural Cambodia; however, the
334 consistently high pathogen prevalence across all treatment arms suggests the suite of
335 interventions assessed in this trial did not prevent environmental exposure to enteric pathogens.

336
337 Our findings are consistent with results from several recent randomised factorial WASH and
338 nutrition efficacy trials reporting protective effects of combined/integrated interventions and null
339 effects of WASH alone on child growth outcomes^{3,4,9}. A small number of experimental trials²⁴
340 and many observational studies^{8,10} have reported increases in child growth and reductions in
341 stunting prevalence with improvements in sanitation coverage and commensurate reductions in
342 OD; among the latter, unmeasured confounding is a likely explanation for observed effects²⁴.

343
344 While gains in the proportion of the population self-reporting access to sanitation were highest in
345 the sanitation only arm (+25 pp), these were only modestly higher than the gains for the control
346 (+19 pp) and nutrition arms (+14 pp); furthermore, sanitation coverage gains for the combined
347 intervention matched the control arm at +19 pp. Comparable secular trends of increasing
348 sanitation coverage in Cambodia have been documented previously: the percentage of children
349 younger than five years of age with access to an improved sanitation facility increased from 5%
350 in 2000 to 17% in 2005, 29% in 2010, and 54% in 2014, the most recent nationally
351 representative DHS survey^{8,23}. Correspondingly swift improvements have been documented

352 specifically in rural areas, where access to any sanitation facility increased from 30% in 2010 to
353 44% in 2014 and improved sanitation coverage rose from 27% to 43%^{8,23}. The rapid pace of
354 WASH development in rural Cambodia makes it challenging to measure the impact of specific
355 programs. However, the lack of differences in other sanitation intervention adherence indicators
356 suggests low overall uptake of the sanitation intervention and only modest increases in sanitation
357 coverage attributable to the intervention, which were likely insufficient to reduce community
358 exposure and transmission. Zoonotic transmission from domestic animals, for instance, was not
359 addressed by this or many other WASH trials, as indicated by nearly 80% of households across
360 all treatment arms lacking access to an area free of animals for children to safely play²⁵.

361
362 The frequency and intensity of contact from program promoters was much greater in the
363 nutrition intervention than the sanitation intervention. Recipients of the nutrition intervention
364 participated in monthly activities, whereas the sanitation intervention consisted of one triggering
365 session with infrequent follow-up visits. The lower contact frequency may explain the
366 discrepancy in intervention adherence. Both arms receiving the nutrition programming reported
367 higher levels of participation in the key intervention activities—including sanitation intervention
368 activities—suggesting higher adherence to the nutrition intervention than the sanitation
369 intervention. Self-reported CLTS participation rates were equally low in both the sanitation-only
370 and control arms at 6%, while 14% of nutrition-only recipients and 25% of combined
371 intervention recipients reported CLTS participation. The comparatively elevated CLTS
372 participation in the nutrition-only arm may reflect biases embedded in the self-reporting process;
373 given the 28+ months that had elapsed since the initial CLTS triggering session and the
374 infrequency of CLTS follow-up visits, households that only received the sanitation intervention
375 may have been less likely to recall programming of any kind than households that participated in
376 the more frequent and intense nutrition intervention activities. Furthermore, the “Growth
377 Together” SBCC campaign, which promoted 13 core health, nutrition, sanitation, and hygiene
378 practices, was fully incorporated into all intervention activities across the three intervention
379 arms, meaning households receiving the higher intensity nutrition programming also encountered
380 the associated SBCC sanitation messaging much more frequently than households in the
381 sanitation-only arm. The SBCC campaign was also promoted nationally on television, such that
382 households in the control arms may have been nearly as exposed to its content as sanitation-only
383 households, while the nutrition-only and combined intervention arms received substantial in-
384 person promotion of the SBCC campaign messaging.

385
386 Due to the nature of the interventions and resource considerations, all trial outcomes were
387 assessed during a single survey round conducted 28 months after initiating the intervention
388 programming, which introduced some limitations. Growth and pathogen outcomes were assessed
389 in children from one to 28 months of age, meaning that older children received the treatments for
390 a longer duration but were initially exposed to less mature intervention conditions than the
391 younger children born later. The timing of outcome ascertainment also precluded detecting
392 effects that may only be realized later in childhood, such as potentially rapid catch-up growth
393 after 24 months of age that may reverse earlier growth faltering²⁶. While a focus on the first
394 1,000 days is justified², investigation of growth and growth-promoting factors after this window
395 may provide additional insight on improved WASH practices and their role in supporting long-
396 term development and health.

397

398 While molecular detection of a specific pathogen in stool unambiguously indicates prior
399 exposure to that pathogen, our data do not necessarily indicate active enteric infection, potential
400 for disease, or direct effects on the individual. We were limited by the suite of pathogen targets
401 selected on our custom TAC assay; we selected these targets *a priori* based on a range of
402 globally observed enteric pathogens, but we cannot know whether these were the most important
403 enteric pathogens in rural Cambodia. There is also evidence that the *invA* gene, which was
404 selected for *Salmonella* spp. detection, is not specific to *Salmonella enterica* and suggest the
405 consideration of other genes, such as *ttrA/C*, for reliable detection of *S. enterica*²⁷.
406 It is highly plausible that this sanitation intervention simply failed to sufficiently reduce
407 environmental exposure to enteric pathogens. For example, only 22% of households in our
408 survey were observed to have a child play environment free of animals, with little difference
409 between treatment and control arms; this is a transmission pathway that our trial and many other
410 WASH trials have not addressed²⁵. The trial design is predicated on the theory that gains in
411 sanitation coverage may lead to improved growth outcomes in children via reductions in the
412 transmission of enteric infection and disease, though links between sanitation coverage and
413 specific outcomes are poorly understood in high-burden settings. The change in community
414 coverage in this trial was limited and likely insufficient to reduce community exposure and
415 transmission.

416
417 There are a few key observations from this study that should be considered in future
418 interventions and effectiveness trials of comparable interventions. Increased frequency, duration,
419 and intensity of CLTS programming could have resulted in greater uptake of sanitation in target
420 communes. Despite the sanitation coverage gains observed in the sanitation-only arm, much of
421 which may have been as a result of the sanitation intervention, we are unable to attribute
422 beneficial effects—i.e., measurable differences in prespecified outcomes—to the sanitation
423 intervention due to the high sanitation gains also observed in the control arm. There may have
424 been other benefits of sanitation gains that were not measured, including in safety and broader
425 measures of well-being²⁸. Future trials may also include additional objective outcome measures,
426 including intermediate measures of environmental contamination that are on the causal pathway
427 between interventions and exposures.

428
429 Our work is consistent with a growing body of research reporting high prevalence of enteric
430 pathogen exposure in early childhood, which may lead to long-term effects on health^{15,18,29,30}.
431 Reducing these exposures in high-burden settings requires transformative interventions that have
432 the potential to dramatically reduce direct and indirect contact with all faeces, including animal
433 faeces²⁴, across multiple pathways.
434

435 Other Information

436 Trial Registry

437 The trial is registered with ISRCTN Registry ([ISRCTN77820875](https://www.isrctn.com/ISRCTN77820875)).

438 Protocol

439 The National Ethics Committee for Health Research in the Cambodian Ministry of Health
440 reviewed and approved the protocols (NECHR #110) prior to the start of data collection. The
441 study also received approvals from the Institutional Review Board at Georgia Institute of
442 Technology (Ref: H19286) and from New England IRB (IRB#: 120190186).

443 Funding

444 This study was funded by the United States Agency for International Development (USAID)
445 under contract number OAA-M-13-00017. The contents of this publication are the sole
446 responsibility of the authors and do not necessarily reflect the views of USAID or the United
447 States Government.

448 Acknowledgements


449 We thank the staff and participants of the study for their important contributions. We also
450 acknowledge contributions by Caroline Akerley, Farran Bush, Isabelle du Plessis, and Juliann
451 Pham.

452 Author contributions

Conceptualisation	IV, OC, JB
Data Curation	IV, KS, AL
Formal Analysis	AL, RA, IV
Funding Acquisition	IV, JB
Investigation	AL, KS, IV, JB
Methodology	AL, IV, OC, JB
Project Administration	IV, KS, JB
Resources	IV, JB
Software	N/A
Supervision	IV, KS, OC, JB
Validation	N/A
Visualisation	N/A
Writing – Original Draft Preparation	AL
Writing – Review & Editing	AL, IV, RA, EK, DH, KS, OC, JB, KL


453

454



455
456 **Figure 1: Theory of change diagram**


457
458
459
460



461
462 **Figure 2: Trial profile**

463


It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .



464
465
466
467

Figure 3: Unadjusted intervention effects on LAZ. Estimates are mean differences (point) with 95% CIs (line)

468 **Figure 4: Impact of interventions on unadjusted prevalence ratio of individual pathogens.**
 469 **Point estimates and 95% confidence intervals were determined using log-linear Poisson**
 470 **models with generalized estimating equations.**



471
472

473 **Table 1: Summary of intervention activities**

	Intervention activity	Frequency
Nutrition-only		
	Community dialogues led by village chief and VHSG to support children's growth	Quarterly
	Caregiver group sessions led by local women trained by staff to promote 13 key stunting prevention behaviours	Monthly
	GMP sessions led by VHSGs to monitor growth and refer children who were sick or not growing well to health centers	Monthly
	Home visits to pregnant women, caregivers of children 9-11 months old, and caregivers of children not growing well to promote childcare and feeding, home hygiene, and handwashing	Monthly
	Village fair help twice per year to offer hands-on learning experiences (health/nutrition, WASH and agricultural using games, latrine marketing and sales	Twice per year
	CCT (cash for antenatal and postnatal care visits and adherence to handwashing stations), vouchers for water filters and food baskets	Up to six payments over first 1,000 days of child life
Sanitation-only		
	CLTS triggering session	Once
	Door-to-door visits to provide information about sanitation/latrines	At least five times per village
	Latrine vouchers to subsidise poor households in villages that reached 75% sanitation coverage to achieve sufficient open defecation free (ODF) coverage	Once, as needed
	Promoted supply-side support by connecting small- and medium-sized enterprises (SMEs) with communes after triggering event	Continuously
Combined		
	All activities described in NUTR and SAN groups above	See above
Control		
	None	N/A

474

475

476 **Table 2: Household and caregiver characteristics**

	Nutrition-only		Sanitation-only		Combined		Control	
	Mean	95% CI	Mean	95% CI	Mean	95% CI	Mean	95% CI
Buddhist	0.98	(0.97, 0.99)	0.97	(0.96, 0.98)	0.96	(0.95, 0.97)	0.99	(0.98, 0.99)
Married or living together	0.95	(0.93, 0.96)	0.94	(0.92, 0.95)	0.94	(0.92, 0.95)	0.95	(0.94, 0.96)
Maternal age (years)	32.0	(31.2, 32.7)	31.1	(30.4, 31.9)	31.9	(31.2, 32.5)	31.0	(30.5, 31.5)
Primary caregiver has attended primary school	0.87	(0.85, 0.89)	0.87	(0.84, 0.89)	0.88	(0.85, 0.89)	0.81	(0.79, 0.83)
Spouse has attended primary school	0.90	(0.88, 0.92)	0.86	(0.83, 0.88)	0.87	(0.85, 0.89)	0.87	(0.85, 0.89)
Household size	5.63	(5.48, 5.78)	5.54	(5.40, 5.69)	5.36	(5.24, 5.47)	5.52	(5.42, 5.62)
Number of children in HH	2.61	(2.52, 2.70)	2.61	(2.51, 2.70)	2.43	(2.36, 2.50)	2.43	(2.37, 2.49)
Has electricity	0.73	(0.70, 0.76)	0.74	(0.71, 0.77)	0.76	(0.73, 0.78)	0.75	(0.72, 0.77)
Owns a mobile phone	0.93	(0.92, 0.95)	0.90	(0.88, 0.92)	0.87	(0.85, 0.89)	0.90	(0.89, 0.92)
Has a finished floor [1]	0.92	(0.90, 0.94)	0.93	(0.91, 0.95)	0.96	(0.94, 0.97)	0.96	(0.95, 0.97)
Wealth index score, excluding WASH variables	0.06	(-0.06, 0.19)	0.19	(0.05, 0.33)	-0.02	(-0.14, 0.09)	-0.12	(-0.21, -0.03)
Improved drinking water source [2]	0.59	(0.56, 0.62)	0.56	(0.52, 0.59)	0.61	(0.58, 0.64)	0.70	(0.68, 0.72)
Has water source on site	0.68	(0.64, 0.71)	0.58	(0.55, 0.61)	0.60	(0.57, 0.63)	0.65	(0.62, 0.67)
Water source is <5 min, roundtrip	0.20	(0.16, 0.25)	0.23	(0.19, 0.28)	0.24	(0.19, 0.29)	0.34	(0.29, 0.39)
Reported minutes to fetch water, roundtrip	12.42	(10.6, 14.2)	14.14	(12.5, 15.8)	14.50	(13.1, 15.9)	13.56	(12.3, 14.8)

477 [1] Finished floor defined as floor made of wood plans, palm/bamboo, parquet or polished wood, vinyl or asphalt strips, ceramic
 478 tiles, cement tiles, or cement. Floor materials were classified by enumerator observation. [2] Improved sources of drinking water
 479 include: piped water into dwelling/yard/plot, public tap or standpipe, tube well or borehole, protected dug well, protected spring,
 480 bottled water, and rainwater.

481
 482

483 **Table 3: Intervention fidelity indicators**

Nutrition	Nutrition-only			Sanitation-only			Combined			Control		
	N	n	%	N	n	%	N	n	%	N	n	%
Participated in any "First 1,000 Days" type activity [1]	817	615	75%	792	145	18%	1,055	813	77%	1,460	383	26%
Participated in any GMP	817	641	78%	792	181	23%	1,055	935	89%	1,460	482	33%
Received home visit VHSG	817	518	63%	792	227	29%	1,055	661	63%	1,460	490	34%
Enrolled in any CCT program for health and nutrition [2]	817	224	27%	792	19	2%	1,055	228	22%	1,460	31	2%
Received any voucher for food basket [3]	817	440	54%	792	1	0%	1,055	554	53%	1,460	6	0%
Received any voucher for water filter [3]	817	41	5%	792	40	5%	1,055	149	14%	1,460	100	7%
Aware of <i>Grow Together</i> campaign [4]	817	353	43%	792	93	12%	1,055	471	45%	1,460	149	10%
Participation in nutrition intervention activities: none (0 of 8)	817	65	8%	792	404	51%	1,055	44	4%	1,460	585	40%
Participation in nutrition intervention activities: low (1-3 of 8)	817	262	32%	792	359	45%	1,055	367	35%	1,460	809	55%
Participation in nutrition intervention activities: med (4-6 of 8)	817	387	47%	792	29	4%	1,055	501	47%	1,460	65	4%
Participation in nutrition intervention activities: high (7-8 of 8)	817	103	13%	792	0	0%	1,055	143	14%	1,460	1	0%
Sanitation												
Any CLTS participation [5]	817	115	14%	792	46	6%	1,055	261	25%	1,460	81	6%
Received any voucher to build latrine [6]	817	66	8%	792	51	6%	1,055	123	12%	1,460	84	6%
[1] "First 1,000 Days" activities were administered in nutrition-only and combined arms and include: community dialogues, caregiver group education sessions, and village fairs. [2] CCT program ended in Jan 2019 and the Government of Cambodia started a new CCT program in July 2019 across study area. CCT program administered in nutrition-only and combined arms only. [3] Vouchers for water filter and food baskets were targeted subsidies distributed to CCT participants in nutrition-only and combined arms. [4] <i>Grow Together</i> campaign was part of the nutrition programming (nutrition-only and combined arms). However, three TV spots were seen across all four arms. [5] In sanitation-only and combined arms, the Ministry of Rural Development confirmed that the project was the only CLTS campaign active in those areas. [6] Latrine vouchers were targeted subsidies given to households in villages that reached 75% sanitation coverage in sanitation-only and combined arms.												

484

485

486 **Table 4: Intervention adherence indicators (28-months after intervention)**

Nutrition	Nutrition-only			Sanitation-only			Combined			Control		
	N	n or mean	% or SD	N	n or mean	% or SD	N	n or mean	% or SD	N	n or mean	% or SD
Visited health facility for at least four antenatal care check-ups	697	632	91%	712	646	91%	910	819	90%	1,257	1116	89%
Brought child for monthly GMP at community or health center	817	641	78%	792	181	23%	1,055	935	89%	1,460	482	33%
Breastfeeding exclusively for children <6 months	171	109	64%	161	110	68%	205	140	68%	272	182	67%
Ever breastfed (all children)	817	797	98%	792	765	97%	1,048	1021	97%	1,457	1420	97%
Solid and semi-solid foods eaten for children >6 months	646	609	94%	631	611	97%	843	792	94%	1,185	1134	96%
Dietary diversity score (0-7)	817	2.24	1.61	792	2.19	1.58	1,048	2.20	1.62	1,457	2.33	1.59
Achieved minimum dietary diversity	817	202	25%	792	177	22%	1,048	247	24%	1,457	350	24%
Achieved minimum meal frequency	817	537	66%	792	518	65%	1,048	680	65%	1,457	977	67%
Achieved minimum acceptable diet	817	170	21%	792	159	20%	1,048	205	20%	1,457	310	21%
Treated drinking water	817	548	67%	792	471	59%	1,055	765	73%	1,460	1,037	71%
Treated drinking water with filter	817	151	18%	792	160	20%	1,055	302	29%	1,460	548	40%
Sanitation												
Had improved sanitation facility [1]	816	452	55%	791	400	51%	1,054	638	61%	1,459	759	52%
Open defecation (OD)	817	112	14%	792	126	16%	1,055	73	7%	1,460	231	16%
Used shared toilet	817	252	31%	791	264	33%	1,054	343	33%	1,459	468	32%
Caregiver reported adults in HH openly defecating	697	92	13%	658	116	18%	973	118	12%	1,208	213	18%
Time to get to toilet, one way (minutes)	171	4.22	4.11	166	3.92	3.83	219	4.74	8.17	291	5.05	8.27
Reported latrine built as a result of CLTS activity	115	51	44%	46	15	33%	261	91	35%	81	28	35%
Reported latrine built using latrine voucher	50	10	20%	15	4	27%	91	37	41%	28	12	43%
Main reason for not constructing latrine: lack of funds	20	17	85%	18	14	78%	60	55	92%	14	14	100%
Main reason for construction latrine: privacy	51	7	14%	15	6	40%	91	6	7%	28	7	25%
Main reason for construction latrine: security	51	10	20%	15	2	13%	91	20	22%	28	4	14%
Main reason for construction latrine: hygiene	51	17	33%	15	5	33%	91	43	47%	28	10	36%
Main reason for construction latrine: OD is harmful	51	5	10%	15	1	7%	91	9	10%	28	5	18%

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .

Child stools properly disposed of [2]	817	581	71%	792	515	65%	1,055	781	74%	1,460	993	68%
Community-level open defecation before intervention	817	28%	27%	792	41%	33%	1,055	28%	28%	1,460	32%	31%
Community-level open defecation after intervention	817	14%	21%	792	16%	21%	1,055	9%	17%	1,460	13%	20%
Environmental hygiene												
Child stools left in the open	817	147	18%	792	170	21%	1,055	160	15%	1,460	315	22%
Child play environment observed to be free of animals	817	182	22%	792	187	24%	1,055	261	25%	1,460	294	20%
Child play environment observed to be free of garbage/HH waste	817	298	36%	792	290	37%	1,055	419	40%	1,460	567	39%
Child play environment observed to be free of sharp objects	817	449	55%	792	427	54%	1,055	639	61%	1,460	818	56%
Child play environment observed to be free of faeces	817	313	38%	792	304	38%	1,055	448	42%	1,460	555	38%
[1] Improved sanitation facilities include: flush/pour flush toilet to a piped sewer system, septic tank or pit latrine, a ventilated improved pit latrine, a pit latrine with slab, and a composting toilet. [2] Proper disposal of children faeces consist of putting or rinsing stool into a sanitation facility or burying it; improper disposal of children faeces includes putting or rinsing stool into a drain or ditch, throwing it into garbage or leaving it in the open.												

487

488

489 **Table 5: Effects of interventions on length and weight (Primary outcome (LAZ) and**
 490 **secondary outcomes (WAZ, WLZ)), comparing intervention arms to control and single**
 491 **intervention arms to combined intervention**

				Unadjusted mean difference (95% CI)	
	N	Mean	SD	Compared to control arm	Compared to combined intervention arm
LAZ					
Nutrition-only	798	-0.95	1.16	0.08 (-0.01, 0.18)	-0.02 (-0.12, 0.09)
Sanitation-only	777	-1.09	1.23	-0.05 (-0.16, 0.05)	-0.16 (-0.27, -0.04)
Combined	1037	-0.94	1.16	0.10 (0.01, 0.20)	--
Control	1443	-1.04	1.20	--	--
WAZ					
Nutrition-only	815	-0.95	1.29	0.10 (0.00, 0.19)	-0.02 (-0.12, 0.08)
Sanitation-only	792	-1.04	1.13	0.01 (-0.07, 0.09)	-0.10 (-0.20, -0.01)
Combined	1044	-0.94	1.11	0.11 (0.03, 0.20)	--
Control	1452	-1.05	1.10	--	--
WLZ					
Nutrition-only	814	-0.60	1.04	0.06 (-0.03, 0.15)	-0.02 (-0.12, 0.08)
Sanitation-only	790	-0.59	0.98	0.06 (-0.02, 0.14)	-0.02 (-0.11, 0.07)
Combined	1043	-0.58	1.03	0.08 (0.00, 0.16)	--
Control	1452	-0.65	0.98	--	--

492
 493
 494

495 **Table 6: Effects of intervention on child health outcomes, comparing intervention arms to**
 496 **control and single intervention arms to combined intervention.**

				Compared to control arm	Compared to combined-intervention arm
	N	Mean	SD	PR (95% CI)	PR (95% CI)
Stunted					
Nutrition-only	801	0.15	0.36	0.84 (0.69, 1.03)	0.93 (0.74, 1.15)
Sanitation-only	782	0.21	0.40	1.12 (0.94, 1.33)	1.23 (1.02, 1.49)
Combined	1046	0.17	0.37	0.91 (0.76, 1.09)	--
Control	1449	0.18	0.39	--	--
Wasted					
Nutrition-only	815	0.07	0.26	0.87 (0.65, 1.17)	1.12 (0.80, 1.57)
Sanitation-only	790	0.07	0.26	0.84 (0.62, 1.14)	1.08 (0.76, 1.53)
Combined	1052	0.07	0.25	0.78 (0.58, 1.04)	--
Control	1457	0.08	0.28	--	--
Underweight					
Nutrition-only	816	0.15	0.35	0.85 (0.71, 1.03)	1.04 (0.84, 1.29)
Sanitation-only	792	0.17	0.38	1.00 (0.85, 1.19)	1.22 (1.00, 1.49)
Combined	1053	0.14	0.35	0.82 (0.68, 0.99)	--
Control	1457	0.17	0.38	--	--
Diarrhoea (7-day recall)					
Nutrition-only	788	0.19	0.39	0.89 (0.74, 1.06)	0.95 (0.78, 1.14)
Sanitation-only	752	0.21	0.41	0.99 (0.84, 1.17)	1.05 (0.88, 1.25)
Combined	1018	0.20	0.40	0.94 (0.80, 1.11)	--
Control	1411	0.21	0.41	--	--
All-cause mortality					
Nutrition-only	1574	0.03	0.16	1.55 (0.71, 3.39)	1.61 (0.68, 3.82)
Sanitation-only	1636	0.03	0.16	1.09 (0.50, 2.40)	1.13 (0.48, 2.68)
Combined	1932	0.03	0.16	0.96 (0.44, 2.10)	--
Control	2688	0.03	0.16	--	--

497

498

499 **Table 7: Enteric pathogen gene prevalence among treatment arms**

	All samples (N=1620)	Nutrition- only (N=305)	Sanitation- only (N=333)	Combined (N=438)	Control (N=544)
Bacteria					
<i>Campylobacter</i> spp.	551 (34%)	104 (34%)	114 (34%)	162 (37%)	171 (31%)
<i>Clostridium difficile</i>	139 (9%)	33 (11%)	25 (8%)	41 (9%)	40 (7%)
EAEC	1029 (64%)	204 (67%)	207 (62%)	281 (64%)	337 (62%)
EPEC	899 (55%)	172 (56%)	173 (52%)	234 (53%)	320 (59%)
aEPEC	703 (43%)	139 (46%)	137 (41%)	173 (39%)	254 (47%)
tEPEC	109 (7%)	15 (5%)	17 (5%)	32 (7%)	45 (8%)
ETEC	422 (26%)	86 (28%)	79 (24%)	114 (26%)	143 (26%)
ETEC-LT	342 (21%)	75 (25%)	68 (20%)	86 (20%)	113 (21%)
ETEC-ST	194 (12%)	39 (13%)	37 (11%)	63 (14%)	55 (10%)
ETEC-LT/ST	114 (7%)	28 (9%)	26 (8%)	35 (8%)	25 (5%)
<i>Salmonella</i> spp.	134 (8%)	28 (9%)	19 (6%)	39 (9%)	48 (9%)
<i>Shigella</i> spp.	186 (11%)	24 (8%)	39 (12%)	52 (12%)	71 (13%)
STEC	132 (8%)	27 (9%)	22 (7%)	45 (10%)	38 (7%)
<i>Vibrio cholerae</i>	10 (1%)	1 (0%)	6 (2%)	1 (0%)	2 (0%)
At least 1 bacterium detected	1410 (87%)	276 (90%)	282 (85%)	386 (88%)	466 (86%)
Mean number of bacteria detected	2.48 (2.42, 2.55)	2.46 (2.32, 2.60)	2.43 (2.28, 2.57)	2.51 (2.39, 2.63)	2.51 (2.40, 2.62)
Viruses					
Adenovirus	287 (18%)	77 (25%)	59 (18%)	78 (18%)	73 (13%)
Astrovirus	7 (0%)	2 (1%)	1 (0%)	3 (1%)	1 (0%)
Enterovirus	558 (34%)	97 (32%)	105 (32%)	169 (39%)	187 (34%)
Norovirus	54 (3%)	8 (3%)	9 (3%)	20 (5%)	17 (3%)
Rotavirus	17 (1%)	2 (1%)	2 (1%)	6 (1%)	7 (1%)
Sapovirus	24 (1%)	4 (1%)	5 (2%)	9 (2%)	6 (1%)
At least 1 virus detected	788 (49%)	152 (50%)	157 (47%)	231 (53%)	248 (46%)
Mean number of viruses detected	1.20 (1.17, 1.23)	1.25 (1.17, 1.33)	1.15 (1.09, 1.21)	1.23 (1.17, 1.30)	1.17 (1.12, 1.23)
Protozoa					
<i>Cryptosporidium</i>	17 (1%)	4 (1%)	1 (0%)	6 (1%)	6 (1%)
<i>Entamoeba</i>	13 (1%)	1 (0%)	1 (0%)	8 (2%)	3 (1%)
<i>Giardia</i>	306 (19%)	52 (17%)	68 (20%)	84 (19%)	102 (19%)
At least 1 protozoan detected	328 (20%)	56 (18%)	69 (21%)	93 (21%)	110 (20%)
Mean number of protozoa detected	1.02 (1.01, 1.04)	1.02 (0.98, 1.05)	1.01 (0.99, 1.04)	1.05 (1.01, 1.10)	1.01 (0.99, 1.03)
STH					
<i>Ascaris lumbricoides</i>	3 (0%)	0 (0%)	0 (0%)	3 (1%)	0 (0%)
<i>Trichuris trichiura</i>	3 (0%)	1 (0%)	1 (0%)	0 (0%)	1 (0%)
<i>Ancylostoma duodenale</i>	17 (1%)	0 (0%)	4 (1%)	4 (1%)	9 (2%)
<i>Necator americanus</i>	20 (1%)	3 (1%)	6 (2%)	5 (1%)	6 (1%)
At least 1	37 (2%)	4 (1%)	10 (3%)	9 (2%)	14 (3%)
Mean number of STH detected	1.16 (1.04, 1.29)	1.00 (1.00, 1.00)	1.10 (0.90, 1.30)	1.33 (1.00, 1.67)	1.14 (0.95, 1.34)

500
501

502 **Table 8: Unadjusted incidence rate ratios of co-detected bacteria, viruses, protozoa, and**
 503 **STHs, comparing intervention arms to control and single intervention arms to combined**
 504 **intervention.**

	N	Compared to control arm			Compared to combined arm	
		Nutrition-only	Sanitation-only	Combined	Nutrition-only	Sanitation-only
Bacteria	1620	1.04 (0.95, 1.13)	0.96 (0.87, 1.05)	1.03 (0.95, 1.12)	0.01 (-0.08, 0.10)	-0.07 (-0.17, 0.02)
Viruses	1620	1.16 (0.99, 1.37)	1.02 (0.86, 1.19)	1.22 (1.05, 1.41)	-0.04 (-0.21, 0.12)	-0.18 (-0.34, -0.02)
Protozoa	1620	0.92 (0.68, 1.23)	1.03 (0.79, 1.35)	1.10 (0.85, 1.41)	-0.18 (-0.48, 0.12)	-0.06 (0.66, -0.34)

STHs were omitted because <5% of samples had detectable STH genes.

505 **Table 9: Unadjusted prevalence ratios (PR) of detected bacteria, viruses, protozoa, and**
 506 **STHs, comparing intervention arms to control and single intervention arms to combined**
 507 **intervention.**
 508

	Compared to control arm			Compared to combined arm	
	Nutrition-only	Sanitation-only	Combined	Nutrition-only	Sanitation-only
Any bacterium	1.06 (1.00, 1.11)	0.99 (0.93, 1.05)	1.03 (0.98, 1.08)	1.03 (0.98, 1.08)	0.96 (0.91, 1.02)
<i>Campylobacter</i> spp.	1.08 (0.89, 1.32)	1.09 (0.90, 1.32)	1.18 (0.99, 1.40)	0.92 (0.76, 1.12)	0.93 (0.76, 1.12)
<i>C.diff</i>	1.47 (0.95, 2.28)	1.02 (0.63, 1.65)	1.27 (0.84, 1.93)	1.16 (0.75, 1.78)	0.80 (0.50, 1.29)
EAEC	1.08 (0.97, 1.20)	1.00 (0.90, 1.12)	1.04 (0.94, 1.14)	1.04 (0.94, 1.16)	0.97 (0.87, 1.08)
EPEC	0.96 (0.85, 1.08)	0.88 (0.78, 1.00)	0.91 (0.81, 1.02)	1.06 (0.93, 1.20)	0.97 (0.85, 1.11)
aEPEC	0.98 (0.84, 1.14)	0.88 (0.75, 1.03)	0.85 (0.73, 0.98)	1.15 (0.97, 1.37)	1.04 (0.88, 1.24)
tEPEC	0.59 (0.34, 1.05)	0.62 (0.36, 1.06)	0.88 (0.57, 1.37)	0.67 (0.37, 1.22)	0.70 (0.39, 1.24)
ETEC	1.07 (0.85, 1.35)	0.90 (0.71, 1.15)	0.99 (0.80, 1.22)	1.08 (0.85, 1.38)	0.91 (0.71, 1.17)
ETEC-LT	1.18 (0.92, 1.53)	0.98 (0.75, 1.29)	0.95 (0.74, 1.21)	1.25 (0.95, 1.65)	1.04 (0.78, 1.38)
ETEC-ST	1.26 (0.86, 1.86)	1.10 (0.74, 1.63)	1.42 (1.01, 2.00)	0.89 (0.61, 1.29)	0.77 (0.53, 1.13)
ETEC-LT/ST	2.00 (1.19, 3.36)	1.70 (1.00, 2.89)	1.74 (1.06, 2.86)	1.15 (0.71, 1.85)	0.98 (0.60, 1.59)
<i>Salmonella</i> spp.	1.04 (0.67, 1.62)	0.65 (0.39, 1.08)	1.01 (0.67, 1.51)	1.03 (0.65, 1.64)	0.64 (0.38, 1.09)
EIEC/ <i>Shigella</i> spp.	0.60 (0.39, 0.94)	0.90 (0.62, 1.29)	0.91 (0.65, 1.27)	0.66 (0.42, 1.05)	0.99 (0.67, 1.46)
STEC	1.27 (0.79, 2.03)	0.95 (0.57, 1.57)	1.47 (0.97, 2.22)	0.86 (0.55, 1.36)	0.64 (0.39, 1.05)
Any virus	1.09 (0.95, 1.26)	1.03 (0.89, 1.20)	1.16 (1.02, 1.31)	0.94 (0.82, 1.09)	0.89 (0.77, 1.03)
Adenovirus	1.88 (1.41, 2.51)	1.32 (0.96, 1.81)	1.33 (0.99, 1.78)	1.42 (1.07, 1.87)	0.99 (0.73, 1.35)
Enterovirus	0.93 (0.76, 1.13)	0.92 (0.75, 1.12)	1.12 (0.95, 1.32)	0.82 (0.67, 1.01)	0.82 (0.67, 1.00)
Any protozoa	0.91 (0.68, 1.21)	1.02 (0.78, 1.34)	1.05 (0.82, 1.34)	0.86 (0.64, 1.16)	0.98 (0.74, 1.29)
<i>Giardia</i>	0.91 (0.67, 1.23)	1.09 (0.83, 1.43)	1.02 (0.79, 1.33)	0.89 (0.65, 1.22)	1.06 (0.80, 1.42)

Gene targets with <5% prevalence were omitted from PR analyses: *V.cholera*, astrovirus, norovirus, rotavirus, sapovirus, *Cryptosporidium*, *Entamoeba*, and all STHs (*Ascaris*, *Trichuris*, *Ancylostoma*, and *Necator*).

509
510

511 **Table 10: Mean difference in log₁₀-transformed gene copy estimates, comparing**
 512 **intervention arms to control and single intervention arms to combined intervention.**

	Compared to control arm			Compared to combined arm	
	Nutrition-only	Sanitation-only	Combined	Nutrition-only	Sanitation-only
Bacteria					
<i>CAMP</i>	0.22 (-0.60 - 1.03)	-0.40 (-1.27 - 0.48)	-0.24 (-1.00 - 0.53)	0.45 (-0.37 - 1.28)	-0.16 (-1.05 - 0.73)
<i>CDIF</i>	0.31 (-1.14 - 1.75)	-0.37 (-1.92 - 1.19)	0.19 (-1.16 - 1.54)	0.11 (-1.29 - 1.52)	-0.56 (-2.08 - 0.96)
<i>EAEC aaic</i>	-0.38 (-1.35 - 0.60)	0.35 (-0.52 - 1.22)	-0.28 (-1.08 - 0.52)	-0.10 (-1.11 - 0.92)	0.63 (-0.28 - 1.54)
<i>EAEC aata</i>	-0.36 (-1.26 - 0.53)	-0.24 (-1.05 - 0.56)	0.12 (-0.69 - 0.93)	-0.48 (-1.43 - 0.47)	-0.36 (-1.24 - 0.51)
<i>EPEC bfpA</i>	-1.08 (-3.47 - 1.31)	-1.13 (-3.16 - 0.89)	1.15 (-0.38 - 2.67)	-2.23 (-4.57 - 0.11)	-2.28 (-4.24 - -0.32)
<i>EPEC eae</i>	-0.12 (-0.80 - 0.56)	-0.54 (-1.17 - 0.09)	0.24 (-0.33 - 0.82)	-0.37 (-1.05 - 0.32)	-0.78 (-1.41 - -0.15)
<i>ETEC LT</i>	-0.62 (-1.64 - 0.40)	-0.47 (-1.56 - 0.61)	-0.03 (-1.06 - 1.00)	-0.59 (-1.65 - 0.47)	-0.44 (-1.57 - 0.68)
<i>ETEC stp</i>	0.13 (-1.73 - 1.98)	1.29 (-0.61 - 3.20)	0.87 (-0.97 - 2.71)	-0.75 (-2.61 - 1.11)	0.42 (-1.49 - 2.33)
<i>SALM</i>	1.42 (-0.04 - 2.87)	0.27 (-0.97 - 1.52)	0.80 (-0.33 - 1.93)	0.62 (-0.93 - 2.17)	-0.53 (-1.89 - 0.83)
<i>IPAH</i>	0.22 (-1.33 - 1.77)	-0.28 (-1.68 - 1.13)	1.17 (0.07 - 2.26)	-0.95 (-2.43 - 0.53)	-1.44 (-2.77 - -0.11)
<i>STEC1</i>	-1.46 (-2.97 - 0.06)	-1.71 (-3.07 - -0.34)	-0.00 (-1.49 - 1.49)	-1.46 (-2.84 - -0.07)	-1.70 (-2.92 - -0.49)
<i>STEC2</i>	-0.20 (-1.45 - 1.04)	0.09 (-1.20 - 1.37)	0.72 (-0.47 - 1.91)	-0.92 (-2.06 - 0.22)	-0.63 (-1.82 - 0.56)
Viruses					
<i>ADEV</i>	0.50 (-0.48 - 1.49)	0.67 (-0.32 - 1.67)	0.48 (-0.42 - 1.38)	0.03 (-0.96 - 1.02)	0.20 (-0.80 - 1.20)
<i>ENTV</i>	-0.40 (-1.09 - 0.28)	-0.35 (-0.97 - 0.26)	-0.26 (-0.77 - 0.24)	-0.14 (-0.84 - 0.56)	-0.09 (-0.72 - 0.54)
Protozoa					
<i>GIAR</i>	-1.73 (-3.02 - -0.44)	0.23 (-1.16 - 1.62)	-0.14 (-1.47 - 1.18)	-1.58 (-3.06 - -0.11)	0.37 (-1.19 - 1.93)
Gene targets with <5% prevalence were omitted from PR analyses: <i>V.cholera</i> , astrovirus, norovirus, rotavirus, sapovirus, <i>Cryptosporidium</i> , <i>Entamoeba</i> , and all STHs (<i>Ascaris</i> , <i>Trichuris</i> , <i>Ancylostoma</i> , and <i>Necator</i>).					

513

References

1. Prendergast, A. J. & Humphrey, J. H. The stunting syndrome in developing countries. *Paediatr Int Child Health* **34**, 250–265 (2014).
2. Victora, C. G. *et al.* Maternal and child undernutrition: consequences for adult health and human capital. *The Lancet* **371**, 340–357 (2008).
3. Luby, S. P. *et al.* Effects of water quality, sanitation, handwashing, and nutritional interventions on diarrhoea and child growth in rural Bangladesh: a cluster randomised controlled trial. *Lancet Glob Health* **6**, e302–e315 (2018).
4. Null, C. *et al.* Effects of water quality, sanitation, handwashing, and nutritional interventions on diarrhoea and child growth in rural Kenya: a cluster-randomised controlled trial. *Lancet Glob Health* **6**, e316–e329 (2018).
5. Ruel, M. T. & Alderman, H. Nutrition-sensitive interventions and programmes: how can they help to accelerate progress in improving maternal and child nutrition? *Lancet* **382**, 536–551 (2013).
6. Humphrey, J. H. Child undernutrition, tropical enteropathy, toilets, and handwashing. *Lancet* **374**, 1032–1035 (2009).
7. Guerrant, R. L., Schorling, J. B., McAuliffe, J. F. & de Souza, M. A. Diarrhea as a cause and an effect of malnutrition: diarrhea prevents catch-up growth and malnutrition increases diarrhea frequency and duration. *Am J Trop Med Hyg* **47**, 28–35 (1992).
8. Ikeda, N., Irie, Y. & Shibuya, K. Determinants of reduced child stunting in Cambodia: analysis of pooled data from three demographic and health surveys. *Bull World Health Organ* **91**, 341–349 (2013).
9. Humphrey, J. H. *et al.* Independent and combined effects of improved water, sanitation, and hygiene, and improved complementary feeding, on child stunting and anaemia in rural Zimbabwe: a cluster-randomised trial. *Lancet Glob Health* **7**, e132–e147 (2019).
10. Vyas, S., Kov, P., Smets, S. & Spears, D. Disease externalities and net nutrition: Evidence from changes in sanitation and child height in Cambodia, 2005–2010. *Econ Hum Biol* **23**, 235–245 (2016).
11. Harris, M., Alzua, M. L., Osbert, N. & Pickering, A. Community-Level Sanitation Coverage More Strongly Associated with Child Growth and Household Drinking Water Quality than Access to a Private Toilet in Rural Mali. *Environ Sci Technol* **51**, 7219–7227 (2017).
12. Pickering, A. J., Djebbari, H., Lopez, C., Coulibaly, M. & Alzua, M. L. Effect of a community-led sanitation intervention on child diarrhoea and child growth in rural Mali: A cluster-randomised controlled trial. *Lancet Glob Health* **3**, e701–e711 (2015).
13. Briceño, B., Coville, A., Gertler, P. & Martinez, S. Are there synergies from combining hygiene and sanitation promotion campaigns: Evidence from a large-scale cluster-randomized trial in rural Tanzania. *PLoS One* **12**, e0186228–e0186228 (2017).
14. Clasen, T. *et al.* Effectiveness of a rural sanitation programme on diarrhoea, soil-transmitted helminth infection, and child malnutrition in Odisha, India: a cluster-randomised trial. *Lancet Glob Health* **2**, e645–e653 (2014).
15. Knee, J. *et al.* Effects of an urban sanitation intervention on childhood enteric infection and diarrhea in Maputo, Mozambique: A controlled before-and-after trial. *Elife* **10**, 2020.08.20.20178608 (2021).

16. Center for Disease Control and Prevention (CDC). *National Health and Nutrition Examination Survey (NHANES): Anthropometry Procedures Manual*. https://wwwn.cdc.gov/nchs/data/nhanes/2017-2018/manuals/2017_Anthropometry_Procedures_Manual.pdf (2017).
17. Mutz, D. C., Pemantle, R. & Pham, P. The Perils of Balance Testing in Experimental Design: Messy Analyses of Clean Data. *Am Stat* **73**, 32–42 (2019).
18. Liu, J. *et al.* Use of quantitative molecular diagnostic methods to identify causes of diarrhoea in children: a reanalysis of the GEMS case-control study. *The Lancet* **388**, 1291–1301 (2016).
19. Rothman, K. J. No Adjustments Are Needed for Multiple Comparisons. *Epidemiology* **1**, (1990).
20. Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal statistical society: series B (Methodological)* **57**, 289–300 (1995).
21. Marschner, I. C. Optimal design of clinical trials comparing several treatments with a control. *Pharm Stat* **6**, 23–33 (2007).
22. Zanello, G., Srinivasan, C. S. & Shankar, B. What Explains Cambodia’s Success in Reducing Child Stunting-2000-2014? *PLoS One* **11**, e0162668 (2016).
23. National Institute of Statistics, Directorate General for Health, and I. I. *Cambodia Demographic and Health Survey 2014*.
24. Pickering, A. J. *et al.* The WASH Benefits and SHINE trials: interpretation of WASH intervention effects on linear growth and diarrhoea. *Lancet Glob Health* **7**, e1139–e1146 (2019).
25. Prendergast, A. J. *et al.* Putting the ‘A’ into WaSH: a call for integrated management of water, animals, sanitation, and hygiene. *Lancet Planet Health* **3**, e336–e337 (2019).
26. Leroy, J. L., Ruel, M., Habicht, J.-P. & Frongillo, E. A. Using height-for-age differences (HAD) instead of height-for-age z-scores (HAZ) for the meaningful measurement of population-level catch-up in linear growth in children less than 5 years of age. *BMC Pediatr* **15**, 145 (2015).
27. Resendiz-Nava, C., Esquivel-Hernandez, Y., Alcaraz-Gonzalez, A., Castaneda-Serrano, P. & Nava, G. M. PCR Assays Based on invA Gene Amplification are not Reliable for Salmonella Detection. *Jundishapur J Microbiol* **In Press**, (2019).
28. Shiras, T. *et al.* Shared latrines in Maputo, Mozambique: exploring emotional well-being and psychosocial stress. *BMC Int Health Hum Rights* **18**, 30 (2018).
29. Platts-Mills, J. A. *et al.* Use of quantitative molecular diagnostic methods to assess the aetiology, burden, and clinical characteristics of diarrhoea in children in low-resource settings: a reanalysis of the MAL-ED cohort study. *Lancet Glob Health* **6**, e1309–e1318 (2018).
30. Rogawski, E. T. *et al.* Use of quantitative molecular diagnostic methods to investigate the effect of enteropathogen infections on linear growth in children in low-resource settings: longitudinal analysis of results from the MAL-ED cohort study. *Lancet Glob Health* **6**, e1319–e1328 (2018).