

## Supplementary material

**Table S1. Derivation of exposure variables used from Avon Longitudinal Study of Parents and Children (ALSPAC)**

Data collection	ALSPAC variable	Question(s)/measure(s)	Responses	Final variable coding
<b>Incontinence</b>				
ALSPAC 'travelling, leisure, and school' questionnaire	<b>How often does the following happen to you?</b>			
	ccp550	<i>Daytime wetting</i> Wet yourself during the day?	1: Never 2: Less than once per week 3: Once a week 4: 2-5-times a week 5: Nearly everyday 6: More than once a day -10, -1: Missing	Recode (1=0) (1/6=1) 0: No 1: Yes
	ccp551	<i>Bedwetting</i> Wet the bed at night?		
ccp552	<i>Soiling</i> Dirty your pants during the day?			
<b>Lower urinary tract symptoms</b>				
ALSPAC 'travelling, leisure, and school' questionnaire	<b>Over the last two weeks, how often have you:</b>			
	ccp520	<i>Urgency</i> Had the sudden feeling you need a wee and had to dash to the toilet?	1: Never 2: A few times 3: Quite often 4: A lot -10, -1: Missing	Recode (1,2=0) (3/4=1) 0: No 1: Yes
	ccp521	<i>Frequent urination</i> Had to go to the toilet for a wee more than seven times a day?		
	ccp522	<i>Low voided volume</i> Passed only a small amount when you went for a wee?		
	ccp524	<i>Voiding postponement</i> Avoided going for a wee until the last moment because you were concentrating on other activities?		
ccp527	<i>Nocturia</i> Woken up to go for a wee?			

**Table S2. Derivation of outcome variables used from Avon Longitudinal Study of Parents and Children (ALSPAC)**

Data collection	ALSPAC Variable	Question(s)/measure(s)	Responses	Final variable coding
<b>Common mental disorders</b>				
Clinical Interview Schedule-Revised* (CIS-R)*	FJCI050	Total CIS-R score	Continuous variable (range: 0-39)	Recode (<12=0) (≥12=1) 0: No 1: Yes
<b>Depression</b>				
Short Moods and Feelings Questionnaire*	CCXD917	High depressive symptoms	Continuous variable (range: 0-26) -10, -1: Missing	Recode (<11=0), (≥11=1) 0: No 1: Yes
CIS-R*	FJCI1001	ICD-10 diagnosis of depression (any severity: mild, moderate, or severe)	0: Not diagnosed 1: Diagnosed -10, -4, -1: Missing	0: No 1: Yes
<b>Anxiety Symptoms</b>				
CIS-R*	FJCI602	Presence of GAD symptoms	Binary variable 0: No 1: Yes -10, -4, -1: Missing	0: No 1: Yes
Anxiety Sensitivity Index*	FJLE220	Physical anxiety subscale score	Continuous variable (range: 10-50) -10, -4, -1: Missing	Continuous variable
	FJLE221	Mental anxiety subscale score	Continuous variable (range: 8-40) -10, -4, -1: Missing	Continuous variable
<b>Self-Harm</b>				
CIS-R*	FJCI369	Have you ever hurt yourself on purpose in any way (e.g., by taking an overdose of pills or by cutting yourself)?	1: No 2: Yes -10, -4, -1: Missing	If FJCI369 = 2, recode FJCI370 (1=0) (2/5=1): 0: No 1: Yes If FJCI369 = 1, recode FJCI370 (=0)
	FJCI370	How many times have you hurt yourself on purpose in the last year?	1: None 2: Once 3: 2-5 4: 6-10 5: >10 -10, -4, -1: Missing	
	FJCI371	During the past week, have you thought about hurting yourself on purpose?	1: No 2: Yes, but never commit suicide 3: Yes -10, -4, -1: Missing	Recode (1=0) (1/2=1) 0: No 1: Yes
<b>Excessive exercise</b>				
ALSPAC 'Your changing life' questionnaire*	cct4105	<b>During the past year:</b> How often did you do any exercise (going to the gym, brisk walking, or any other sports activity)?	1: 5 or more times a week 2: 1-4 times a week 3: 1-3 times a month 4: Less than once a month 5: Never	<b>Any excessive exercise:</b> <i>Exercise must be accompanied by intent to control weight and/or impairment.</i>  Recode cct4105 (5=0) (1/4=1) Recode cct4106 (3=0) (1/2=1) Recode cct4107 (3=0) (1/2=1) Recode cct4108 (0=0) (1/3=1)
	cct4106	Was it difficult for you to do your work or schoolwork because of the amount of time that you were exercising?	1: Yes, sometimes 2: Yes, frequently 3: No	

Supplementary material for: *Prospective relationships between continence problems and common mental health disorders in adolescents from a UK cohort*

Data collection	ALSPAC Variable	Question(s)/measure(s)	Responses	Final variable coding
	cct4107	Did you exercise in order to lose weight or avoid gaining weight?	1: Yes, sometimes 2: Yes, frequently 3: No	0: No (if cct41005, cct4106, cct41007 or cct4108 = 0) 1: Yes (if cct41007 = 1 & cct4108 = 1, or if cct41007 = 1 & cct4106 = 1)  <b>DSM-5<sup>†</sup> excessive exercise:</b>  Recode cct4105 (3/5=0) (1/2=1) Recode cct4106 (3=0) (1/2=1) Recode cct4107 (3=0) (1/2=1), replace = 0 if cct4105 = 0 Recode cct4108 (0/1=0) (2/3=1), replace = 0 if cct4105 = 0  0: (if cct41005, cct4106, cct41007 or cct4108 = 0) 1: Yes (if cct41007 = 1 & cct4108 = 1, or if cct41007 = 1 & cct4106 = 1)
	cct4108	Did you feel guilty after missing an exercise session?	0: No 1: Yes, sometimes 2: Yes, frequently 3: Did not miss any exercise sessions	
<b>Fasting</b>				
ALSPAC 'Your changing life' questionnaire	cct4110	During the past year, how often did you fast (not eat for at least a day) to lose weight or avoid gaining weight?	1: Never 2: Less than once a month 3: 1-3 times a month 4: Once a week 5: 2 or more times a week	<b>Any fasting:</b> Recode (1=0) (2/5=1) 0: No 1: Yes  <b>DSM-5<sup>†</sup> fasting:</b> Recode (1/3=0) (4/5=1) 0: No 1: Yes
<b>Purging</b>				
ALSPAC 'Your changing life' questionnaire	cct4112	How often did you make yourself throw up (vomit) to lose weight or avoid gaining weight?	1: Never 2: Less than once a month 3: 1-3 times a month 4: Once a week 5: 2-6 times a week 6: Every day	<b>Any purging</b> Recode cct4112 (1=0) (2/6=1) Recode cct4115 (3=0) (1/2=1) Recode cct4116 (1=0) (2/6=1), replace = 0 if cct4115 = 0
	cct4115	Did you take laxatives or other tablets or medicines (diet pills or water tablets) to lose weight or avoid gaining weight?	1: Yes, laxative 2: Yes, other 3: Never	0: No (if cct4112 & cct4116=0) 1: Yes (if cct4112 or cct4116=1)
	cct4116	How often did you take laxatives or other tablets or medicines to lose weight or avoid gaining weight?	1: Never 2: Less than once a month 3: 1-3 times a month 4: Once a week 5: 2-6 times a week 6: Every day	<b>DSM-5<sup>†</sup> purging</b> Recode cct4112 (1/3=0) (4/6=1) Recode cct4115 (3=0) (1/2=1) Recode cct4116 (1/3=0) (4/6=1), replace = 0 if cct4115 = 0  0: No (if cct4112 & cct4116=0) 1: Yes (if cct4112 or cct4116=1)
<b>Binge-eating</b>				
		<b>During the past year:</b>		

Supplementary material for: *Prospective relationships between continence problems and common mental health disorders in adolescents from a UK cohort*

Data collection	ALSPAC Variable	Question(s)/measure(s)	Responses	Final variable coding
ALSPAC 'Your changing life' questionnaire	cct4120	During the past year, how often did you go on an eating binge?	1: Less than once a month 2: 1-3 times a month 3: Once a week 4: More than once a week 5: Never	<b>Any binge-eating</b> <i>Binge-eating must be accompanied by loss of control.</i>  Recode cct4120 (5=0), (1/4=1) Recode cct4125 (3=0) (1/2=1)
	cct4125	Did you feel out of control, like you couldn't stop eating even if you wanted to stop?	1: Yes, usually 2: Yes, sometimes 3: No	0: No (if cct4120 or cct4125=0) 1: Yes (if cct4112 & cct4125=1)  <b>DSM-5<sup>†</sup> Binge-eating</b> Recode cct4120 (5=0) (1/2=0) (3/4=1) Recode cct4125 (3=0) (1/2=1)  0: No (if cct4120 or cct4125=0) 1: Yes (if cct4112 & cct4125=1)
<b>Any disordered eating at any frequency</b>				<b>Presence of any fasting, purging, binge-eating, or excessive exercise:</b> 0: No 1: Yes
<b>Any disordered eating at DSM-5<sup>†</sup> frequency</b>				<b>Presence of any DSM-5<sup>†</sup> fasting, DSM-5<sup>†</sup> purging, DSM-5<sup>†</sup> binge-eating, or DSM-5<sup>†</sup> excessive exercise:</b> 0: No 1: Yes

\*Completed at Teen Focus 4 Clinic. †Behaviour occurs  $\geq 1$  per week. *Abbreviation: ALSPAC, Avon Longitudinal Study of Parents and Children; DSM-5, Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition.*

**Table S3. Derivation of confounder variables used from Avon Longitudinal Study of Parents and Children (ALSPAC)**

Data collection	ALSPAC Variable	Question(s)/measure(s)	Responses	Final variable coding
<b>Child Sex</b>				
Recorded at birth	kz021	Collected from fieldworkers visiting maternity units at birth	1: male 2: female -1: not known	Recode: 0: male 1: female
<b>Parental social class</b>				
1991 British Office of Population and Census Statistics job codes	c755 c765	Derived variable from questions: <ul style="list-style-type: none"> <li>Actual job, occupation, trade or profession</li> <li>Please tick which of the following apply to you: foreman, manager, supervisor, leading hand, self-employed, none of these</li> <li>Type of industry or service given (main things done in job)</li> </ul>	1: I 2: II 3: III (non-manual) 4: III (manual) 5: IV 6: V 65: Armed forces -1: missing	Recode (1/3=0) (4/6=1) (else=.)  Select highest parental social class i.e., if one parent is non-manual then social class=0:  0: Non-manual: professional, managerial, or skilled professions 1: Manual: partly or unskilled occupations
<b>Ethnicity</b>				
	c804		-1: Missing 1: White 2: Non-White	Recode: 0: White 1: Non-White
<b>Maternal education</b>				
	c645a	Derived variable: mother's highest educational qualification	1: CSE/none 2: Vocational 3: O-level 4: A-level 5: Degree -1: Missing	Recode (4/5=0) (3=1) (1/2=2) 0 = A-level or greater 1 = O-level 2 = CSE, vocational or less
<b>Home ownership</b>				
	q2010	<b>Is your home:</b> <ol style="list-style-type: none"> <li>being bought/mortgaged</li> <li>being bought from council</li> <li>owned - with no mortgage to pay</li> <li>rented from council</li> <li>rented from private landlord - furnished</li> <li>rented from private landlord - unfurnished</li> <li>rented from housing association</li> <li>other (please tick &amp; describe)</li> </ol>	0: Being bought/mortgaged 1: Being bought from council 2: Owned with no mortgage to pay 3: Rented from council 4: Rented from private landlord furnished 5: Rented from private landlord unfurnished 6: Rented from housing association 7: other	Recode (0/2=0) (4/5=0) (3=1) (6/7=1) 0 = Mortgage/owned/private rented 1 = rent/other
<b>Material hardship</b>				
ALSPAC hardship items	r9000 r9001 r9002 r9003 r9004	<b>How difficult at the moment do you find it to afford these items:</b> <ol style="list-style-type: none"> <li>food</li> <li>clothing</li> <li>heating</li> <li>rent or mortgage</li> <li>things you need for your children</li> </ol>	1: Very difficult 2: Fairly difficult 3: Slightly difficult 4: Not difficult 5: Paid directly by social security	Recode r9000-r9004 (5=4) 20-(r9000+r9001+r9002+r9003 +9004) = summed score (range: 0-15)  <b>Mode imputation:</b> Replace summed score =. with mode score if at least at least one of r9000-9004 answered

Supplementary material for: *Prospective relationships between continence problems and common mental health disorders in adolescents from a UK cohort*

Data collection	ALSPAC Variable	Question(s)/measure(s)	Responses	Final variable coding (Higher scores = more material hardship)
<b>Family size</b>				
	b032	Derived using parity as proxy.	Continuous variable (range: 0-22)	Recode (0/2=0) (3/22=1) 0 = <3 children 1 = ≥3 children
<b>Developmental delay at 18-month</b>				
Adapted from Denver Developmental Screening Test ( <i>Pediatrics</i> . 1992;89(1):91-7)	kd680	Derived variable: Total ALSPAC development score at 18 months (complete case)	Continuous score -102: Not in correct age range -101: Missing	Continuous score (-101, -102 = missing)
<b>Child IQ</b>				
Adapted from Wechsler Intelligence Scale for Children (3 <sup>rd</sup> ed)	f8ws112	Derived variable: Total IQ score at 8-years	Continuous score -3: Not enough subtests done -2 Did not start WISC	Continuous score (-3, -2 = missing)
<b>Stressful life events</b>				
ALSPAC 'life events inventory' questionnaire		<b>Since the study child's 9<sup>th</sup> birthday:</b>		Recode r5000-r5044 (4=0) (0/3=1) Create summed score (range: 0-45) <b>Mode imputation:</b> Replace summed score =. with mode score if at least at least one of r5000-5044 answered
	r5000	Respondent's husband/partner died	0: Relevant text but no box ticked 1: Yes, since study child was 9 or 10 2: Yes, since child's 11 <sup>th</sup> birthday 3: Yes, both when the study child was 10/11 and since 9 <sup>th</sup> birthday 4: No, did not happen -10: Not completed -1: No response	
	r5001	One of respondent's children has died		
	r5002	Respondent's friend/relative has died		
	r5003	One of respondent's children has been ill		
	r5004	Respondent's husband/partner has been ill		
	r5005	One of respondent's children has been ill		
	r5006	Respondent's friend/relative has been ill		
	r5007	Respondent has been admitted to hospital		
	r5008	Respondent has been in trouble with the law		
	r5009	Respondent has been divorced		
	r5010	Respondent's husband/partner didn't want their child		
	r5011	Respondent has been very ill		
	r5012	Respondent's husband/partner lost his job		
	r5013	Respondent's husband/partner had problems at work		
	r5014	Respondent had problems at work		
	r5015	Respondent lost their job		
	r5016	Respondent's husband/partner went away		
	r5017	Respondent's husband/partner was in trouble		
	r5018	Respondent separated from husband/partner		
	r5019	Respondent's income was reduced		
	r5020	Respondent argued with their husband/partner		
	r5021	Respondent argued with their family/friends		
	r5022	Respondent moved house		
	r5023	Respondent's husband/partner was physically cruel to them		
	r5024	Respondent became homeless		
	r5025	Respondent had a major financial problem		
r5026	Respondent got married			

Supplementary material for: *Prospective relationships between continence problems and common mental health disorders in adolescents from a UK cohort*

Data collection	ALSPAC Variable	Question(s)/measure(s)	Responses	Final variable coding
	r5027	Respondent's husband/partner was physically cruel to their children		
	r5028	Respondent was physically cruel to their children		
	r5029	Respondent attempted suicide		
	r5030	Respondent was convicted of an offence		
	r5031	Respondent became pregnant		
	r5032	Respondent started a new job		
	r5033	Respondent returned to work		
	r5034	Respondent had a miscarriage		
	r5035	Respondent had an abortion		
	r5036	Respondent has taken an examination		
	r5037	Respondent's husband/partner was emotionally cruel to them		
	r5038	Respondent's husband/partner has been emotionally cruel to their children		
	r5039	Respondent has been emotionally cruel to their children		
	r5040	Respondent's house/car was burgled		
	r5041	Respondent found a new partner		
	r5042	One of respondent's child started school		
	r5043	Respondent's husband/partner started a new job		
	r5044	Respondent's pet died		
<b>Maternal depression</b>				
Edinburgh Postnatal Depression Scale when child was 11-years ( <i>Br J Psychiatry J Ment Sci.</i> 1987;150:782–6)		<b>In the past seven days:</b>		
	r4010	I have been able to laugh and see the funny side of things	1: As much as I always could 2: Not quite so much now 3: Definitely not so much now 4: Not at all -10, -1: Missing	Recode r4010, r4011, r4013: (1=0) (2=1) (2=3) (4=3) Recode r4012, r4014-r4019: (4=0), (3=1) (2=2), (1=3) Sum all 10 items to generate continuous score (range:0-30)  <b>Mode imputation:</b> Replace summed score =. with mode score if at least at least one of r4010-r4019 answered
	r4011	I have looked forward with enjoyment to things	1: As much as I ever did 2: Rather less than I used to 3: Definitely less than I used to 4: Hardly at all -10, -1: missing	
	r4012	I have blamed myself unnecessarily when things when wrong	1: Yes, most of the time 2: Yes, some of the time 3: Not very often 4: Never -10, -1: Missing	
	r4013	I have been anxious or worried for no good reason	1: No, not at all 2: Hardly ever 3: Yes, sometimes 4: Yes, often -10, -1: Missing	
r4014	I have felt scared or panicky for no good reason	1: Yes, quite a lot 2: Yes, sometimes		

Supplementary material for: *Prospective relationships between continence problems and common mental health disorders in adolescents from a UK cohort*

Data collection	ALSPAC Variable	Question(s)/measure(s)	Responses	Final variable coding
			3: No, not much 4: No, not at all -10, -1: Missing	
	r4015	Things have been getting on top of me	1: Yes, most of the time I haven't been able to cope 2: Yes, sometimes I haven't been coping 3: No, most of the time I have coped 4: No, I have been coping as well as ever -10, -1: Missing	
	r4016	I have been so unhappy that I have had difficulty sleeping	1: Yes, most of the time 2: Yes, sometimes 3: Not very often 4: No, not at all -10, -1: Missing	
	r4017	I have felt sad or miserable	1: Yes, most of the time 2: Yes, sometimes 3: Not very often 4: No, not at all -10, -1: Missing	
	r4018	I have felt so unhappy that I've been crying	1: Yes, most of the time 2: Yes, quite often 3: Only occasionally 4: Never -10, -1: Missing	
	r4019	The thought of harming myself has occurred to me	1: Yes, quite often 2: Yes, Sometimes 3: Hardly ever 4: Never -10, -1: Missing	
<b>Maternal anxiety</b>				
Crown Crisp Experimental Index when child was 11-years ( <i>Br J Med Psychol.</i> 1988;61(3):255–66)	r4000	Do you feel upset for no obvious reason?	1: Very often 2: Often 3: Not very often 4: Never -10, -1: missing	Recode r4000 r4005 r4006 (3=0) (4=0) (1=2) (2=2) Recode r4001 r4002 r4004 r4007 (4=0) (3=1) (1=2) (2=2) Recode r4003 (4=0) (1=2) (2=2) (3=2) Generate total score: sum of recoded r4000-r4007 (range: 0-24)  <b>Mode imputation:</b> Replace summed score =. with mode score if at least at least one of r4007-r4007 answered
	r4001	Have you felt as though you might faint?		
	r4002	Do you feel uneasy and restless?		
	r4003	Do you sometimes feel panicky?		
	r4004	Do you worry a lot?		
	r4005	Do you feel strung-up inside?		
	r4006	Do you ever have the feeling you are going to pieces?		
	r4007	Do you have bad dreams which upset you when you wake up?		
ku847	Wet self during the day			
ku848	Wet the bed at night			
<b>Child body mass index</b>				
Clinical measures of height and weight at 13.5-years	fg3134	Z-score derived using LMS parameters and 1990 British Growth Reference <sup>1,2</sup> .	Continuous score -110, -106, -101: Missing	Recode z-scores in line with WHO thresholds <sup>3</sup> :



Supplementary material for: *Prospective relationships between continence problems and common mental health disorders in adolescents from a UK cohort*

Data collection	ALSPAC Variable	Question(s)/measure(s)	Responses	Final variable coding
<b>Earlier emotional and behavioural problems</b>				
Adapted from Strengths and Difficulties Questionnaire ( <i>Eur Child Adolesc Psychiatry</i> . 1998;7(3):125–30)	kw660b	Derived variable: SDQ total difficulties score (prorated).	Continuous score (range 0-40) -10, -6, -5: Missing	Continuous score Missing = -10, -6, -5
<i>Abbreviations: ALSPAC, Avon Longitudinal Study of Parents and Children; LMS, lambda-mu-sigma; BMI, body mass index.</i>				

**Table S4. Amount of missing information for each variable in the substantive models**

Variable	N missing (/7332)	%
Daytime wetting	2188	29.84
Bedtime wetting	2187	29.83
Soiling	2195	29.94
Urgency	2179	29.72
Frequent urination	2194	29.92
Low voided volume	2210	30.14
Voiding postponement	2195	29.94
Nocturia	2206	30.09
Common Mental Disorder	3537	48.24
ICD-10 Depression	3537	48.24
High depressive symptoms (SMFQ)	3674	50.11
GAD symptoms	3537	48.24
Physical anxiety score	3711	50.61
Mental anxiety score	3593	49.00
Self-harm act	3537	48.24
Self-harm thoughts	3537	48.24
Any disordered eating	4636	63.23
Excessive exercise	4689	63.95
Fasting	4645	63.35
Purging	4648	63.39
Binge-eating	4650	63.42
DSM-5 frequency disordered eating	4635	63.22
Sex	0	0.00
Parental social class	844	11.51
Ethnicity	664	9.06
Maternal education	556	7.58
Home ownership	1159	15.81
Material hardship	1514	20.65
Family size/Parity	550	7.50
Developmental level	2090	28.51
IQ	0	0.00
Maternal stressful life events	1459	19.90
Maternal depression	1463	19.95
Maternal anxiety	1468	20.02
BMI	2055	28.03
Earlier behaviour and emotional problems (SDQ)	1637	22.33

**Table S5. Descriptive information for secondary outcomes (imputed and complete case samples)**

Variable	Imputed sample (n=7,332)		Mental health sample (n=1528)		Disordered eating sample (n=1375)	
	% or mean (se)		% or mean (n or SD)		% or mean (n or SD)	
Physical anxiety	25.44	0.11	25.40		7.51	
Mental anxiety	21.84	0.08	21.68		4.81	
Self-harm thoughts	11.74%	0.56	9.42%		144	
Excessive exercise	21.89%	0.81			22.55%	310
Fasting	11.49%	0.69			10.76%	148
Purging	6.08%	0.44			5.53%	76
Binge-eating	13.42%	0.72			11.71%	161
DSM-5 disordered eating	11.85%	0.64			9.67%	133

DSM-5 - Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition

**Table S6. Sequential adjustments for primary outcomes using imputed data (n=7,332)**

Exposure	Outcome	Unadjusted		Adjusted 1		Adjusted 2		Adjusted 3		Adjusted 4		Adjusted 5	
		OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p
Daytime wetting	Common mental disorder	2.38 (1.59, 3.57)	<.001	1.99 (1.32, 3.01)	0.001	1.88 (1.23, 2.86)	0.003	1.89 (1.24, 2.89)	0.003	1.79 (1.17, 2.74)	0.007	1.59 (1.03, 2.47)	0.036
Bedwetting	Common mental disorder	1.77 (1.06, 2.96)	0.029	1.90 (1.12, 3.22)	0.017	1.67 (0.97, 2.89)	0.065	1.68 (0.97, 2.89)	0.063	1.66 (0.96, 2.88)	0.071	1.39 (0.78, 2.47)	0.259
Soiling	Common mental disorder	1.56 (1.08, 2.26)	0.019	1.34 (0.92, 1.97)	0.128	1.32 (0.90, 1.95)	0.157	1.33 (0.90, 1.96)	0.151	1.30 (0.88, 1.92)	0.192	1.25 (0.84, 1.86)	0.278
Urgency	Common mental disorder	2.07 (1.40, 3.06)	<.001	1.99 (1.33, 2.98)	0.001	1.74 (1.15, 2.64)	0.009	1.75 (1.16, 2.66)	0.008	1.67 (1.10, 2.54)	0.017	1.54 (1.00, 2.36)	0.05
Frequent urination	Common mental disorder	1.83 (1.07, 3.10)	0.027	1.74 (1.01, 3.00)	0.047	1.55 (0.89, 2.71)	0.12	1.56 (0.89, 2.72)	0.116	1.53 (0.87, 2.67)	0.139	1.41 (0.80, 2.50)	0.231
Low voided volume	Common mental disorder	1.75 (1.16, 2.65)	0.008	1.64 (1.08, 2.49)	0.021	1.56 (1.02, 2.39)	0.039	1.57 (1.02, 2.40)	0.039	1.55 (1.01, 2.38)	0.047	1.45 (0.94, 2.26)	0.095
Voiding postponement	Common mental disorder	2.06 (1.63, 2.60)	<.001	2.03 (1.60, 2.58)	<.001	1.95 (1.52, 2.49)	<.001	1.95 (1.52, 2.49)	<.001	1.93 (1.51, 2.46)	<.001	1.88 (1.46, 2.41)	<.001
Nocturia	Common mental disorder	1.44 (1.07, 1.94)	0.017	1.41 (1.04, 1.91)	0.028	1.30 (0.95, 1.77)	0.103	1.30 (0.95, 1.78)	0.101	1.28 (0.93, 1.76)	0.124	1.19 (0.86, 1.64)	0.303
Exposure	Outcome	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p
Daytime wetting	ICD-10 depression	2.48 (1.52, 4.04)	<.001	2.08 (1.27, 3.40)	0.004	1.95 (1.18, 3.24)	0.01	2.00 (1.20, 3.32)	0.008	1.96 (1.18, 3.24)	0.009	1.77 (1.05, 2.98)	0.032
Bedwetting	ICD-10 depression	0.61 (0.22, 1.70)	0.34	0.64 (0.23, 1.79)	0.389	0.52 (0.18, 1.50)	0.224	0.52 (0.18, 1.52)	0.232	0.52 (0.18, 1.52)	0.232	0.44 (0.15, 1.31)	0.14
Soiling	ICD-10 depression	1.18 (0.67, 2.05)	0.565	1.01 (0.57, 1.78)	0.968	1.00 (0.56, 1.79)	0.991	1.03 (0.58, 1.85)	0.915	1.03 (0.57, 1.84)	0.933	0.99 (0.55, 1.78)	0.964
Urgency	ICD-10 depression	2.43 (1.56, 3.78)	<.001	2.33 (1.49, 3.65)	<.001	2.00 (1.25, 3.19)	0.004	2.08 (1.30, 3.34)	0.002	2.05 (1.28, 3.30)	0.003	1.94 (1.19, 3.14)	0.008
Frequent urination	ICD-10 depression	1.33 (0.67, 2.67)	0.413	1.26 (0.63, 2.54)	0.516	1.08 (0.53, 2.22)	0.822	1.12 (0.55, 2.30)	0.752	1.11 (0.54, 2.28)	0.774	1.04 (0.50, 2.14)	0.919
Low voided volume	ICD-10 depression	1.82 (1.10, 2.99)	0.019	1.69 (1.02, 2.81)	0.041	1.60 (0.95, 2.67)	0.075	1.62 (0.97, 2.71)	0.064	1.62 (0.97, 2.70)	0.067	1.53 (0.91, 2.57)	0.107
Voiding postponement	ICD-10 depression	1.74 (1.29, 2.35)	<.001	1.70 (1.25, 2.31)	0.001	1.61 (1.18, 2.20)	0.003	1.62 (1.18, 2.22)	0.003	1.61 (1.18, 2.21)	0.003	1.58 (1.15, 2.16)	0.005
Nocturia	ICD-10 depression	1.32 (0.89, 1.96)	0.163	1.29 (0.86, 1.92)	0.212	1.16 (0.77, 1.75)	0.472	1.19 (0.79, 1.80)	0.398	1.19 (0.79, 1.79)	0.414	1.11 (0.73, 1.69)	0.628
Exposure	Outcome	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p
Daytime wetting	High depressive symptoms	2.26 (1.54, 3.32)	<.001	2.03 (1.38, 2.98)	<.001	1.95 (1.32, 2.87)	0.001	1.91 (1.29, 2.84)	0.001	1.84 (1.24, 2.74)	0.002	1.70 (1.13, 2.56)	0.011
Bedwetting	High depressive symptoms	1.42 (0.88, 2.32)	0.154	1.48 (0.90, 2.41)	0.12	1.29 (0.77, 2.15)	0.324	1.26 (0.75, 2.11)	0.381	1.25 (0.74, 2.11)	0.403	1.10 (0.63, 1.89)	0.741
Soiling	High depressive symptoms	1.83 (1.30, 2.59)	0.001	1.68 (1.19, 2.38)	0.004	1.70 (1.19, 2.43)	0.004	1.59 (1.11, 2.29)	0.012	1.56 (1.08, 2.25)	0.018	1.52 (1.05, 2.20)	0.029
Urgency	High depressive symptoms	1.27 (0.86, 1.90)	0.231	1.23 (0.83, 1.85)	0.302	1.05 (0.69, 1.61)	0.821	0.97 (0.63, 1.50)	0.905	0.94 (0.61, 1.45)	0.771	0.87 (0.56, 1.35)	0.535
Frequent urination	High depressive symptoms	1.35 (0.85, 2.13)	0.202	1.30 (0.82, 2.07)	0.262	1.16 (0.72, 1.87)	0.543	1.09 (0.67, 1.76)	0.727	1.06 (0.66, 1.73)	0.799	1.01 (0.62, 1.64)	0.981
Low voided volume	High depressive symptoms	2.21 (1.55, 3.14)	<.001	2.13 (1.49, 3.03)	<.001	2.02 (1.41, 2.91)	<.001	1.98 (1.37, 2.86)	<.001	1.95 (1.34, 2.82)	<.001	1.87 (1.28, 2.73)	0.001
Voiding postponement	High depressive symptoms	1.68 (1.35, 2.10)	<.001	1.66 (1.33, 2.07)	<.001	1.59 (1.26, 1.99)	<.001	1.58 (1.25, 1.99)	<.001	1.56 (1.24, 1.97)	<.001	1.53 (1.21, 1.93)	<.001
Nocturia	High depressive symptoms	1.55 (1.19, 2.03)	0.001	1.53 (1.17, 2.01)	0.002	1.41 (1.07, 1.86)	0.015	1.34 (1.02, 1.78)	0.038	1.33 (1.00, 1.76)	0.047	1.26 (0.95, 1.68)	0.109
Exposure	Outcome	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p
Daytime wetting	GAD symptoms	4.11 (2.51, 6.74)	<.001	3.57 (2.17, 5.87)	<.001	3.30 (1.98, 5.49)	<.001	3.30 (1.97, 5.51)	<.001	3.18 (1.89, 5.33)	<.001	3.01 (1.78, 5.09)	<.001
Bedwetting	GAD symptoms	2.50 (1.22, 5.09)	0.012	2.64 (1.29, 5.40)	0.008	2.39 (1.16, 4.92)	0.019	2.36 (1.15, 4.85)	0.02	2.32 (1.12, 4.78)	0.023	2.16 (1.03, 4.52)	0.042
Soiling	GAD symptoms	1.69 (0.96, 2.96)	0.067	1.49 (0.84, 2.62)	0.168	1.44 (0.82, 2.53)	0.203	1.39 (0.79, 2.44)	0.253	1.35 (0.76, 2.39)	0.304	1.32 (0.74, 2.35)	0.348
Urgency	GAD symptoms	2.64 (1.54, 4.52)	<.001	2.54 (1.47, 4.39)	0.001	2.31 (1.33, 4.03)	0.003	2.24 (1.29, 3.90)	0.005	2.16 (1.23, 3.79)	0.008	2.09 (1.19, 3.67)	0.011
Frequent urination	GAD symptoms	2.80 (1.48, 5.30)	0.002	2.69 (1.42, 5.10)	0.003	2.56 (1.34, 4.88)	0.005	2.48 (1.30, 4.74)	0.006	2.44 (1.27, 4.69)	0.008	2.37 (1.23, 4.57)	0.01

Supplementary material for: *Prospective relationships between continence problems and common mental health disorders in adolescents from a UK cohort*

Low voided volume	GAD symptoms	1.22 (0.62, 2.42)	0.564	1.14 (0.57, 2.28)	0.701	1.11 (0.56, 2.20)	0.767	1.09 (0.55, 2.18)	0.805	1.07 (0.53, 2.16)	0.841	1.03 (0.51, 2.07)	0.935
Voiding postponement	GAD symptoms	1.74 (1.22, 2.50)	0.003	1.70 (1.19, 2.45)	0.004	1.64 (1.14, 2.36)	0.008	1.63 (1.13, 2.36)	0.009	1.61 (1.12, 2.33)	0.01	1.59 (1.10, 2.30)	0.013
Nocturia	GAD symptoms	2.02 (1.31, 3.09)	0.001	1.98 (1.28, 3.04)	0.002	1.87 (1.21, 2.90)	0.005	1.83 (1.18, 2.82)	0.007	1.80 (1.17, 2.78)	0.008	1.73 (1.12, 2.68)	0.014
<b>Exposure</b>	<b>Outcome</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>
Daytime wetting	Self-harm act	1.66 (0.94, 2.95)	0.082	1.37 (0.77, 2.46)	0.282	1.26 (0.69, 2.28)	0.449	1.23 (0.68, 2.24)	0.498	1.16 (0.63, 2.12)	0.637	1.07 (0.58, 1.98)	0.826
Bedwetting	Self-harm act	1.98 (1.11, 3.54)	0.021	2.12 (1.17, 3.83)	0.013	1.94 (1.06, 3.53)	0.031	1.92 (1.05, 3.50)	0.034	1.87 (1.02, 3.44)	0.044	1.71 (0.93, 3.16)	0.086
Soiling	Self-harm act	1.36 (0.83, 2.22)	0.216	1.17 (0.71, 1.92)	0.541	1.14 (0.69, 1.88)	0.619	1.11 (0.67, 1.83)	0.691	1.07 (0.64, 1.77)	0.802	1.04 (0.62, 1.73)	0.885
Urgency	Self-harm act	0.70 (0.34, 1.45)	0.337	0.66 (0.31, 1.38)	0.265	0.57 (0.28, 1.20)	0.138	0.56 (0.27, 1.17)	0.121	0.52 (0.25, 1.10)	0.089	0.49 (0.23, 1.05)	0.065
Frequent urination	Self-harm act	1.63 (0.89, 2.98)	0.112	1.54 (0.83, 2.85)	0.167	1.43 (0.77, 2.67)	0.255	1.40 (0.75, 2.62)	0.284	1.37 (0.73, 2.58)	0.33	1.32 (0.70, 2.50)	0.396
Low voided volume	Self-harm act	0.88 (0.50, 1.55)	0.645	0.80 (0.45, 1.43)	0.456	0.77 (0.43, 1.37)	0.377	0.76 (0.42, 1.35)	0.346	0.74 (0.42, 1.33)	0.313	0.71 (0.39, 1.27)	0.245
Voiding postponement	Self-harm act	1.17 (0.85, 1.63)	0.335	1.14 (0.82, 1.59)	0.437	1.09 (0.78, 1.53)	0.604	1.10 (0.78, 1.54)	0.593	1.08 (0.77, 1.52)	0.665	1.06 (0.75, 1.49)	0.752
Nocturia	Self-harm act	1.35 (0.92, 1.97)	0.126	1.31 (0.89, 1.93)	0.173	1.24 (0.84, 1.83)	0.288	1.22 (0.82, 1.80)	0.319	1.20 (0.81, 1.77)	0.373	1.14 (0.77, 1.70)	0.513
<b>Exposure</b>	<b>Outcome</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>
Daytime wetting	Any disordered eating	2.06 (1.41, 3.01)	<.001	1.58 (1.06, 2.35)	0.024	1.56 (1.05, 2.31)	0.029	1.61 (1.08, 2.41)	0.019	1.58 (1.06, 2.37)	0.026	1.48 (0.98, 2.25)	0.064
Bedwetting	Any disordered eating	1.51 (0.98, 2.30)	0.059	1.71 (1.09, 2.69)	0.021	1.72 (1.09, 2.72)	0.02	1.74 (1.10, 2.76)	0.017	1.72 (1.09, 2.72)	0.02	1.61 (1.00, 2.59)	0.052
Soiling	Any disordered eating	1.68 (1.21, 2.33)	0.002	1.35 (0.96, 1.92)	0.088	1.34 (0.94, 1.91)	0.101	1.38 (0.97, 1.98)	0.073	1.36 (0.95, 1.94)	0.091	1.33 (0.92, 1.91)	0.131
Urgency	Any disordered eating	1.33 (0.92, 1.94)	0.132	1.25 (0.83, 1.87)	0.278	1.24 (0.82, 1.88)	0.303	1.27 (0.84, 1.94)	0.259	1.25 (0.82, 1.90)	0.296	1.21 (0.79, 1.86)	0.378
Frequent urination	Any disordered eating	1.36 (0.90, 2.07)	0.144	1.27 (0.81, 2.00)	0.304	1.27 (0.80, 2.03)	0.305	1.30 (0.82, 2.08)	0.264	1.29 (0.81, 2.07)	0.282	1.24 (0.77, 2.00)	0.368
Low voided volume	Any disordered eating	1.66 (1.11, 2.48)	0.014	1.52 (0.98, 2.36)	0.061	1.53 (0.98, 2.38)	0.061	1.56 (1.00, 2.44)	0.05	1.56 (1.00, 2.45)	0.05	1.52 (0.97, 2.40)	0.07
Voiding postponement	Any disordered eating	1.41 (1.13, 1.75)	0.003	1.38 (1.09, 1.74)	0.007	1.38 (1.09, 1.74)	0.007	1.37 (1.09, 1.73)	0.008	1.36 (1.08, 1.72)	0.01	1.36 (1.07, 1.74)	0.013
Nocturia	Any disordered eating	1.76 (1.34, 2.31)	<.001	1.76 (1.32, 2.36)	<.001	1.78 (1.32, 2.38)	<.001	1.81 (1.34, 2.43)	<.001	1.80 (1.33, 2.42)	<.001	1.72 (1.27, 2.34)	0.001

GAD – Generalised Anxiety Disorder; ICD-10 - International Classification of Diseases, Tenth Revision.

Adjusted 1 confounders: sex. Adjusted 2 confounders: sex and socioeconomic indicators (parental occupational social class, maternal educational attainment, family size, ethnicity, home ownership status, material hardship). Adjusted 3 confounders: sex, socioeconomic indicators, child IQ and developmental level. Adjusted 4 confounders: sex, socioeconomic indicators, child IQ, developmental level, maternal stressful life events, maternal depression and maternal anxiety. Adjusted 5 confounders: sex, socioeconomic indicators, child IQ, developmental level, maternal stressful life events, maternal depression, maternal anxiety, child Body Mass Index and earlier emotional/behaviour problems (fully adjusted model).

**Table S7. Sequential adjustments for primary outcomes using complete case data (mental health outcomes n=1,528; disordered eating outcome n=1,375)**

Exposure	Outcome	Unadjusted		Adjusted 1		Adjusted 2		Adjusted 3		Adjusted 4		Adjusted 5	
		OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p
Daytime wetting	Common mental disorder	2.44 (1.30, 4.56)	0.005	2.08 (1.10, 3.92)	0.024	2.09 (1.10, 3.95)	0.024	2.07 (1.09, 3.93)	0.025	2.02 (1.06, 3.84)	0.032	1.95 (1.02, 3.74)	0.044
Bedwetting	Common mental disorder	1.01 (0.39, 2.60)	0.991	1.13 (0.43, 2.97)	0.800	1.07 (0.41, 2.84)	0.884	1.07 (0.40, 2.83)	0.891	1.09 (0.41, 2.90)	0.865	0.89 (0.33, 2.41)	0.820
Soiling	Common mental disorder	1.63 (0.87, 3.04)	0.125	1.38 (0.74, 2.60)	0.314	1.34 (0.71, 2.53)	0.371	1.32 (0.70, 2.50)	0.393	1.35 (0.71, 2.56)	0.361	1.31 (0.69, 2.50)	0.415
Urgency	Common mental disorder	2.23 (1.22, 4.07)	0.009	2.15 (1.17, 3.97)	0.014	2.09 (1.13, 3.87)	0.019	2.08 (1.13, 3.86)	0.019	2.04 (1.10, 3.79)	0.024	2.00 (1.07, 3.73)	0.030
Frequent urination	Common mental disorder	1.62 (0.70, 3.73)	0.260	1.54 (0.66, 3.61)	0.315	1.52 (0.64, 3.57)	0.340	1.50 (0.64, 3.54)	0.351	1.51 (0.64, 3.57)	0.347	1.44 (0.60, 3.43)	0.412
Low voided volume	Common mental disorder	1.21 (0.61, 2.41)	0.588	1.10 (0.55, 2.20)	0.798	1.11 (0.55, 2.23)	0.776	1.10 (0.55, 2.22)	0.785	1.07 (0.53, 2.17)	0.847	0.98 (0.48, 1.99)	0.945
Voiding postponement	Common mental disorder	1.60 (1.05, 2.42)	0.028	1.59 (1.04, 2.43)	0.031	1.56 (1.02, 2.39)	0.039	1.57 (1.03, 2.40)	0.037	1.48 (0.96, 2.27)	0.073	1.45 (0.94, 2.23)	0.092
Nocturia	Common mental disorder	1.29 (0.79, 2.11)	0.309	1.30 (0.79, 2.14)	0.299	1.30 (0.79, 2.14)	0.308	1.29 (0.78, 2.13)	0.322	1.28 (0.77, 2.11)	0.344	1.26 (0.75, 2.09)	0.381
Exposure	Outcome	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p
Daytime wetting	ICD-10 depression	3.22 (1.58, 6.60)	0.001	2.85 (1.38, 5.87)	0.004	2.98 (1.44, 6.16)	0.003	3.03 (1.46, 6.28)	0.003	3.04 (1.46, 6.33)	0.003	2.94 (1.40, 6.16)†	0.004
Bedwetting	ICD-10 depression	0.35 (0.05, 2.58)	0.303	0.38 (0.05, 2.81)	0.344	0.36 (0.05, 2.64)	0.313	0.36 (0.05, 2.66)	0.315	0.36 (0.05, 2.71)	0.323	0.30 (0.04, 2.27) †	0.245
Soiling	ICD-10 depression	1.57 (0.70, 3.52)	0.275	1.38 (0.61, 3.11)	0.438	1.36 (0.60, 3.08)	0.468	1.39 (0.61, 3.16)	0.437	1.38 (0.61, 3.15)	0.442	1.35 (0.59, 3.10)†	0.477
Urgency	ICD-10 depression	3.09 (1.56, 6.11)	0.001	2.99 (1.50, 5.95)	0.002	2.89 (1.44, 5.80)	0.003	2.93 (1.46, 5.87)	0.002	2.95 (1.47, 5.93)	0.002	2.90 (1.43, 5.85)†	0.003
Frequent urination	ICD-10 depression	1.20 (0.36, 3.98)	0.764	1.15 (0.35, 3.83)	0.818	1.15 (0.34, 3.85)	0.824	1.17 (0.35, 3.92)	0.802	1.18 (0.35, 3.95)	0.791	1.13 (0.33, 3.81)†	0.850
Low voided volume	ICD-10 depression	1.91 (0.89, 4.10)	0.099	1.77 (0.82, 3.83)	0.145	1.79 (0.82, 3.89)	0.141	1.81 (0.83, 3.93)	0.136	1.79 (0.82, 3.90)	0.142	1.63 (0.74, 3.58)†	0.221
Voiding postponement	ICD-10 depression	1.40 (0.80, 2.44)	0.237	1.39 (0.79, 2.43)	0.251	1.36 (0.78, 2.39)	0.282	1.36 (0.77, 2.39)	0.287	1.35 (0.77, 2.38)	0.298	1.30 (0.74, 2.30)†	0.362
Nocturia	ICD-10 depression	1.51 (0.82, 2.77)	0.187	1.52 (0.82, 2.80)	0.184	1.52 (0.82, 2.82)	0.180	1.54 (0.83, 2.85)	0.169	1.55 (0.83, 2.86)	0.167	1.52 (0.82, 2.83)†	0.185
Exposure	Outcome	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p
Daytime wetting	High depressive symptoms	1.88 (1.03, 3.41)	0.038	1.70 (0.93, 3.11)	0.083	1.71 (0.93, 3.14)	0.084	1.65 (0.89, 3.03)	0.110	1.61 (0.87, 2.97)	0.128	1.58 (0.86, 2.93)	0.143
Bedwetting	High depressive symptoms	0.97 (0.43, 2.23)	0.951	1.04 (0.45, 2.39)	0.926	0.96 (0.41, 2.24)	0.926	0.93 (0.40, 2.19)	0.871	0.92 (0.39, 2.17)	0.853	0.83 (0.35, 1.99)	0.682
Soiling	High depressive symptoms	2.18 (1.29, 3.68)	0.004	1.99 (1.17, 3.37)	0.011	1.97 (1.15, 3.37)	0.014	1.81 (1.05, 3.12)	0.031	1.84 (1.07, 3.17)	0.027	1.83 (1.06, 3.16)	0.030
Urgency	High depressive symptoms	1.05 (0.55, 2.00)	0.880	1.02 (0.53, 1.94)	0.957	0.95 (0.49, 1.82)	0.868	0.92 (0.48, 1.79)	0.815	0.88 (0.45, 1.71)	0.707	0.88 (0.45, 1.70)	0.695
Frequent urination	High depressive symptoms	1.24 (0.56, 2.73)	0.600	1.20 (0.54, 2.66)	0.656	1.17 (0.52, 2.63)	0.712	1.11 (0.49, 2.53)	0.801	1.09 (0.48, 2.49)	0.842	1.07 (0.46, 2.45)	0.880
Low voided volume	High depressive symptoms	1.82 (1.05, 3.15)	0.032	1.73 (0.99, 3.00)	0.052	1.76 (1.01, 3.08)	0.046	1.72 (0.98, 3.02)	0.059	1.68 (0.96, 2.95)	0.072	1.60 (0.91, 2.83)	0.103
Voiding postponement	High depressive symptoms	1.55 (1.07, 2.24)	0.020	1.54 (1.07, 2.24)	0.022	1.51 (1.04, 2.21)	0.031	1.55 (1.06, 2.26)	0.023	1.49 (1.02, 2.19)	0.040	1.47 (1.00, 2.16)	0.047
Nocturia	High depressive symptoms	1.37 (0.90, 2.09)	0.146	1.38 (0.90, 2.11)	0.142	1.37 (0.89, 2.11)	0.152	1.32 (0.85, 2.03)	0.217	1.32 (0.85, 2.05)	0.213	1.31 (0.85, 2.04)	0.222
Exposure	Outcome	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p
Daytime wetting	GAD symptoms	3.41 (1.55, 7.50)	0.002	2.96 (1.34, 6.54)	0.007	3.02 (1.36, 6.71)	0.007	2.98 (1.34, 6.64)	0.008	2.87 (1.28, 6.42)	0.010	2.80 (1.24, 6.32)†	0.013
Bedwetting	GAD symptoms	2.19 (0.76, 6.31)	0.149	2.46 (0.84, 7.18)	0.100	2.34 (0.79, 6.88)	0.123	2.31 (0.79, 6.82)	0.128	2.29 (0.78, 6.78)	0.134	2.04 (0.68, 6.11)†	0.203
Soiling	GAD symptoms	1.48 (0.58, 3.80)	0.411	1.28 (0.50, 3.29)	0.612	1.25 (0.48, 3.24)	0.648	1.22 (0.47, 3.16)	0.689	1.25 (0.48, 3.24)	0.648	1.25 (0.48, 3.26)†	0.649

Supplementary material for: *Prospective relationships between continence problems and common mental health disorders in adolescents from a UK cohort*

		Unadjusted		Adjusted 1		Adjusted 2		Adjusted 3		Adjusted 4		Adjusted 5	
Urgency	GAD symptoms	2.45 (1.08, 5.58)	0.032	2.36 (1.03, 5.39)	0.042	2.32 (1.00, 5.34)	0.049	2.30 (1.00, 5.30)	0.051	2.21 (0.95, 5.11)	0.065	2.20 (0.95, 5.13)†	0.066
Frequent urination	GAD symptoms	3.81 (1.54, 9.44)	0.004	3.69 (1.48, 9.19)	0.005	3.66 (1.45, 9.25)	0.006	3.61 (1.43, 9.15)	0.007	3.60 (1.41, 9.14)	0.007	3.44 (1.34, 8.81)†	0.010
Low voided volume	GAD symptoms	1.53 (0.60, 3.93)	0.373	1.40 (0.55, 3.62)	0.481	1.42 (0.55, 3.66)	0.473	1.40 (0.54, 3.62)	0.490	1.37 (0.53, 3.56)	0.518	1.27 (0.49, 3.31)†	0.629
Voiding postponement	GAD symptoms	1.12 (0.57, 2.22)	0.741	1.11 (0.56, 2.20)	0.769	1.08 (0.54, 2.16)	0.823	1.09 (0.55, 2.18)	0.802	1.03 (0.51, 2.06)	0.934	1.00 (0.50, 2.00)†	0.991
Nocturia	GAD symptoms	1.56 (0.78, 3.10)	0.206	1.57 (0.79, 3.13)	0.202	1.57 (0.78, 3.14)	0.205	1.55 (0.77, 3.10)	0.221	1.52 (0.76, 3.07)	0.238	1.51 (0.75, 3.06)†	0.246
Exposure	Outcome	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p
Daytime wetting	Self-harm act	1.42 (0.60, 3.38)	0.430	1.23 (0.51, 2.94)	0.645	1.29 (0.54, 3.10)	0.572	1.28 (0.53, 3.09)	0.583	1.21 (0.50, 2.92)	0.678	1.22 (0.50, 2.98)†	0.657
Bedwetting	Self-harm act	2.59 (1.12, 5.99)	0.026	2.91 (1.24, 6.81)	0.014	2.99 (1.27, 7.08)	0.013	2.99 (1.26, 7.07)	0.013	2.98 (1.25, 7.12)	0.014	2.77 (1.15, 6.67)†	0.023
Soiling	Self-harm act	1.53 (0.72, 3.28)	0.272	1.34 (0.62, 2.88)	0.459	1.27 (0.59, 2.77)	0.541	1.26 (0.58, 2.75)	0.565	1.31 (0.60, 2.86)	0.494	1.32 (0.61, 2.90)†	0.482
Urgency	Self-harm act	0.56 (0.17, 1.81)	0.334	0.53 (0.16, 1.73)	0.295	0.51 (0.16, 1.67)	0.266	0.51 (0.16, 1.67)	0.266	0.48 (0.15, 1.57)	0.224	0.49 (0.15, 1.61)†	0.238
Frequent urination	Self-harm act	1.01 (0.31, 3.33)	0.989	0.96 (0.29, 3.20)	0.950	0.92 (0.27, 3.09)	0.890	0.92 (0.27, 3.09)	0.888	0.89 (0.26, 3.04)	0.855	0.88 (0.26, 3.01)†	0.833
Low voided volume	Self-harm act	0.92 (0.36, 2.33)	0.857	0.84 (0.33, 2.15)	0.720	0.84 (0.33, 2.15)	0.719	0.83 (0.32, 2.13)	0.699	0.81 (0.32, 2.09)	0.666	0.76 (0.30, 1.97)†	0.576
Voiding postponement	Self-harm act	1.15 (0.66, 1.99)	0.624	1.13 (0.65, 1.97)	0.654	1.09 (0.62, 1.90)	0.769	1.10 (0.63, 1.92)	0.747	1.01 (0.58, 1.78)	0.963	0.99 (0.56, 1.74)†	0.970
Nocturia	Self-harm act	1.24 (0.68, 2.27)	0.481	1.25 (0.68, 2.29)	0.474	1.23 (0.67, 2.27)	0.504	1.22 (0.66, 2.26)	0.517	1.19 (0.64, 2.20)	0.588	1.20 (0.65, 2.23)†	0.560
Exposure	Outcome	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p
Daytime wetting	Any disordered eating	2.07 (1.15, 3.73)	0.016	1.67 (0.91, 3.08)	0.100	1.72 (0.93, 3.20)	0.085	1.78 (0.96, 3.32)	0.068	1.75 (0.94, 3.27)	0.077	1.58 (0.84, 2.99)	0.160
Bedwetting	Any disordered eating	1.45 (0.74, 2.85)	0.275	1.66 (0.81, 3.39)	0.163	1.64 (0.80, 3.37)	0.177	1.68 (0.82, 3.44)	0.158	1.67 (0.81, 3.44)	0.163	1.41 (0.67, 2.98)	0.365
Soiling	Any disordered eating	1.35 (0.82, 2.22)	0.241	1.17 (0.69, 1.97)	0.559	1.15 (0.68, 1.94)	0.611	1.19 (0.70, 2.02)	0.526	1.18 (0.70, 2.01)	0.534	1.12 (0.66, 1.92)	0.674
Urgency	Any disordered eating	1.24 (0.75, 2.07)	0.405	1.18 (0.69, 2.02)	0.540	1.21 (0.71, 2.07)	0.484	1.24 (0.72, 2.13)	0.434	1.22 (0.71, 2.10)	0.472	1.22 (0.70, 2.11)	0.483
Frequent urination	Any disordered eating	1.16 (0.62, 2.16)	0.647	1.13 (0.59, 2.18)	0.712	1.14 (0.59, 2.21)	0.693	1.17 (0.60, 2.27)	0.639	1.16 (0.60, 2.25)	0.660	1.08 (0.55, 2.12)	0.830
Low voided volume	Any disordered eating	1.78 (1.08, 2.94)	0.024	1.69 (1.00, 2.86)	0.051	1.67 (0.98, 2.83)	0.059	1.74 (1.02, 2.97)	0.043	1.74 (1.02, 2.97)	0.043	1.65 (0.96, 2.85)	0.071
Voiding postponement	Any disordered eating	1.35 (0.98, 1.86)	0.065	1.33 (0.95, 1.87)	0.092	1.34 (0.95, 1.88)	0.091	1.32 (0.94, 1.86)	0.106	1.29 (0.92, 1.82)	0.145	1.28 (0.90, 1.81)	0.166
Nocturia	Any disordered eating	1.51 (1.02, 2.22)	0.037	1.59 (1.06, 2.39)	0.025	1.59 (1.05, 2.39)	0.027	1.62 (1.07, 2.44)	0.022	1.61 (1.07, 2.43)	0.023	1.58 (1.04, 2.41)	0.032

GAD – Generalised Anxiety Disorder; ICD-10 - International Classification of Diseases, Tenth Revision. †Analyses performed on n=1494 due to 34 observations being dropped (underweight BMI predicted outcome perfectly).

Adjusted 1 confounders: sex. Adjusted 2 confounders: sex and socioeconomic indicators (parental occupational social class, maternal educational attainment, family size, ethnicity, home ownership status, material hardship). Adjusted 3 confounders: sex, socioeconomic indicators, child IQ and developmental level. Adjusted 4 confounders: sex, socioeconomic indicators, child IQ, developmental level, maternal stressful life events, maternal depression and maternal anxiety. Adjusted 5 confounders: sex, socioeconomic indicators, child IQ, developmental level, maternal stressful life events, maternal depression, maternal anxiety, child Body Mass Index and earlier emotional/behaviour problems (fully adjusted model).

**Table S8. Sequential adjustments for secondary outcomes using imputed data (n=7,332)**

Exposure	Outcome	Unadjusted		Adjusted 1		Adjusted 2		Adjusted 3		Adjusted 4		Adjusted 5	
		B (95% CI)	p	B (95% CI)	p	B (95% CI)	p	B (95% CI)	p	B (95% CI)	p	B (95% CI)	p
Daytime wetting	Physical anxiety	2.15 (0.77, 3.52)	0.00	1.19 (-0.16, 2.54)	0.08	1.13 (-0.20, 2.45)	0.09	1.10 (-0.23, 2.43)	0.10	0.99 (-0.34, 2.32)	0.14	0.92 (-0.42, 2.25)	0.17
		0.19 (-1.51, 1.89)	0.82	0.42 (-1.23, 2.08)	0.61	0.32 (-1.31, 1.95)	0.70	0.30 (-1.34, 1.93)	0.72	0.29 (-1.35, 1.93)	0.73	0.16 (-1.48, 1.81)	0.84
Bedwetting	Physical anxiety	2.26 (1.04, 3.47)	<.00	1.51 (0.32, 2.69)	0.01	1.47 (0.29, 2.65)	0.01	1.42 (0.23, 2.60)	0.01	1.36 (0.17, 2.54)	0.02	1.34 (0.15, 2.52)	0.02
		1.93 (0.69, 3.16)	0.00	1.67 (0.44, 2.89)	0.00	1.59 (0.35, 2.82)	0.01	1.53 (0.29, 2.77)	0.01	1.44 (0.19, 2.68)	0.02	1.36 (0.12, 2.60)	0.03
Soiling	Physical anxiety	1.10 (-0.51, 2.70)	0.17	0.80 (-0.77, 2.38)	0.31	0.76 (-0.78, 2.31)	0.33	0.71 (-0.84, 2.26)	0.36	0.64 (-0.91, 2.20)	0.41	0.60 (-0.94, 2.15)	0.44
		1.94 (0.56, 3.31)	0.00	1.56 (0.20, 2.91)	0.02	1.58 (0.23, 2.92)	0.02	1.55 (0.21, 2.90)	0.02	1.48 (0.13, 2.83)	0.03	1.43 (0.08, 2.79)	0.03
Urgency	Physical anxiety	1.89 (1.17, 2.61)	<.00	1.75 (1.05, 2.45)	<.00	1.72 (1.02, 2.41)	<.00	1.71 (1.02, 2.41)	<.00	1.67 (0.98, 2.36)	<.00	1.64 (0.95, 2.33)	<.00
		1.10 (0.20, 2.00)	0.01	0.95 (0.07, 1.84)	0.03	0.91 (0.03, 1.79)	0.04	0.87 (-0.01, 1.75)	0.05	0.84 (-0.05, 1.72)	0.06	0.81 (-0.07, 1.69)	0.07
Frequent urination	Physical anxiety	2.15 (0.77, 3.52)	0.00	1.19 (-0.16, 2.54)	0.08	1.13 (-0.20, 2.45)	0.09	1.10 (-0.23, 2.43)	0.10	0.99 (-0.34, 2.32)	0.14	0.92 (-0.42, 2.25)	0.17
		0.19 (-1.51, 1.89)	0.82	0.42 (-1.23, 2.08)	0.61	0.32 (-1.31, 1.95)	0.70	0.30 (-1.34, 1.93)	0.72	0.29 (-1.35, 1.93)	0.73	0.16 (-1.48, 1.81)	0.84
Low voided volume voiding postponement	Physical anxiety	2.26 (1.04, 3.47)	<.00	1.51 (0.32, 2.69)	0.01	1.47 (0.29, 2.65)	0.01	1.42 (0.23, 2.60)	0.01	1.36 (0.17, 2.54)	0.02	1.34 (0.15, 2.52)	0.02
		1.93 (0.69, 3.16)	0.00	1.67 (0.44, 2.89)	0.00	1.59 (0.35, 2.82)	0.01	1.53 (0.29, 2.77)	0.01	1.44 (0.19, 2.68)	0.02	1.36 (0.12, 2.60)	0.03
Nocturia	Physical anxiety	1.10 (-0.51, 2.70)	0.17	0.80 (-0.77, 2.38)	0.31	0.76 (-0.78, 2.31)	0.33	0.71 (-0.84, 2.26)	0.36	0.64 (-0.91, 2.20)	0.41	0.60 (-0.94, 2.15)	0.44
		1.94 (0.56, 3.31)	0.00	1.56 (0.20, 2.91)	0.02	1.58 (0.23, 2.92)	0.02	1.55 (0.21, 2.90)	0.02	1.48 (0.13, 2.83)	0.03	1.43 (0.08, 2.79)	0.03

  

Exposure	Outcome	Unadjusted		Adjusted 1		Adjusted 2		Adjusted 3		Adjusted 4		Adjusted 5	
		B (95% CI)	p	B (95% CI)	p	B (95% CI)	p	B (95% CI)	p	B (95% CI)	p	B (95% CI)	p
Daytime wetting	Mental anxiety	1.58 (0.65, 2.51)	0.00	1.41 (0.48, 2.33)	0.00	1.32 (0.40, 2.25)	0.00	1.31 (0.39, 2.24)	0.00	1.23 (0.30, 2.17)	0.01	1.08 (0.15, 2.02)	0.02
		0.65 (-0.52, 1.83)	0.27	0.70 (-0.48, 1.87)	0.24	0.60 (-0.57, 1.78)	0.31	0.60 (-0.57, 1.77)	0.31	0.59 (-0.58, 1.77)	0.32	0.37 (-0.80, 1.55)	0.53
Bedwetting	Mental anxiety	1.43 (0.59, 2.28)	0.00	1.29 (0.45, 2.14)	0.00	1.27 (0.42, 2.11)	0.00	1.27 (0.42, 2.12)	0.00	1.22 (0.37, 2.07)	0.00	1.17 (0.32, 2.02)	0.00
		1.00 (0.16, 1.83)	0.01	0.95 (0.12, 1.78)	0.02	0.85 (0.02, 1.69)	0.04	0.86 (0.03, 1.69)	0.04	0.79 (-0.05, 1.63)	0.06	0.67 (-0.16, 1.51)	0.11
Soiling	Mental anxiety	0.55 (-0.54, 1.64)	0.32	0.49 (-0.59, 1.58)	0.37	0.44 (-0.63, 1.50)	0.42	0.43 (-0.62, 1.49)	0.42	0.38 (-0.68, 1.44)	0.47	0.29 (-0.77, 1.34)	0.59
		1.60 (0.66, 2.54)	0.00	1.53 (0.58, 2.47)	0.00	1.50 (0.55, 2.44)	0.00	1.50 (0.55, 2.44)	0.00	1.44 (0.49, 2.39)	0.00	1.36 (0.41, 2.31)	0.00
Urgency	Mental anxiety	1.45 (0.99, 1.92)	<.00	1.43 (0.96, 1.89)	<.00	1.38 (0.92, 1.84)	<.00	1.39 (0.93, 1.85)	<.00	1.36 (0.90, 1.82)	<.00	1.31 (0.85, 1.78)	<.00
		0.52 (-0.18, 1.22)	0.14	0.49 (-0.20, 1.19)	0.16	0.44 (-0.25, 1.13)	0.20	0.45 (-0.25, 1.14)	0.20	0.42 (-0.27, 1.11)	0.23	0.34 (-0.35, 1.03)	0.33

  

Exposure	Outcome	Unadjusted		Adjusted 1		Adjusted 2		Adjusted 3		Adjusted 4		Adjusted 5	
		OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p
Daytime wetting	Self-harm thoughts	2.48 (1.58, 3.91)	<.00	2.16 (1.36, 3.42)	0.00	2.09 (1.30, 3.36)	0.00	2.05 (1.27, 3.30)	0.00	1.96 (1.21, 3.18)	0.00	1.83 (1.12, 2.98)	0.01
		1.36 (0.70, 2.66)	0.36	1.43 (0.73, 2.80)	0.30	1.31 (0.66, 2.61)	0.44	1.29 (0.64, 2.58)	0.47	1.26 (0.63, 2.53)	0.51	1.14 (0.56, 2.31)	0.71
Bedwetting	Self-harm thoughts	1.70 (1.11, 2.61)	0.01	1.52 (0.98, 2.35)	0.06	1.50 (0.96, 2.33)	0.07	1.45 (0.93, 2.25)	0.09	1.41 (0.90, 2.19)	0.13	1.37 (0.88, 2.14)	0.16
		1.19 (0.70, 2.03)	0.50	1.14 (0.67, 1.96)	0.62	1.01 (0.59, 1.73)	0.97	0.98 (0.57, 1.67)	0.93	0.93 (0.54, 1.62)	0.80	0.88 (0.51, 1.54)	0.66





Supplementary material for: *Prospective relationships between continence problems and common mental health disorders in adolescents from a UK cohort*

Bedwetting	Binge-eating	1.29 (0.73, 2.28)	0.38	1.40 (0.77, 2.53)	0.26	1.32 (0.72, 2.41)	0.36	1.31 (0.71, 2.40)	0.38	1.27 (0.69, 2.35)	0.43	1.15 (0.61, 2.14)	0.67
Soiling	Binge-eating	2.06 (1.35, 3.14)	0.00	1.71 (1.11, 2.64)	0.01	1.70 (1.10, 2.63)	0.01	1.65 (1.06, 2.58)	0.02	1.62 (1.05, 2.52)	0.03	1.58 (1.02, 2.46)	0.04
Urgency	Binge-eating	1.69 (1.05, 2.73)	0.03	1.60 (0.97, 2.63)	0.06	1.51 (0.91, 2.52)	0.10	1.47 (0.88, 2.46)	0.14	1.42 (0.85, 2.38)	0.17	1.36 (0.81, 2.28)	0.23
Frequent urination	Binge-eating	1.18 (0.66, 2.12)	0.58	1.09 (0.59, 2.00)	0.78	1.03 (0.55, 1.90)	0.93	1.00 (0.54, 1.85)	0.99	0.98 (0.53, 1.84)	0.96	0.94 (0.50, 1.75)	0.83
Low voided volume	Binge-eating	2.04 (1.31, 3.18)	0.00	1.88 (1.19, 2.97)	0.00	1.84 (1.17, 2.91)	0.00	1.83 (1.15, 2.90)	0.01	1.85 (1.17, 2.93)	0.00	1.79 (1.12, 2.85)	0.01
Voiding postponement	Binge-eating	1.43 (1.06, 1.94)	0.02	1.39 (1.02, 1.90)	0.03	1.36 (1.00, 1.87)	0.05	1.36 (0.99, 1.86)	0.05	1.35 (0.98, 1.85)	0.06	1.33 (0.97, 1.83)	0.08
Nocturia	Binge-eating	1.65 (1.18, 2.31)	0.00	1.61 (1.14, 2.29)	0.00	1.58 (1.11, 2.24)	0.01	1.54 (1.08, 2.19)	0.01	1.52 (1.07, 2.17)	0.02	1.45 (1.01, 2.07)	0.04
<b>Exposure</b>	<b>Outcome</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>
Daytime wetting	Excessive exercise	1.77 (1.14, 2.76)	0.01	1.34 (0.84, 2.14)	0.21	1.39 (0.86, 2.24)	0.18	1.46 (0.90, 2.37)	0.12	1.46 (0.90, 2.36)	0.12	1.40 (0.86, 2.27)	0.17
Bedwetting	Excessive exercise	1.43 (0.88, 2.32)	0.14	1.61 (0.95, 2.71)	0.07	1.76 (1.03, 3.00)	0.04	1.81 (1.06, 3.09)	0.03	1.80 (1.05, 3.09)	0.03	1.76 (1.01, 3.05)	0.04
Soiling	Excessive exercise	1.37 (0.95, 1.97)	0.09	1.09 (0.75, 1.58)	0.65	1.08 (0.74, 1.57)	0.69	1.15 (0.78, 1.68)	0.48	1.14 (0.77, 1.67)	0.51	1.11 (0.76, 1.64)	0.58
Urgency	Excessive exercise	0.87 (0.57, 1.33)	0.52	0.79 (0.51, 1.23)	0.30	0.84 (0.54, 1.32)	0.44	0.89 (0.56, 1.40)	0.60	0.88 (0.56, 1.39)	0.59	0.87 (0.55, 1.37)	0.54
Frequent urination	Excessive exercise	1.21 (0.74, 1.99)	0.44	1.11 (0.66, 1.87)	0.68	1.18 (0.70, 2.01)	0.53	1.25 (0.72, 2.15)	0.42	1.24 (0.72, 2.14)	0.43	1.21 (0.70, 2.11)	0.49
Low voided volume	Excessive exercise	1.63 (1.08, 2.45)	0.02	1.48 (0.95, 2.31)	0.08	1.53 (0.98, 2.39)	0.06	1.59 (1.01, 2.49)	0.04	1.58 (1.00, 2.49)	0.04	1.56 (0.98, 2.46)	0.05
Voiding postponement	Excessive exercise	1.35 (1.05, 1.75)	0.01	1.32 (1.01, 1.72)	0.04	1.35 (1.03, 1.78)	0.03	1.35 (1.02, 1.78)	0.03	1.34 (1.02, 1.78)	0.03	1.36 (1.03, 1.80)	0.03
Nocturia	Excessive exercise	1.56 (1.16, 2.11)	0.00	1.54 (1.12, 2.11)	0.00	1.62 (1.17, 2.23)	0.00	1.69 (1.22, 2.33)	0.00	1.69 (1.22, 2.33)	0.00	1.64 (1.18, 2.28)	0.00
<b>Exposure</b>	<b>Outcome</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>
Daytime wetting	DSM-5 DE	2.35 (1.42, 3.89)	0.00	1.88 (1.12, 3.14)	0.01	1.99 (1.17, 3.39)	0.01	2.06 (1.20, 3.54)	0.00	2.03 (1.18, 3.49)	0.01	1.83 (1.04, 3.20)	0.03
Bedwetting	DSM-5 DE	1.39 (0.72, 2.70)	0.32	1.51 (0.76, 3.00)	0.23	1.47 (0.73, 2.98)	0.28	1.47 (0.72, 2.97)	0.28	1.44 (0.71, 2.92)	0.31	1.25 (0.60, 2.61)	0.54
Soiling	DSM-5 DE	2.13 (1.37, 3.32)	0.00	1.79 (1.13, 2.82)	0.01	1.85 (1.16, 2.94)	0.01	1.81 (1.13, 2.91)	0.01	1.78 (1.10, 2.86)	0.01	1.72 (1.07, 2.78)	0.02
Urgency	DSM-5 DE	1.44 (0.87, 2.37)	0.15	1.35 (0.81, 2.26)	0.24	1.21 (0.71, 2.04)	0.47	1.17 (0.69, 1.98)	0.55	1.15 (0.68, 1.95)	0.59	1.09 (0.64, 1.88)	0.74
Frequent urination	DSM-5 DE	2.26 (1.27, 4.01)	0.00	2.15 (1.18, 3.89)	0.01	2.01 (1.08, 3.71)	0.02	1.97 (1.06, 3.64)	0.03	1.94 (1.05, 3.60)	0.03	1.85 (0.99, 3.47)	0.05
Low voided volume	DSM-5 DE	1.85 (1.11, 3.08)	0.01	1.70 (1.00, 2.89)	0.04	1.66 (0.97, 2.85)	0.06	1.68 (0.98, 2.88)	0.06	1.66 (0.97, 2.86)	0.06	1.59 (0.91, 2.77)	0.09
Voiding postponement	DSM-5 DE	1.52 (1.08, 2.14)	0.01	1.48 (1.05, 2.09)	0.02	1.45 (1.02, 2.06)	0.03	1.43 (1.01, 2.03)	0.04	1.42 (1.00, 2.02)	0.05	1.40 (0.98, 2.01)	0.06

Supplementary material for: *Prospective relationships between continence problems and common mental health disorders in adolescents from a UK cohort*

Nocturia	DSM-5 DE	1.56 (1.06, 2.32)	0.02 6	1.53 (1.02, 2.29)	0.04 1	1.43 (0.94, 2.16)	0.09 1	1.39 (0.92, 2.11)	0.12 1	1.38 (0.91, 2.09)	0.13 1	1.28 (0.83, 1.97)	0.25 6
----------	----------	-------------------	-----------	-------------------	-----------	-------------------	-----------	-------------------	-----------	-------------------	-----------	-------------------	-----------

DSM-5 DE- Disordered eating at Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition frequency (at least once a week).

Adjusted 1 confounders: sex. Adjusted 2 confounders: sex and socioeconomic indicators (parental occupational social class, maternal educational attainment, family size, ethnicity, home ownership status, material hardship). Adjusted 3 confounders: sex, socioeconomic indicators, child IQ and developmental level. Adjusted 4 confounders: sex, socioeconomic indicators, child IQ, developmental level, maternal stressful life events, maternal depression and maternal anxiety. Adjusted 5 confounders: sex, socioeconomic indicators, child IQ, developmental level, maternal stressful life events, maternal depression, maternal anxiety, child Body Mass Index and earlier emotional/behaviour problems (fully adjusted model).

**Table S9. Sequential adjustments for secondary outcomes using complete case data (mental health outcomes n=1,528; disordered eating outcomes n=1,375)**

Exposure	Outcome	Unadjusted		Adjusted 1		Adjusted 2		Adjusted 3		Adjusted 4		Adjusted 5	
		B (95% CI)	p	B (95% CI)	p	B (95% CI)	p	B (95% CI)	p	B (95% CI)	p	B (95% CI)	p
Daytime wetting	Physical anxiety	1.87 (-0.16, 3.89)	0.07	1.18 (-0.80, 3.17)	0.24	1.14 (-0.85, 3.12)	0.26	1.09 (-0.89, 3.08)	0.28	1.07 (-0.91, 3.06)	0.28	1.12 (-0.87, 3.10)	0.27
		-1.28 (-3.67, 1.11)	0.29	-0.88 (-3.21, 1.46)	0.46	-1.15 (-3.49, 1.19)	0.33	-1.17 (-3.51, 1.18)	0.32	-1.13 (-3.48, 1.21)	0.34	-1.23 (-3.58, 1.11)	0.30
Bedwetting	Physical anxiety	1.13 (-0.69, 2.94)	0.22	0.47 (-1.31, 2.25)	0.60	0.45 (-1.34, 2.23)	0.62	0.35 (-1.44, 2.14)	0.70	0.35 (-1.44, 2.14)	0.70	0.35 (-1.44, 2.13)	0.70
			0.01		0.01		0.02		0.02		0.02		0.02
Urgency	Physical anxiety	2.50 (0.60, 4.39)	0	2.29 (0.44, 4.14)	5	2.11 (0.25, 3.96)	6	2.09 (0.24, 3.95)	7	2.08 (0.22, 3.94)	9	2.11 (0.25, 3.97)	6
		0.31 (-2.14, 2.76)	0.80	0.10 (-2.30, 2.50)	0.93	0.04 (-2.36, 2.44)	0.97	0.00 (-2.40, 2.40)	0.99	-0.02 (-2.43, 2.38)	0.98	0.01 (-2.39, 2.41)	0.99
Frequent urination	Physical anxiety	1.38 (-0.47, 3.22)	0.14	0.99 (-0.81, 2.80)	0.27	1.10 (-0.70, 2.90)	0.23	1.05 (-0.75, 2.86)	0.25	1.01 (-0.80, 2.81)	0.27	0.93 (-0.87, 2.74)	0.31
			0.00		0.00		0.00		0.00		0.00		0.01
Low voided volume Voiding postponement	Physical anxiety	1.65 (0.48, 2.82)	6	1.60 (0.45, 2.74)	6	1.56 (0.42, 2.71)	8	1.59 (0.44, 2.74)	7	1.56 (0.41, 2.71)	8	1.52 (0.37, 2.67)	0
		-1.04 (-0.29, 2.37)	0.12	1.05 (-0.25, 2.35)	0.11	1.06 (-0.24, 2.36)	0.10	1.02 (-0.28, 2.32)	0.12	1.03 (-0.27, 2.33)	0.12	1.03 (-0.27, 2.33)	0.12
Nocturia	Physical anxiety	2.37	5	2.35	3	2.36	9	2.32	4	2.33	1	2.33	0
Exposure	Outcome	B (95% CI)	p	B (95% CI)	p	B (95% CI)	p	B (95% CI)	p	B (95% CI)	p	B (95% CI)	p
Daytime wetting	Mental anxiety	0.97 (-0.32, 2.27)	0.14	0.93 (-0.37, 2.23)	0.15	0.87 (-0.43, 2.17)	0.18	0.87 (-0.43, 2.17)	0.19	0.84 (-0.46, 2.14)	0.20	0.80 (-0.50, 2.10)	0.22
		-0.70 (-2.23, 0.83)	0.37	-0.67 (-2.21, 0.86)	0.38	-0.80 (-2.33, 0.74)	0.30	-0.79 (-2.32, 0.75)	0.31	-0.77 (-2.30, 0.76)	0.32	-0.94 (-2.47, 0.59)	0.23
Bedwetting	Mental anxiety	1.09 (-0.07, 2.25)	0.06	1.06 (-0.11, 2.22)	0.07	1.03 (-0.13, 2.20)	0.08	1.03 (-0.14, 2.20)	0.08	1.03 (-0.14, 2.20)	0.08	0.96 (-0.20, 2.13)	0.10
			0.02		0.02		0.02		0.02		0.03		0.03
Urgency	Mental anxiety	1.39 (0.18, 2.60)	4	1.38 (0.17, 2.59)	6	1.36 (0.15, 2.58)	8	1.36 (0.15, 2.58)	8	1.33 (0.12, 2.55)	2	1.29 (0.08, 2.51)	7
		0.33 (-1.24, 1.90)	0.68	0.31 (-1.26, 1.88)	0.69	0.35 (-1.22, 1.92)	0.66	0.35 (-1.22, 1.93)	0.65	0.33 (-1.24, 1.90)	0.68	0.29 (-1.28, 1.85)	0.72
Frequent urination	Mental anxiety	1.29 (0.11, 2.46)	2	1.26 (0.09, 2.44)	6	1.25 (0.07, 2.43)	8	1.23 (0.05, 2.41)	1	1.19 (0.01, 2.37)	8	1.11 (-0.07, 2.29)	0.06
			0.00		0.00		0.00		0.00		0.00		0.00
Low voided volume Voiding postponement	Mental anxiety	1.12 (0.37, 1.87)	3	1.12 (0.37, 1.87)	3	1.08 (0.33, 1.84)	5	1.09 (0.34, 1.85)	4	1.05 (0.30, 1.81)	6	1.02 (0.27, 1.78)	8
		0.46 (-0.39, 1.31)	0.29	0.46 (-0.39, 1.31)	0.29	0.45 (-0.40, 1.30)	0.30	0.45 (-0.40, 1.30)	0.30	0.45 (-0.40, 1.30)	0.29	0.42 (-0.43, 1.27)	0.32
Nocturia	Mental anxiety	1.31	4	1.31	3	1.30	0	1.30	0	1.30	8	1.27	8
Exposure	Outcome	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p
Daytime wetting	Self-harm thoughts	1.94 (0.93, 4.05)	0.07	1.69 (0.80, 3.54)	0.16	1.74 (0.82, 3.68)	0.14	1.72 (0.81, 3.63)	0.15	1.69 (0.80, 3.58)	0.17	1.63 (0.76, 3.49)†	0.20
			8		8		7		9		1		8
Bedwetting	Self-harm thoughts	1.43 (0.55, 3.71)	0.46	1.58 (0.60, 4.15)	0.35	1.59 (0.60, 4.22)	0.35	1.58 (0.59, 4.21)	0.36	1.60 (0.60, 4.27)	0.34	1.39 (0.51, 3.74)†	0.52
			4		0		2		1		8		0
Soiling	Self-harm thoughts	2.35 (1.25, 4.42)	0.00	2.06 (1.09, 3.90)	0.02	2.02 (1.06, 3.85)	0.03	1.98 (1.04, 3.78)	0.03	2.00 (1.05, 3.83)	0.03	1.98 (1.03, 3.81)†	0.04
			8		5		2		9		5		0
Urgency	Self-harm thoughts	1.21 (0.54, 2.71)	0.64	1.16 (0.52, 2.61)	0.72	1.11 (0.49, 2.51)	0.80	1.11 (0.49, 2.51)	0.80	1.08 (0.47, 2.46)	0.85	1.05 (0.46, 2.40)†	0.90
			0		0		3		7		5		8

Supplementary material for: *Prospective relationships between continence problems and common mental health disorders in adolescents from a UK cohort*

Exposure	Outcome	Unadjusted		Adjusted 1		Adjusted 2		Adjusted 3		Adjusted 4		Adjusted 5	
		OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p
Frequent urination	Self-harm thoughts	1.52 (0.58, 3.96)	0.39	1.46 (0.55, 3.83)	0.44	1.42 (0.53, 3.79)	0.48	1.40 (0.52, 3.76)	0.50	1.42 (0.53, 3.80)	0.49	1.36 (0.50, 3.68)†	0.54
Low voided volume	Self-harm thoughts	1.52 (0.74, 3.14)	0.25	1.41 (0.68, 2.92)	0.35	1.42 (0.68, 2.97)	0.34	1.39 (0.66, 2.90)	0.38	1.37 (0.65, 2.87)	0.40	1.26 (0.60, 2.65)†	0.54
Voiding postponement	Self-harm thoughts	1.42 (0.88, 2.30)	0.15	1.41 (0.87, 2.29)	0.16	1.38 (0.84, 2.25)	0.20	1.39 (0.85, 2.28)	0.18	1.35 (0.82, 2.22)	0.23	1.30 (0.79, 2.15)†	0.29
Nocturia	Self-harm thoughts	1.90 (1.15, 3.14)	0.01	1.92 (1.16, 3.19)	0.01	1.94 (1.16, 3.23)	0.01	1.92 (1.15, 3.20)	0.01	1.90 (1.14, 3.18)	0.01	1.90 (1.13, 3.19)†	0.01
<b>Exposure</b>	<b>Outcome</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>
Daytime wetting	Fasting	1.01 (0.39, 2.60)	0.98	0.80 (0.31, 2.08)	0.64	0.85 (0.33, 2.22)	0.73	0.78 (0.30, 2.06)	0.62	0.74 (0.28, 1.97)	0.55	0.64 (0.24, 1.71)♦	0.37
Bedwetting	Fasting	1.35 (0.52, 3.52)	0.54	1.49 (0.55, 4.00)	0.43	1.50 (0.55, 4.06)	0.42	1.40 (0.51, 3.84)	0.51	1.42 (0.51, 3.91)	0.50	1.10 (0.38, 3.13)♦	0.86
Soiling	Fasting	1.85 (0.96, 3.53)	0.06	1.62 (0.84, 3.14)	0.15	1.73 (0.88, 3.38)	0.11	1.57 (0.80, 3.09)	0.18	1.57 (0.80, 3.10)	0.19	1.58 (0.79, 3.13)♦	0.19
Urgency	Fasting	2.13 (1.13, 4.01)	0.01	2.07 (1.08, 3.97)	0.02	2.09 (1.08, 4.05)	0.02	1.92 (0.98, 3.74)	0.05	1.85 (0.94, 3.63)	0.07	1.87 (0.94, 3.72)♦	0.07
Frequent urination	Fasting	1.60 (0.70, 3.65)	0.26	1.58 (0.68, 3.68)	0.29	1.54 (0.65, 3.63)	0.32	1.41 (0.59, 3.36)	0.43	1.40 (0.58, 3.35)	0.45	1.30 (0.54, 3.13)♦	0.56
Low voided volume	Fasting	1.74 (0.89, 3.42)	0.10	1.62 (0.81, 3.22)	0.17	1.65 (0.82, 3.30)	0.15	1.54 (0.76, 3.09)	0.23	1.56 (0.77, 3.17)	0.21	1.48 (0.72, 3.01)♦	0.28
Voiding postponement	Fasting	1.44 (0.91, 2.27)	0.11	1.41 (0.88, 2.24)	0.14	1.48 (0.92, 2.36)	0.10	1.48 (0.92, 2.39)	0.10	1.43 (0.89, 2.32)	0.14	1.39 (0.86, 2.26)♦	0.17
Nocturia	Fasting	1.58 (0.92, 2.69)	0.09	1.63 (0.94, 2.82)	0.08	1.57 (0.90, 2.74)	0.11	1.50 (0.85, 2.62)	0.16	1.47 (0.84, 2.58)	0.17	1.40 (0.79, 2.49)♦	0.24
<b>Exposure</b>	<b>Outcome</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>
Daytime wetting	Purging	2.16 (0.83, 5.64)	0.11	1.74 (0.66, 4.59)	0.26	1.79 (0.67, 4.76)	0.24	1.81 (0.68, 4.83)	0.23	1.73 (0.64, 4.64)	0.27	1.58 (0.58, 4.30)♦	0.37
Bedwetting	Purging	1.58 (0.47, 5.26)	0.45	1.73 (0.51, 5.90)	0.38	1.60 (0.46, 5.53)	0.46	1.65 (0.48, 5.72)	0.43	1.52 (0.44, 5.32)	0.51	1.30 (0.36, 4.65)♦	0.68
Soiling	Purging	2.06 (0.91, 4.67)	0.08	1.81 (0.79, 4.14)	0.16	1.86 (0.80, 4.31)	0.14	1.87 (0.80, 4.36)	0.14	1.91 (0.82, 4.47)	0.13	1.89 (0.80, 4.45)♦	0.14
Urgency	Purging	1.77 (0.74, 4.24)	0.20	1.69 (0.70, 4.08)	0.24	1.62 (0.66, 3.96)	0.29	1.60 (0.65, 3.96)	0.30	1.55 (0.63, 3.85)	0.34	1.55 (0.62, 3.88)♦	0.34
Frequent urination	Purging	1.26 (0.38, 4.17)	0.70	1.23 (0.37, 4.11)	0.74	1.14 (0.33, 3.86)	0.83	1.14 (0.33, 3.88)	0.83	1.10 (0.32, 3.80)	0.88	1.01 (0.29, 3.53)♦	0.98
Low voided volume	Purging	2.17 (0.96, 4.93)	0.06	2.01 (0.88, 4.63)	0.09	2.08 (0.90, 4.82)	0.08	2.10 (0.91, 4.89)	0.08	2.11 (0.90, 4.92)	0.08	1.98 (0.84, 4.64)♦	0.11
Voiding postponement	Purging	1.36 (0.73, 2.53)	0.32	1.32 (0.71, 2.47)	0.38	1.34 (0.72, 2.52)	0.35	1.33 (0.71, 2.50)	0.37	1.25 (0.66, 2.37)	0.49	1.22 (0.64, 2.31)♦	0.54
Nocturia	Purging	2.13 (1.12, 4.08)	0.02	2.20 (1.14, 4.25)	0.01	2.11 (1.08, 4.11)	0.02	2.11 (1.08, 4.13)	0.02	2.08 (1.06, 4.08)	0.03	2.01 (1.02, 3.97)♦	0.04
<b>Exposure</b>	<b>Outcome</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>	<b>OR (95% CI)</b>	<b>p</b>
Daytime wetting	Binge-eating	3.14 (1.62, 6.11)	0.00	2.66 (1.35, 5.23)	0.00	2.74 (1.38, 5.43)	0.00	2.79 (1.40, 5.54)	0.00	2.71 (1.36, 5.41)	0.00	2.39 (1.18, 4.83)	0.01

Supplementary material for: *Prospective relationships between continence problems and common mental health disorders in adolescents from a UK cohort*

		Unadjusted		Adjusted 1		Adjusted 2		Adjusted 3		Adjusted 4		Adjusted 5	
Bedwetting	Binge-eating	0.94 (0.33, 2.70)	0.91	1.00 (0.35, 2.92)	0.99	0.96 (0.33, 2.81)	0.94	0.98 (0.33, 2.86)	0.96	0.98 (0.33, 2.86)	0.96	0.75 (0.25, 2.25)	0.60
Soiling	Binge-eating	1.16 (0.56, 2.38)	0.68	1.03 (0.49, 2.13)	0.94	1.02 (0.49, 2.12)	0.96	1.02 (0.49, 2.14)	0.96	1.02 (0.48, 2.13)	0.96	0.96 (0.45, 2.02)	0.90
Urgency	Binge-eating	1.20 (0.58, 2.48)	0.61	1.15 (0.55, 2.39)	0.71	1.15 (0.55, 2.41)	0.70	1.14 (0.54, 2.40)	0.72	1.11 (0.52, 2.34)	0.78	1.07 (0.50, 2.27)	0.86
Frequent urination	Binge-eating	1.20 (0.50, 2.88)	0.68	1.17 (0.48, 2.85)	0.72	1.20 (0.49, 2.93)	0.69	1.20 (0.49, 2.95)	0.68	1.19 (0.48, 2.92)	0.70	1.12 (0.45, 2.77)	0.80
Low voided volume	Binge-eating	2.39 (1.31, 4.37)	0.00	2.27 (1.23, 4.20)	0.00	2.27 (1.22, 4.21)	0.00	2.31 (1.24, 4.29)	0.00	2.35 (1.26, 4.40)	0.00	2.27 (1.21, 4.28)	0.01
Voiding postponement	Binge-eating	1.29 (0.82, 2.02)	0.27	1.26 (0.79, 1.99)	0.32	1.27 (0.80, 2.02)	0.30	1.26 (0.79, 2.00)	0.32	1.21 (0.76, 1.93)	0.41	1.20 (0.75, 1.91)	0.45
Nocturia	Binge-eating	1.52 (0.90, 2.57)	0.11	1.56 (0.92, 2.65)	0.10	1.57 (0.92, 2.68)	0.09	1.57 (0.92, 2.69)	0.09	1.56 (0.91, 2.67)	0.10	1.49 (0.86, 2.56)	0.15
			3		2		8		7		4		4
Exposure	Outcome	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p
Daytime wetting	Excessive exercise	2.07 (1.12, 3.82)	0.02	1.69 (0.90, 3.18)	0.10	1.69 (0.89, 3.19)	0.10	1.81 (0.95, 3.43)	0.06	1.81 (0.96, 3.44)	0.06	1.74 (0.91, 3.34)	0.09
Bedwetting	Excessive exercise	1.15 (0.53, 2.47)	0.72	1.26 (0.57, 2.79)	0.56	1.27 (0.57, 2.83)	0.56	1.32 (0.59, 2.96)	0.49	1.31 (0.59, 2.95)	0.50	1.21 (0.53, 2.78)	0.64
Soiling	Excessive exercise	1.25 (0.72, 2.18)	0.42	1.09 (0.62, 1.93)	0.76	1.05 (0.59, 1.87)	0.86	1.13 (0.63, 2.02)	0.67	1.13 (0.63, 2.03)	0.67	1.09 (0.60, 1.96)	0.77
Urgency	Excessive exercise	0.75 (0.40, 1.43)	0.38	0.70 (0.36, 1.34)	0.28	0.72 (0.37, 1.39)	0.32	0.76 (0.39, 1.48)	0.42	0.77 (0.40, 1.49)	0.43	0.77 (0.39, 1.51)	0.44
Frequent urination	Excessive exercise	1.15 (0.57, 2.30)	0.69	1.12 (0.55, 2.30)	0.75	1.15 (0.56, 2.36)	0.71	1.21 (0.58, 2.50)	0.61	1.21 (0.58, 2.50)	0.61	1.15 (0.55, 2.40)	0.71
Low voided volume	Excessive exercise	1.56 (0.91, 2.69)	0.10	1.46 (0.83, 2.56)	0.18	1.44 (0.82, 2.53)	0.20	1.54 (0.87, 2.72)	0.14	1.52 (0.86, 2.70)	0.15	1.45 (0.81, 2.58)	0.21
Voiding postponement	Excessive exercise	1.42 (1.00, 2.02)	0.04	1.41 (0.98, 2.02)	0.06	1.40 (0.97, 2.01)	0.07	1.38 (0.96, 1.99)	0.08	1.38 (0.95, 1.99)	0.08	1.37 (0.94, 1.99)	0.09
Nocturia	Excessive exercise	1.39 (0.91, 2.13)	0.12	1.44 (0.93, 2.24)	0.10	1.48 (0.95, 2.31)	0.08	1.55 (0.99, 2.42)	0.05	1.55 (0.99, 2.43)	0.05	1.55 (0.99, 2.44)	0.05
			7		5		2		5		3		7
Exposure	Outcome	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p
Daytime wetting	DSM-5 DE	2.03 (0.93, 4.44)	0.07	1.66 (0.75, 3.69)	0.20	1.83 (0.82, 4.12)	0.14	1.82 (0.81, 4.10)	0.15	1.80 (0.80, 4.06)	0.15	1.50 (0.65, 3.47)	0.34
Bedwetting	DSM-5 DE	0.85 (0.26, 2.79)	0.78	0.91 (0.27, 3.04)	0.87	0.87 (0.26, 2.98)	0.83	0.88 (0.26, 3.00)	0.83	0.88 (0.26, 3.02)	0.84	0.64 (0.18, 2.28)	0.49
Soiling	DSM-5 DE	1.87 (0.96, 3.67)	0.06	1.66 (0.84, 3.29)	0.14	1.74 (0.86, 3.51)	0.12	1.69 (0.83, 3.42)	0.14	1.68 (0.83, 3.42)	0.14	1.58 (0.77, 3.23)	0.21
Urgency	DSM-5 DE	1.11 (0.50, 2.49)	0.79	1.06 (0.47, 2.38)	0.89	1.04 (0.45, 2.39)	0.92	0.98 (0.42, 2.26)	0.95	0.96 (0.41, 2.24)	0.92	0.93 (0.40, 2.18)	0.87
Frequent urination	DSM-5 DE	2.14 (0.98, 4.71)	0.05	2.14 (0.95, 4.79)	0.06	2.23 (0.98, 5.08)	0.05	2.19 (0.96, 5.02)	0.06	2.16 (0.94, 4.97)	0.06	2.09 (0.91, 4.83)	0.08
Low voided volume	DSM-5 DE	1.75 (0.87, 3.53)	0.11	1.64 (0.80, 3.33)	0.17	1.68 (0.81, 3.45)	0.16	1.67 (0.81, 3.45)	0.16	1.65 (0.80, 3.41)	0.17	1.53 (0.73, 3.24)	0.26
			5		5		1		8		9		0

		<b>Unadjusted</b>		<b>Adjusted 1</b>		<b>Adjusted 2</b>		<b>Adjusted 3</b>		<b>Adjusted 4</b>		<b>Adjusted 5</b>	
Voiding postponement	DSM-5 DE	1.24 (0.76, 2.03)	0.39	1.21 (0.73, 1.99)	0.46	1.27 (0.76, 2.13)	0.35	1.25 (0.75, 2.09)	0.38	1.20 (0.71, 2.01)	0.49	1.17 (0.70, 1.98)	0.54
			2		2		2		7		4		7
			0.23		0.21		0.25		0.27		0.28		0.39
Nocturia	DSM-5 DE	1.42 (0.80, 2.52)	1	1.45 (0.81, 2.60)	3	1.42 (0.78, 2.58)	1	1.40 (0.77, 2.55)	2	1.39 (0.76, 2.54)	2	1.31 (0.71, 2.42)	0

DSM-5 DE- Disordered eating at Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition frequency (at least once a week). †Analyses performed on n=1494 due to 34 observations being dropped (underweight BMI predicted outcome perfectly). ♦Analyses performed on n=1349 due to 26 observations being dropped (underweight BMI predicted outcome perfectly).

Adjusted 1 confounders: sex. Adjusted 2 confounders: sex and socioeconomic indicators (parental occupational social class, maternal educational attainment, family size, ethnicity, home ownership status, material hardship). Adjusted 3 confounders: sex, socioeconomic indicators, child IQ and developmental level. Adjusted 4 confounders: sex, socioeconomic indicators, child IQ, developmental level, maternal stressful life events, maternal depression and maternal anxiety. Adjusted 5 confounders: sex, socioeconomic indicators, child IQ, developmental level, maternal stressful life events, maternal depression, maternal anxiety, child Body Mass Index and earlier emotional/behaviour problems (fully adjusted model).