

## PICRUSt Pathway Abundance Analysis Significant Results

Pathway	Taxa w/pathway (per MetaCyc)	Adjusted p-value
6-hydroxymethyl-dihydropterin diphosphate biosynthesis III (Chlamydia)	<i>Chlamydia trachomatis</i>	0.029
8-amino-7-oxononanoate biosynthesis I	<i>Candidatus Nitrososphaera gargensis Ga9.2</i> <i>Escherichia coli K-12 substr. MG1655</i> <i>Francisella tularensis novicida U112</i> <i>Haemophilus influenzae Rd KW20</i> <i>Helicobacter pylori 26695</i> <i>Mycobacterium tuberculosis H37Rv</i> <i>Prochlorococcus marinus MIT 9211</i>	0.007
biotin biosynthesis I	<i>Escherichia coli K-12 substr. MG1655</i> <i>Francisella tularensis novicida U112</i> <i>Haemophilus influenzae Rd KW20</i> <i>Helicobacter pylori 26695</i> <i>Prochlorococcus marinus MIT 9211</i>	0.007
CMP-3-deoxy-D-manno-octulosonate biosynthesis I	<i>Arabidopsis thaliana col</i> <i>Escherichia coli BL21(DE3)</i> <i>Escherichia coli K-12 substr. MG1655</i> <i>Zea mays</i>	0.003
flavin biosynthesis I (bacteria and plants)	<i>Arabidopsis thaliana col</i> <i>Bacillus subtilis</i> <i>Escherichia coli K-12 substr. MG1655</i>	0.029
fucose degradation	<i>Escherichia coli K-12 substr. MG1655</i>	0.045
Kdo transfer to lipid IVA III (Chlamydia)	<i>Chlamydia pneumoniae</i> <i>Chlamydia psittaci</i> <i>Chlamydia psittaci 6BC</i>	0.003
L-lysine biosynthesis I	<i>Arthrobacter globiformis</i> <i>Azotobacter vinelandii</i> <i>Bordetella pertussis</i> <i>Corynebacterium glutamicum</i> <i>Corynebacterium glutamicum ATCC 13032</i> <i>Escherichia coli K-12 substr. MG1655</i> <i>Haemophilus influenzae</i> <i>Haemophilus influenzae Rd KW20</i> <i>Helicobacter pylori</i> <i>Mycobacterium tuberculosis</i> <i>Mycobacterium tuberculosis H37Rv</i> <i>Rhodospirillum rubrum</i>	0.034
L-lysine biosynthesis VI	<i>Arabidopsis thaliana col</i> <i>Chlamydia trachomatis</i> <i>Glycine max</i> <i>Methanocaldococcus jannaschii</i> <i>Methanothermobacter thermautotrophicus</i> <i>Nicotiana tabacum</i> <i>Synechocystis</i> <i>Zea mays</i>	0.026
lipid IVA biosynthesis	<i>Caulobacter vibrioides NA1000</i> <i>Escherichia coli K-12 substr. MG1655</i>	0.003
methylerythritol phosphate pathway I	<i>Brucella abortus</i> <i>Escherichia coli K-12 substr. MG1655</i>	0.005
methylerythritol phosphate pathway II	<i>Arabidopsis thaliana col</i> <i>Botryococcus braunii</i> <i>Croton stellatopilosus</i> <i>Plasmodium falciparum</i> <i>Plasmodium falciparum HB3</i> <i>Solanum lycopersicum</i> <i>Thermosynechococcus vestitus BP-1</i>	0.005

mycolate biosynthesis	<i>Mycobacterium tuberculosis H37Rv</i>	0.023
NAD salvage pathway II	<i>Escherichia coli K-12 substr. MG1655</i> <i>Salmonella enterica enterica serovar Typhimurium</i>	0.009
oleate biosynthesis IV (anaerobic)	<i>Aerococcus viridans</i> <i>Cereibacter sphaerooides 2.4.1</i> <i>Clostridium beijerinckii</i>	0.023
palmitate biosynthesis II (bacteria and plants)	<i>Arabidopsis thaliana col</i> <i>Brassica napus</i> <i>Escherichia coli K-12 substr. MG1655</i> <i>Spinacia oleracea</i>	0.025
palmitoleate biosynthesis I (from (5Z)-dodec-5-enoate)	<i>Escherichia coli K-12 substr. MG1655</i> <i>Helicobacter pylori 26695</i>	0.003
stearate biosynthesis II (bacteria and plants)	<i>Arabidopsis thaliana col</i> <i>Brassica napus</i> <i>Spinacia oleracea</i>	0.012
superpathway of (Kdo)2-lipid A biosynthesis	<i>Escherichia coli K-12 substr. MG1655</i>	0.028
superpathway of fatty acid biosynthesis initiation (E. coli)	<i>Escherichia coli K-12 substr. MG1655</i>	0.030

**Table, Supplemental Digital Content 5.** PICRUSt2 inferred pathways and associated species. These pathway abundances are all significantly different between RSV+ and RSV- samples. Species identified as being differentially expressed in RSV+ and RSV- decedents in our differential abundance analysis are bolded.