

Title: Understanding disruptions in cancer care to reduce increased cancer burden

Running title: Disruptions in cancer care during COVID-19

Authors: Kia L. Davis, ScD, MPH¹, Nicole Ackermann, MPH¹, Lisa M. Klesges, PhD¹, Nora Leahy, MPH, CHES¹, Callie, Walsh-Bailey, MPH², Sarah Humble, MS¹, Bettina Drake, PhD, MPH¹, Vetta L. Sanders Thompson, PhD²

¹ Washington University in St. Louis, School of Medicine, Department of Surgery, Public Health Sciences, St. Louis, MO, United States

² Washington University in St. Louis, Brown School, St. Louis MO, United States

Corresponding Author: Kia Davis, ScD, MPH, Division of Public Health Sciences, Department of Surgery, Washington University School of Medicine, 660 S. Euclid Ave. Campus Box 8100, St. Louis, MO 63110, phone: 314-747-8069, email: daviskl@wustl.edu

Conflict of Interest Disclosures: The authors declare no potential conflicts of interest.

Abstract

Background: This study seeks to understand how and for whom COVID-19 disrupted cancer care to understand the potential for cancer health disparities across the cancer prevention and control continuum.

Methods: In this cross-sectional study, participants age 30+ residing in an 82-county region in Missouri and Illinois completed an online survey from June-August 2020. Descriptive statistics were calculated for all variables separately and by care disruption status. Logistic regression modeling was conducted to determine the correlates of care disruption.

Results: Participants (N=680) reported 21% to 57% of cancer screening or treatment appointments were canceled from March 2020 through the end of 2020. Approximately 34% of residents stated they would need to know if their doctor's office is taking the appropriate COVID-related safety precautions to return to care. Higher education (OR=1.26, 95% CI:1.11-1.43), identifying as female (OR=1.60, 95% CI:1.12-2.30), experiencing more discrimination in healthcare settings (OR= 1.40, 95% CI:1.13-1.72), and having scheduled a telehealth appointment (OR=1.51, 95% CI:1.07-2.15) were associated with higher odds of care disruption. Factors associated with care disruption were not consistent across races. Higher odds of care disruption for White residents were associated with higher education, female identity, older age, and having scheduled a telehealth appointment, while higher odds of care disruption for Black residents were associated only with higher education.

Conclusion(s): This study provides an understanding of the factors associated with cancer care disruption and what patients need to return to care. Results may inform outreach and engagement strategies to reduce delayed cancer screenings and encourage returning to cancer care.

Funding Support: This study was supported by the National Cancer Institute's Administrative Supplements for P30 Cancer Center Support Grants (P30CA091842-18S2 and P30CA091842-19S4). Kia L. Davis, Lisa Klesges, and Bettina Drake were supported by the National Cancer Institute's P50CA244431 and Kia L. Davis was also supported by the Breast Cancer Research Foundation. Callie Walsh-Bailey was supported by NIMHD T37 MD014218. The content does not necessarily represent the official view of these funding agencies and is solely the responsibility of the authors.

Availability of data and material

The dataset generated for the study is not publicly available but is available by request. Interested individuals should contact the corresponding author with a brief description of how the data will be used and proof of IRB approval or exemption. Then a de-identified dataset will be shared.

Key Words: Early detection of cancer, COVID-19 pandemic, Cancer health disparities, Social determinants of health, Community health, Delivery of health care, Epidemiology, Access to care

It is made available under a [CC-BY-NC 4.0 International license](#) .

Disruptions in cancer care during COVID-19

Introduction

The COVID-19 pandemic abruptly upended cancer care in many countries including the US. The need to reduce community spread and reserve hospital capacity for the most severe COVID-19 cases led to rescheduling or postponement of cancer care appointments.¹⁻⁵ These control measures significantly decreased cancer-related patient encounters in the early phase of the pandemic, particularly for cancer screenings.² Comparing March to July 2020 with the same period in 2019, there was a substantial decrease in cancer screenings, biopsies, surgeries, office visits, and therapy; the decreases varied by service location and cancer type.² For example, breast cancer screenings decreased by 89.2% and colorectal by 84.5%.⁶ Patients reported delays in receiving cancer care, including follow-up clinic appointments and cancer therapies, such as radiation, infusion therapies, and surgeries.^{7,8}

Cancer care delays due to the COVID-19 pandemic are anticipated to lead to increased cancer morbidity and mortality.^{9,10} One study found an association between surgical and screening delays and increased cancer mortality among patients diagnosed with colorectal, lung, and prostate cancer during the pandemic.⁴ Delayed mammography and computed tomography for lung cancer were associated with advanced stage of cancer at diagnosis.⁴ Another study determined delayed surgery for lung cancer was associated with worse survival.¹¹ For breast screenings, some evidence suggests that patients were reluctant to return for mammograms after care disruptions.¹² Thus, cancer care disruptions during COVID-19 could have detrimental future impacts on cancer outcomes and may require changes to public health and clinical strategies across the cancer prevention and control continuum.

It is unclear if patients felt comfortable returning to care in the context of rapidly changing information and guidelines related to COVID-19 and even now that guidelines are more consistent and vaccines are available. There is concern about whether patients will prioritize immediate unmet social needs that might be a result of or exacerbated by COVID-19, such as food insecurity, employment loss, and housing challenges, over disease prevention. Furthermore, people of color, including African Americans, Latinx, and Native communities, as well as those employed in low-wage occupations, are likely to have greater concerns over COVID-19 safety, in addition to the immediate concerns noted above.¹³ Rural communities that already experience limited access to cancer care, have less capacity to manage COVID-19.¹⁴ Finally, hospitals rapidly increased the use of telehealth to continue cancer care during COVID-19, but older people and those who lived in low-income and rural areas, or were less likely to have commercial insurance were less likely to participate.^{15,16} This combination of factors may exacerbate existing disparities.¹³

This survey study was conducted by NCI-designated Siteman Cancer Center to elucidate: 1) to what extent cancer care appointments (including preventive screenings and treatment) in the bi-state Midwestern catchment area were postponed or canceled, 2) patients' needs for returning to care, and 3) correlates of care disruption across the catchment area. The cancer burden is significantly greater in this catchment area than the US averages for multiple cancers. Moreover, racial and geographical disparities persist such that African American patients have higher incidence and mortality for lung, colorectal, late-stage breast cancer diagnoses, and prostate cancers compared to White patients. Rural counties also have higher mortality (but not incidence) for melanoma, breast, and prostate cancer compared to urban areas.¹⁷

Thus, we explore how socio-contextual factors impact cancer health disparities across the continuum of cancer control and prevention during COVID-19 in this bi-state Midwestern catchment area. Race, ethnicity, social class, and gender are social identities that shape many contextual factors related to cancer and COVID outcomes and are considered in this report. We stratify our results by race because of the differential impact of COVID-19 on communities of color and the over-representation of socioeconomic factors such as low-income, low-wage work often experienced by communities of color.¹⁸⁻²⁰

Methods

Data Source

Data were collected from June through August 2020 as part of Siteman Cancer Center's Community Outreach and Engagement efforts. The survey focused on understanding cancer prevention and control behaviors throughout the Siteman catchment area. The Siteman catchment area includes 82 counties throughout Missouri (40) and Illinois (42) and is diverse concerning race (21% people of color), geography (15% rural), and healthcare access (29% live in medically underrepresented areas).²¹

Data Collection

The Washington University in St. Louis, MO Institutional Review Board approved and exempted this study (ID#202006089). We recruited participants through Qualtrics® Online Panels, which emailed potential participants a survey link.²² We screened potential participants for the following eligibility criteria: age 30 or older and residing in eastern or southeastern Missouri or central or southern Illinois. Recruitment oversampled for males (35%), people of color (35%) (defined as all races & ethnicities except for non-Hispanic White), and non-metro area residents (20%) (defined as a score of 4 or greater for census-designated rural-urban continuum [RUCC] codes)²³ to allow for analyses by these groups. The median survey completion time was 20.3 minutes. All participants received an agreed-upon incentive from Qualtrics.

Measures

Outcome variable. Supplemental Table 1 provides detailed information about the measures used in this study. Our outcome of interest, care disruption, was defined as any delay in health or cancer care. Catchment area residents who reported that they decided not to attend an appointment not already canceled due to COVID-19 or they or their doctor/clinic postponed any cancer screening (Pap test, stool blood test, colonoscopy, mammogram, or PSA test) appointment were categorized as experiencing care disruption.

Explanatory variables. We included predictor variables that could result in differential access to care due to social stratification: age, race,²⁴ ethnicity, gender identity,²⁵ sex assigned at birth, sexual orientation, education,²⁴ income,²⁶ residence in non-metro area, pre-COVID employment, health insurance status, job loss due to COVID-19,²⁷ and access to a private vehicle. We also assessed self-report healthcare discrimination using a 7-item scale assessing how many times a participant experienced certain kinds of treatment (overall Cronbach's alpha = 0.92).²⁸ We also controlled for whether they scheduled a telehealth appointment.²⁹ All items were adapted from standardized measures, except for sex assigned at birth and access to a private vehicle, which were created by the study team.

We asked if residents participated in a telehealth medical appointment since the COVID-19 pandemic started and whether it was for a general medical appointment or cancer care. While this measure is not directly associated with social stratification, it could be correlated with Internet and other technology access and also predict whether someone was more likely to cancel a scheduled in-person appointment. Finally, we developed a single item to understand what patients who may have experienced care disruption would need *most* to be able to reschedule the appointment. These options included transportation, time to schedule, and knowing: how they would pay for the appointment, if the doctor's office or clinic was taking appropriate COVID-related safety precautions, if the doctor's office was still open or scheduling appointments, or that they could bring someone with them; we also included an "other" option with an open-ended response field.

Analytic Procedures

Descriptive statistics were obtained for all variables separately and by care disruption status (any care disruption compared to no disruption). Next, logistic regression modeling was conducted to determine the associations with care disruption across the catchment area. For all analyses, "prefer not to answer" responses were recoded as missing. We dropped those who reported that canceling an appointment did not apply to them (n=84) with more males, uninsured people, and those without telehealth appointments reflected in this exclusion. We also used sex at birth and not gender identity in the model due to the near-complete overlap between the two variables and the small sample size for some of the gender-diverse categories (N<6). Additionally, we recoded the job loss variable into the following 3 categories: yes, resident was laid off; no, resident was not laid off; and combined categories of don't know/not sure/prefer not to answer/not applicable. Finally, we conducted a stratified logistic regression analysis to determine if the associations of care disruption among non-Hispanic Black residents differed when compared to non-Hispanic White residents. Metro/non-metro area was excluded from the stratified non-Hispanic Black and non-Hispanic White models due to a small number of non-Hispanic Black residents in non-metro areas. We do not present other race/ethnicity in the race-stratified models due to the small sample size of participants with non-missing variables for the model in this category (N=71).

Results

Sociodemographic and care disruption descriptive information

Unadjusted sociodemographic characteristics of this sample of residents from the Siteman Cancer Center catchment area (n=680) are presented in Table 1. Residents were 46 years old on average. The diverse study sample included 41% respondents of color, and 28% of the respondents live in a non-metro area. The majority of residents identified as female (68%), lived in metro areas (73%), and had a 4-year college or graduate degree (38%). Compared to our catchment area, this sample had a higher proportion of women (68% vs. 51%) and college graduates (38% vs. 30%). We also had a higher proportion of people of color (41% vs. 21%) and residents who lived in rural areas (28% vs. 15%) due to intentional oversampling.

In this sample, approximately 55% of respondents experienced disruption to their scheduled healthcare appointments. Those who experienced care disruption were more likely to be female and have higher levels of educational attainment compared to those who did not experience care disruption.

Disruptions in cancer care during COVID-19

Table 1. Characteristics of residents across Missouri and Southern Illinois by care disruption status (July-August 2020).

Variable	Category	Total Sample (N=680) – N (%)	No Care Disruption (N=304) – N (%)	Care Disruption (N=376) – N (%)
Race	White	399 (58.7%)	186 (61.2%)	213 (56.7%)
	Black or African American	212 (31.2%)	90 (29.6%)	122 (32.5%)
	Asian/ Native Hawaiian or Other Pacific Islander	21 (3.1%)	12 (4.0%)	9 (2.4%)
	Other, including multiple groups	48 (7.1%)	16 (5.3%)	32 (8.5%)
Hispanic, Latino/a, or Spanish origin	Yes	15 (2.2%)	5 (1.6%)	10 (2.7%)
	No	664 (97.8%)	299 (98.4%)	365 (97.3%)
Gender Identity*	Woman	464 (68.2%)	192 (63.2%)	272 (72.3%)
	Man	206 (30.1%)	110 (36.2%)	96 (25.5%)
	Transgender / Gender Diverse	5 (0.7%)	1 (0.3%)	4 (1.1%)
	Prefer not to answer	5 (0.7%)	1 (0.3%)	4 (1.1%)
Sex assigned at birth*	Female	472 (69.4%)	193 (63.5%)	279 (74.2%)
	Male	204 (30.0%)	110 (36.2%)	94 (25.0%)
	Prefer not to answer	4 (0.6%)	1 (0.3%)	3 (0.8%)
Sexual Orientation	LGBTQIA+	76 (11.2%)	25 (8.2%)	51 (13.6%)
	Straight or Heterosexual	590 (86.8%)	272 (89.5%)	318 (84.6%)
	Prefer not to answer	14 (2.1%)	7 (2.3%)	7 (1.9%)
Education*	Less than High School or GED	31 (4.6%)	17 (5.6%)	14 (3.7%)
	Grade 12 or GED (High school graduate)	120 (17.7%)	64 (21.1%)	56 (14.9%)
	Some college, but did not graduate	159 (23.4%)	78 (25.7%)	81 (21.5%)
	Associates Degree or Technical School Certification	111 (16.4%)	42 (13.9%)	69 (18.4%)
	College 4 years or more (College graduate)	143 (21.1%)	63 (20.8%)	80 (21.3%)
	Graduate or professional school	115 (16.9%)	39 (12.9%)	76 (20.2%)
	Annual Household Income	\$0 to \$9,999	57 (8.4%)	32 (10.6%)
\$10,000 to \$14,999		53 (7.8%)	19 (6.3%)	34 (9.1%)
\$15,000 to \$19,999		36 (5.3%)	14 (4.6%)	22 (5.9%)
\$20,000 to \$34,999		105 (15.5%)	43 (14.2%)	62 (16.5%)

It is made available under a [CC-BY-NC 4.0 International license](https://creativecommons.org/licenses/by-nc/4.0/) .

Disruptions in cancer care during COVID-19

	\$35,000 to \$49,999	110 (16.2%)	50 (16.5%)	60 (16.0%)
	\$50,000 to \$74,999	121 (17.9%)	60 (19.8%)	61 (16.3%)
	\$75,000 to \$99,999	91 (13.4%)	45 (14.9%)	46 (12.3%)
	\$100,000 or more	105 (15.5%)	40 (13.2%)	65 (17.3%)
Metro or Non-Metro Area (RUCC codes by ZIP Code)	Metro	493 (72.5%)	222 (73.0%)	271 (72.1%)
	Non-Metro	187 (27.5%)	82 (27.0%)	105 (27.9%)
Employment (pre-COVID)	Employed Full-time	321 (47.4%)	148 (49.2%)	173 (46.0%)
	Employed Part-time	72 (10.6%)	30 (10.0%)	42 (11.2%)
	Unemployed	61 (9.0%)	29 (9.6%)	32 (8.5%)
	Homemaker	65 (9.6%)	22 (7.3%)	43 (11.4%)
	Student	4 (0.6%)	2 (0.7%)	2 (0.5%)
	Retired	84 (12.4%)	42 (14.0%)	42 (11.2%)
	Disabled	62 (9.2%)	23 (7.6%)	39 (10.4%)
	Self-Employed/Other	8 (1.2%)	5 (1.7%)	3 (0.8%)
Insurance	Private	314 (46.2%)	135 (44.4%)	179 (47.6%)
	Medicare/Medicare + Medicaid	126 (18.5%)	59 (19.4%)	67 (17.8%)
	Medicaid	120 (17.7%)	47 (15.5%)	73 (19.4%)
	Other/Unknown	22 (3.2%)	12 (4.0%)	10 (2.7%)
	Currently do not have insurance	98 (14.4%)	51 (16.8%)	47 (12.6%)
Telehealth appointment*	Yes	233 (34.3%)	85 (28.0%)	148 (39.4%)
	No	447 (65.7%)	219 (72.0%)	228 (60.6%)
Telehealth appointment type	Cancer Care	6 (2.6%)	0 (0%)	6 (4.1%)
	General Health Care	218 (94.0%)	81 (96.4%)	137 (92.6%)
	Both	8 (3.5%)	3 (3.6%)	5 (3.4%)
Access to Private Vehicle (own or others)	Yes	611 (89.9%)	275 (90.5%)	336 (89.4%)
	No	69 (10.2%)	29 (9.5%)	40 (10.6%)
Laid off Job or had to close own business*	Yes	135 (19.9%)	50 (16.5%)	85 (22.6%)
	No	423 (62.2%)	204 (67.1%)	219 (58.2%)
	Don't Know/Not Sure/Prefer Not to Answer	15 (2.2%)	2 (0.7%)	13 (3.5%)
	Not Applicable	107 (15.7%)	48 (15.8%)	59 (15.7%)
Variable		Mean (SD)		
Age		46.2 (12.6)	46.0 (13.3)	46.5 (12.0)
Discrimination ^{1*}		1.8 (0.8)	1.7 (0.8)	1.9 (0.9)

*Statistically significant difference (p<0.05; Chi-square or Fischer's test for categorical, t-test or Wilcoxon rank sum for continuous)

^Missing values: 1 Hispanic/Latina(a)/Spanish origin; 1 Education; 2 Income; 3 Employment

¹Average score of 7 items on a scale of (1) never, (2) once, (3) 2 or 3 times, and (4) 4 times or more; higher scores indicate more discrimination

Disruptions in cancer care during COVID-19

The number of residents scheduled for a cancer screening appointment, and whose appointment was postponed by the patient or their doctor/clinic is presented in Figure 1. There were 480 possible appointments scheduled between March 2020 through the end of 2020 for either a mammogram, pap test, blood stool test, colonoscopy, or PSA test. Appointment cancelations varied from 21%-57% by screening type. Additionally, in our sample, 25% of residents canceled a scheduled in-person dental appointment, 31% avoided seeking care in a hospital (e.g. labor and delivery, emergency room, etc.), and 46% of residents canceled a scheduled in-person general medical appointment (data not presented).




Figure footnotes: N shown is the number who were planning to have a screening test between March 2020 and the end of 2020; For Cancer-related care, this question was asked only of those who self-reported ever being diagnosed as having cancer and the number scheduled for the test is unknown

Figure 1. Care Disruption by Cancer Screening/Appointment Type across Missouri and Southern Illinois (July-August 2020)

Patient needs for rescheduling

In addition, we asked participants who experienced any care disruption what they would need most to reschedule their appointments (n=376). The largest proportion of participants said they would need to know if their doctor's office or clinic is taking the appropriate COVID-related safety precautions (33.8%), followed by not needing anything (18.1%). Some participants needed to know if their doctor's office is making appointments for general or routine care (13.3%) or stated they were dealing with other things and not ready to reschedule yet (10.6%). Approximately 8.2% stated they needed to have time to reschedule the appointment. All other needs were reported by less than 5% of respondents.

Correlates of care disruption

Logistic regression results for the overall and race-specific models are presented in Table 2. In the overall model, higher odds of care disruption were associated with higher educational attainment (OR=1.26, 95% CI: 1.11- 1.43), female (OR=1.60, 95% CI: 1.12- 2.30), reporting experiencing more discrimination in healthcare settings (OR=1.40, 95% CI: 1.13- 1.72), and having scheduled a telehealth appointment (OR=1.51, 95% CI: 1.07-2.15). The correlates of care disruption were not consistent across race. Among Black residents, only higher levels of educational attainment (OR=1.45, 95% CI: 1.13- 1.85) were associated with greater odds of care disruption. Whereas, among White residents, higher odds of care disruption were associated with higher levels of educational attainment (OR=1.39, 95% CI: 1.17- 1.65), female (OR=1.90, 95% CI: 1.17- 3.08), older age (OR=1.02, 95% CI: 1.001- 1.04), and having scheduled a telehealth appointment (OR=1.62, 95% CI: 1.01-2.59).

Table 2. Odds of any care disruption compared to no care disruption by social factors across Missouri and Southern Illinois (July-August 2020).

Variable	Overall Sample (N=663)		Non-Hispanic Black or African American (N=205)		Non-Hispanic White (N=387)	
	Odds Ratio	95% CI	Odds Ratio	95% CI	Odds Ratio	95% CI
Race/Ethnicity						
Non-Hispanic Black or African American	1.15	0.77, 1.72	--	--	--	--
Other Race/Ethnicity	1.40	0.79, 2.45	--	--	--	--
Non-Hispanic White (ref)	--	--	--	--	--	--
Sex Assigned at Birth ^{1,3}						
Female	1.60	1.12, 2.30	1.11	0.56, 2.19	1.90	1.17, 3.08
Male (ref)	--	--	--	--	--	--
Sexual Orientation						
LGBTQIA+	1.53	0.88, 2.65	0.68	0.27, 1.72	1.65	0.73, 3.73
Straight or Heterosexual (ref)	--	--	--	--	--	--
Area designation (by ZIP code)						
Non-Metro	1.23	0.82, 1.84	--	--	--	--
Metro (ref)	--	--	--	--	--	--
Telehealth Appointment ^{1,3}						
Yes	1.51	1.07, 2.15	1.06	0.57, 1.99	1.62	1.01, 2.59
No (ref)	--	--	--	--	--	--
Access to Private Vehicle (own or others)						
Yes	0.74	0.41, 1.33	0.79	0.33, 1.90	0.74	0.27, 1.98
No (ref)	--	--	--	--	--	--
Health Insurance						
Medicare/Medicare +	0.71	0.41, 1.24	0.88	0.31, 2.47	0.75	0.37, 1.52
Medicaid	1.02	0.59, 1.75	0.76	0.32, 1.77	1.43	0.65, 3.15
Other/Unknown	0.63	0.24, 1.66	0.75	0.15, 3.92	0.38	0.09, 1.65

Currently do not have insurance	0.66	0.38, 1.13	0.58	0.22, 1.52	0.87	0.42, 1.80
Private (ref)	--	--	--	--	--	--
Laid off Job or had to close own business						
Yes	1.55	0.994, 2.41	1.55	0.75, 3.20	1.52	0.80, 2.89
No (ref)	--	--	--	--	--	--
Don't Know/Not Sure/Prefer Not to Answer/Not Applicable	1.42	0.89, 2.25	2.12	0.87, 5.18	1.06	0.59, 1.93
Education ^{1, 2, 3}	1.26	1.11, 1.43	1.45	1.13, 1.85	1.39	1.17, 1.65
Income	0.99	0.89, 1.09	0.93	0.78, 1.13	0.99	0.87, 1.13
Discrimination ¹	1.40	1.13, 1.72	1.26	0.89, 1.78	1.29	0.96, 1.74
Age ³	1.01	0.995, 1.03	0.99	0.96, 1.02	1.02	1.001, 1.04

¹Statistically significant (p<0.05) overall variable effect – overall model

²Statistically significant (p<0.05) overall variable effect – Non-Hispanic Black or African American model

³Statistically significant (p<0.05) overall variable effect – Non-Hispanic White model

Discussion

Using primary data collected from residents across the 82-county Siteman catchment area in Missouri and Illinois, we learned that 21% to 57% of cancer screening or treatment appointments were canceled from March 2020 through the end of 2020. Across all races, residents with higher educational attainment had 1.25 higher odds of care disruption for general or cancer care compared to residents with lower educational attainment; this association remained significant among Black and White residents. Additionally, White residents of older age, assigned female at birth, or having scheduled a telehealth appointment, also had higher odds of care disruption. Finally, knowing their doctor's office or clinic is taking the appropriate COVID-related safety precautions was the greatest reported need for returning to care (33.8%).

Delays in cancer screening can lead to stage shifts where patients are diagnosed at later stages and thus have a higher risk for cancer morbidity and mortality. Understanding which screenings were impacted and for whom and identifying patient concerns can inform community outreach and engagement efforts. This allows programs to target groups most likely to have delayed screening and draft messaging that can alleviate patient concerns and in turn facilitate a return to care.

Mammograms and Pap tests are an area of increased interest for our catchment area given the high number of women scheduled for screening. Approximately 38% of the 170 women who were scheduled for mammograms had delayed or canceled appointments. Similarly, 45% of the 188 women scheduled for Pap tests had delayed or canceled appointments. Delays in colorectal cancer screening impacted a smaller number of people, but colorectal cancer screening is an important area given the high proportion of cancellations, overall low number of scheduled appointments in general, and high colorectal cancer disparities in the region. Of the 51 people scheduled for a colonoscopy, 57% delayed or canceled appointments, and of the 38 scheduled for a blood stool test, 29% delayed or canceled appointments as well. To help healthcare systems reduce the cancer screening deficit, community outreach and engagement strategies need to address these needs. For example, employing mobile strategies such as the use of mobile mammography and home-based cervical and colorectal cancer screening tests could serve those most impacted.

These data are consistent with prior literature that suggests a reduction in general medical and cancer-related appointments.^{2,5,7,30} This study allows us to understand the magnitude of the impact across Missouri and southern Illinois. Future research exploring whether those with higher educational attainment were more likely to cancel appointments because they were more likely to have better access to scheduling future appointments could further elucidate the extent of educational disparities in healthcare access.

These cross-sectional data cannot infer causality however, many of the correlates of interest (e.g., race, educational attainment) pre-date COVID-19 and the need to consider postponing clinical care. Thus, it is unlikely these results are subject to reverse causation. Also, those excluded due to missing data were more likely to be uninsured. If uninsured persons were also more likely to have postponed appointments, this could potentially bias results about care disruptions by insurance status towards the null and underestimating the impact.

Disruptions in cancer care during COVID-19

Despite these limitations, this is a significant study that can improve our understanding of COVID-19 impacts on cancer prevention and control and offer specific insights into the region. In our data, those with higher education were more likely to postpone care. This indicates that any trends seen in increasing late-stage diagnosis might occur across socioeconomic categories. Additionally, while Black and White people of higher educational attainment both had increased odds of care disruption, having a scheduled telehealth visit was significantly associated with higher odds of care disruption only for White residents. This suggests that while White people were canceling in-person care, this care may have been substituted with telehealth appointments. Many of these screenings cannot be done virtually, yet this warrants further investigation to understand if care disruption does not always equate to being disconnected from healthcare for some and the subsequent impact on racial disparities in cancer care.

Acknowledgments

The authors would like to thank the survey participants for their time, effort, and contribution to the study.

References

1. Nelson B. COVID-19 is shattering US cancer care. *BMJ*. 2020;369:m1544.
2. Patt D, Gordan L, Diaz M, et al. Impact of COVID-19 on cancer care: How the pandemic is delaying cancer diagnosis and treatment for American seniors. *JCO Clinical Cancer Informatics*. 2020(4):1059-1071.
3. Ueda M, Martins R, Hendrie PC, et al. Managing cancer care during the COVID-19 pandemic: agility and collaboration toward a common goal. *J Natl Compr Canc Netw*. 2020:1-4.
4. Zheng NS, Warner JL, Osterman TJ, et al. A retrospective approach to evaluating potential adverse outcomes associated with delay of procedures for cardiovascular and cancer-related diagnoses in the context of COVID-19. *J Biomed Inform*. 2020;113:103657.
5. Wenger NS, Stanton AL, Baxter-King R, Sepucha K, Vavreck L, Naeim A. The impact of COVID-19 on routine medical care and cancer screening. *J Gen Intern Med*. 2022;37(6):1450-1456.
6. Warner ET, Restrepo E, Benjamin C, et al. Patient-reported impact of the COVID-19 pandemic on breast cancer screening, diagnosis, and treatment: A national survey. *Clinical Cancer Research*. 2020;26(18 Supplement):S11-02.
7. London JW, Fazio-Eynullayeva E, Palchuk MB, Sankey P, McNair C. Effects of the COVID-19 pandemic on cancer-related patient encounters. *JCO Clinical Cancer Informatics*. 2020(4):657-665.
8. Riera R, Bagattini Â M, Pacheco RL, Pachito DV, Roitberg F, Ilbawi A. Delays and disruptions in cancer health care due to COVID-19 pandemic: systematic review. *JCO Glob Oncol*. 2021;7:311-323.
9. Blay JY, Boucher S, Le Vu B, et al. Delayed care for patients with newly diagnosed cancer due to COVID-19 and estimated impact on cancer mortality in France. *ESMO Open*. 2021;6(3):100134.
10. Malagón T, Yong JHE, Tope P, Miller WH, Jr., Franco EL. Predicted long-term impact of COVID-19 pandemic-related care delays on cancer mortality in Canada. *Int J Cancer*. 2022;150(8):1244-1254.
11. Mayne NR, Elser HC, Darling AJ, et al. Estimating the impact of extended delay to surgery for stage i non-small-cell lung cancer on survival. *Ann Surg*. 2021;273(5):850-857.
12. Miller MM, Meneveau MO, Rochman CM, et al. Impact of the COVID-19 pandemic on breast cancer screening volumes and patient screening behaviors. *Breast Cancer Res Treat*. 2021;189(1):237-246.
13. Cancino RS SZ, Mesa R, Tomlinson GE, Wang J. The impact of COVID-19 on cancer screening: Challenges and opportunities. *JMIR Cancer*. 2020;6:e21697.
14. Segel JE, Ross HI, Edwards JL, Braun KA, Davis LA. The unique challenges facing rural providers in the COVID-19 Pandemic. *Population Health Management*. 2021;24(3):304-306.
15. Darcourt JG, Aparicio K, Dorsey PM, et al. Analysis of the implementation of telehealth visits for care of patients with cancer in Houston during the COVID-19 pandemic. *JCO oncology practice*. 2021;17(1):e36-e43.

Disruptions in cancer care during COVID-19

16. Jaffe DH, Lee L, Huynh S, Haskell TP. Health inequalities in the use of telehealth in the United States in the lens of COVID-19. *Popul Health Manag.* 2020;23(5):368-377.
17. National Cancer Institute, Control USCFDPa. State Cancer Profiles. 2022; <https://statecancerprofiles.cancer.gov/index.html>. Accessed October 1, 2022.
18. Acosta AM, Garg S, Pham H, et al. Racial and ethnic disparities in rates of COVID-19–associated hospitalization, intensive care unit admission, and in-hospital death in the United States from March 2020 to February 2021. *JAMA Network Open.* 2021;4(10):e2130479-e2130479.
19. Athavale P, Kumar V, Clark J, Mondal S, Sur S. Differential impact of COVID-19 risk factors on ethnicities in the United States. *Frontiers in public health.* 2021;9:743003-743003.
20. Millett GA, Jones AT, Benkeser D, et al. Assessing differential impacts of COVID-19 on black communities. *Annals of epidemiology.* 2020;47:37-44.
21. United States Census Bureau. Population and Housing Unit Estimates. 2022; <https://www.census.gov/programs-surveys/popest.html>. Accessed Oct 1, 2022.
22. Qualtrics. In. Provo, Utah, USA Version June 2020.
23. United States Department of Agriculture. Rural-Urban Continuum Codes. 2019; <https://www.ers.usda.gov/data-products/rural-urban-continuum-codes.aspx>. Accessed May 14, 2020.
24. Blake K. Health Information National Trends Survey (HINTS) 5, Cycle 2 Dataset. In:2019.
25. Killermann S. How you can make the gender question on an application form more inclusive. 2020; <https://www.itspronouncedmetrosexual.com/2012/06/how-can-i-make-the-gender-question-on-an-application-form-more-inclusive/>. Accessed May 21, 2020.
26. Health Information National Trends Survey (HINTS) 5, Cycle 2. In:2018.
27. Grasso DJ, Briggs-Gowan MJ, Ford JD, Carter AS. The Epidemic – Pandemic Impacts Inventory (EPII). University of Connecticut School of Medicine. In:2020.
28. Peek ME, Nunez-Smith M, Drum M, Lewis TT. Adapting the everyday discrimination scale to medical settings: reliability and validity testing in a sample of African American patients. *Ethn Dis.* 2011;21(4):502-509.
29. Penedo FJ, Cohen L, Bower J, Antoni MH. COVID-19: Impact of the pandemic and HRQOL in cancer patients and survivors. [Unpublished questionnaire]. 2020; <https://clinicaltrials.gov/ct2/show/NCT04447222>.
30. Czeisler MÉ, Marynak K, Clarke KE, et al. Delay or avoidance of medical care because of COVID-19–related concerns—United States, June 2020. *Morbidity and mortality weekly report.* 2020;69(36):1250.
31. Centers for Disease Control and Prevention. Behavioral Risk Factor Surveillance System Survey Questionnaire. In. Atlanta, Georgia: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention; 2018.
32. Wadhera RK, Choi E, Shen C, Yeh RW, Joynt Maddox KE. Trends, causes, and outcomes of hospitalizations for homeless individuals: A retrospective cohort study. *Med Care.* 2019;57(1):21-27.
33. Wadhera RK, Khatana SAM, Choi E, et al. Disparities in care and mortality among homeless adults hospitalized for cardiovascular conditions. *JAMA Intern Med.* 2020;180(3):357-366.

Disruptions in cancer care during COVID-19

34. Qi AC, Peacock K, Luke AA, Barker A, Olsen MA, Joynt Maddox KE. Associations between social risk factors and surgical site infections after colectomy and abdominal hysterectomy. *JAMA Netw Open*. 2019;2(10):e1912339.

Disruptions in cancer care during COVID-19

Appendix
Supplemental Table 1: Survey item information

Construct	Survey item and response options	Variable coding and interpretation	Measure source
Scheduled care and care disruptions			
Care delay (primary outcome)	Due to COVID-19, did you decide to: Cancel a scheduled in-person general medical appointment not already cancelled due to COVID-19; Not included, but related items: Cancel a scheduled in-person dental appointment not already cancelled due to COVID-19; Avoid seeking care in a hospital (e.g. labor and delivery, emergency room, etc.) <i>Response options: a) yes, b) no, c) does not apply</i>	Yes for general medical appointment = care disruption <i>(Dental and hospital care not included in care disruption primary outcome)</i>	Adapted from Penedo et al., 2020 ²⁹
Care delay (primary outcome)	Did you or your doctor postpone your mammography because of COVID-19? <i>Response options: a) yes (it was me), b) yes (it was my doctor or clinic), c) no, d) don't know/not sure</i>	Yes (a or b) = care disruption	Penedo et al., 2020 ²⁹
Care delay (primary outcome)	Did you or your doctor postpone your Pap test because of COVID-19? <i>Response options: a) yes (it was me), b) yes (it was my doctor or clinic), c) no, d) don't know/not sure</i>	Yes (a or b) = care disruption	Penedo et al., 2020 ²⁹
Care delay (primary outcome)	Did you or your doctor postpone your stool blood test because of COVID-19? <i>Response options: a) yes (it was me), b) yes (it was my doctor or clinic), c) no, d) don't know/not sure</i>	Yes (a or b) = care disruption	Penedo et al., 2020 ²⁹
Care delay (primary outcome)	Did you or your doctor postpone your colonoscopy because of COVID-19? <i>Response options: a) yes (it was me), b) yes (it was my doctor or clinic), c) no, d) don't know/not sure</i>	Yes (a or b) = care disruption	Penedo et al., 2020 ²⁹
Care delay (primary outcome)	Did you or your doctor postpone your PSA test because of COVID-19? <i>Response options: a) yes (it was me), b) yes (it was my doctor or clinic), c) no, d) don't know/not sure</i>	Yes (a or b) = care disruption	Penedo et al., 2020 ²⁹
Cancer Care delay	Were you scheduled for any cancer-related medical care that you or your doctor had to cancel or postpone during the COVID-19 restrictions? <i>Response options: a) yes, I had to cancel or postpone, b) yes, my doctor had to cancel or postpone, c) no, d) don't know/not sure</i> <i>*Not included in current manuscript</i>		Penedo et al., 2020 ²⁹
Telehealth appointment	Did you participate in a Telehealth medical appointment (e.g., Zoom, Facetime) since the COVID-19 pandemic?	Yes = participated in telehealth appointment	Adapted from Penedo et al., 2020 ²⁹

Disruptions in cancer care during COVID-19

	<i>Response options: a) yes, b) no</i>		
Telehealth appointment	(if yes to previous item): Did you participate in a Telehealth appointment for: <i>Response options: a) cancer care, b) general health care, c) both</i>	Response options correlate to telehealth care type	Team created
Appointment rescheduling needs	If you were unable to keep a doctor's appointment for general health care or cancer screening during the COVID-19 pandemic, what do you need <u>most</u> to be able to reschedule that appointment? I need: <i>Response options (select one): a) To know my doctor's office or clinic is taking the appropriate COVID related safety precautions, b) Time to schedule the appointment, c) To know if my doctor's office is still open, d) To know my doctor's office is making appointments for general or routine care, e) To know how I'm going to get to the doctor's office (transportation), f) To know how I'm going to pay for the doctor's appointment, g) To know that someone can come with me, h) I don't need anything, i) I am dealing with other things and not ready to reschedule yet, j) Other (please specify)</i>	Selection of any response a-g, j indicates a need for rescheduling; h or i response considered no need	Adapted from Penedo et al., 2020 ²⁹
Demographic and social stratification variables			
Age	What is your age?	Age measured as a continuous variable in years	Team created
Race	Which of these groups would you say best represents your racial identity? (Please mark all that apply) <i>Response options: a) American Indian or Alaska Native, b) White, c) Black or African American, d) Asian, e) Middle East and North African (MENA), f) Native Hawaiian or Other Pacific Islander, g) Another group (please specify)</i>	Combined race and ethnicity recoded into 3 categories: Non-Hispanic Black/African American, Non-Hispanic White, any other race/ethnicity	Adapted from BRFSS 2018 ³¹
Ethnicity	Are you Hispanic, Latino/a, or Spanish origin? <i>Response options: a) yes, b) no</i>		Adapted from HINTS 2018 ²⁴
Gender	What is your gender identity? <i>Response options: a) Man, b) Woman, c) Transgender, d) Queer/Non-binary, e) Agender/No gender, f) Something else (please specify), g) Prefer not to answer</i>	Recoded into 3 categories: a = man, b = woman, c-f = transgender/gender diverse	Adapted from Killermann, 2020 ²⁵
Sex assigned at birth	What is your sex assigned at birth, on your birth certificate? We ask this question to better understand what health screenings are relevant for you. <i>Response options: a) Female, b) Male, c) Prefer not to answer</i>	Sex assigned at birth correlate to response options	Team created

Disruptions in cancer care during COVID-19

Sexual orientation	What is your sexual orientation? <i>Response options: a) Asexual, b) Bisexual, c) Gay, d) Lesbian, e) Pansexual, f) Straight or heterosexual, g) Something else (please specify), h) Prefer not to answer</i>	Recoded into 3 categories: a-e, g = LGBTQIA+, f = straight or heterosexual, h = prefer not to answer	Adapted from Penedo et al., 2020 ²⁹
Education	What is the highest grade or year of school you completed? <i>Response options: a) Never attended school or only attended kindergarten, b) Grades 1 through 8 (Elementary), c) Grades 9 through 11 (Some high school), d) Grade 12 or GED (High school graduate), e) Some college, but did not graduate, f) Associate degree or Technical School Certification, g) College 4 years or more (College graduate), h) Graduate or professional school, i) Other (please specify)</i>	Recoded into 7 categories: a-c = less than high school, d = high school graduate or GED, e = some college, f = associate degree or technical school certificate, g = college graduate, h = graduate or professional school	Adapted from BRFSS 2018 ³¹
Income	Thinking about members of your family living in this household, what is your combined annual income, meaning the total pre-tax income from all sources earned in the past year? <i>Response options: a) \$0 to \$9,999, b) \$10,000 to \$14,999, c) \$15,000 to \$19,999, d) \$20,000 to \$34,999, e) \$35,000 to \$49,999, f) \$50,000 to \$74,999, g) \$75,000 to \$99,999, h) \$100,000 to \$199,999, i) \$200,000 or more</i>	Recoded into 9 categories: less than \$10,000, \$10,000-\$14,999, \$15,000-\$19,999, \$20,000-\$34,999, \$35,000-\$49,999, \$50,000-\$74,999, \$75,000-\$99,999, and \$100,000+.	HINTS 2018 ²⁴
Rurality	What is the zip code where you live? <i>Response option: numerical entry</i>	Zip codes with RUCC code designation of 4 or greater classified as non-metro	Rural-Urban Continuum Codes, 2019 ²³
Pre-COVID employment	Which category best describes your occupational status in February 2020 prior to the stay-at-home orders put in place as a result of the COVID-19 pandemic? <i>Response option: a) Employed Full-time, b) Employed Part-time, c) Unemployed, d) Homemaker, e) Student, f) Retired, g) Disabled, h) Other (please specify)</i>	Response options correlate to employment status	Adapted from Penedo et al., 2020 ²⁹
Individual job loss due to COVID-19	Have you or anyone in your household experienced the following: Laid off job or had to close own business <i>Response options: a) Yes (me), b) Yes (person in home), c) No, d) Don't know/Not sure, e) Prefer not to answer, f) Not applicable</i>	Yes (a) = job loss	Adapted from Grasso et al., 2020 ²⁷

Disruptions in cancer care during COVID-19

Health insurance status	<p>Are you currently covered by any of the following types of health insurance or health coverage plan?</p> <p><i>Response options: a) Indian Health Service or Tribal Health Services, b) A plan purchased through a current or former employer or union, c) A plan that you or another family member buys on your own from an insurance company, including Marketplace plans, d) Medicare, for people 65 and older, or people with certain disabilities, e) Medicaid or any kind of government-assistance plan for those with low incomes or a disability, f) TRICARE, VA, or other Military, g) Any other type of health insurance or health coverage plan (please specify), h) I currently do not have health insurance</i></p>	<p>Recoded into 5 categories: b or c = private insurance, d = Medicare, e = Medicaid, h = uninsured, a,f,g = other/unknown</p>	<p>BRFSS 2018³¹ coding based on Wadhera et al., (2019, 2020) & Qi et al., 2019³²⁻³⁴</p>
Access to a private vehicle	<p>What kind of transportation do you most often use to get to the places you need to go?</p> <p><i>Response options: a) Either my own or someone else's private car, van, truck or motorcycle, b) Bus, c) Light rail like the Metrolink, d) Call-a-ride, e) Taxi/Uber/Lyft, f) Other (please specify)</i></p>	<p>a = access to private vehicle</p>	<p>Adapted from BRFSS 2018³¹</p>
Healthcare discrimination	<p>7-item scale; hen getting health care, how often has each experience happened to you: a) Treated with less courtesy than other people, b) Treated with less respect than other people, c) Received poorer services than other people, d) Had a doctor or nurse act as if he or she thinks you were not smart, e) Had a doctor or nurse act as if he or she was afraid of you, f) Had a doctor or nurse act as if he or she was better than you, g) Felt like a doctor or nurse was not listening to what you were saying</p> <p><i>Response options: 1) never, 2) once, 3) 2 or 3 times, 4) 4 or more times</i></p>	<p>Mean scores calculated across items; higher score indicates more discrimination</p>	<p>Peek et al., 2011²⁸</p>

Note: Affirmative responses to any one of the seven care delay items indicated the presence of care disruption.