


Bariatric surgery: preparations and quality of life consequences

Liis Lozano^{1,2}, Triin Põder^{1,3}, George A. Lozano^{4*} 

¹ Tartu Health Care College, Nooruse 5, Tartu, Estonia, 50411

² Current address: Department of Hematology and Oncology, Tartu Clinic Hospital, Puusepa 8, Tartu, Estonia, 51014. Email liislozano@kliinikum.ee

³ Current address: Raatuse Health Centre, Raatuse 21, Tartu, Estonia, 50603. Email triin.poder7@gmail.com

⁴ Estonian Centre of Evolutionary Ecology, 15-8 Tähe Street, Tartu, Estonia, 50108. Tel +37255633099. dr.george.lozano@gmail.com

Abstract

Introduction.- Obesity is a major threat to global health. When more conventional methods have failed, obesity can be addressed via bariatric surgery. Here we examine the reasons why patients choose bariatric surgery and the behavioral consequences thereafter.

Methods.- A qualitative study with a phenomenological design was used to analyze detailed interview responses from recipients of bariatric surgery.

Results.- Before surgery, (a) bariatric surgery was chosen for obvious reasons: patients had been overweight, had obesity-related health problems, and had difficulties moving. (b) Information was obtained from medical practitioners and online discussion groups. (c) Fear, anxiety, and apprehension were common. Friends and family were sometimes supportive and sometimes disapproving. After surgery, (1) many chronic health problems disappeared or eased significantly. (2) Subjects had difficulties adjusting to small portions and altered food preferences. (3) Physical activity increased. (4) Self-esteem increased but the worry of regaining the weight remained. (5) As before surgery, there were both supportive and condemning attitudes by relatives, friends, and society. (6) A desire for further nutritional and psychological counseling was indicated. (7) Relationships and quality of sex improved in most cases but not always. Single women, particularly, became more active in potential relationships.

Discussion.-With a few exceptions, our results agree with the literature, supporting the idea that bariatric surgery leads to extensive physical, psychological, and social changes. Hence, patients ought to be better prepared for these changes, and medical practitioners ought to be aware of the magnitude of the changes this surgery will bring about in their patients' lives.

Keywords: Bariatric surgery, Quality of life, Pre-op counseling, Post-op support.

INTRODUCTION

Over the past fifty years, obesity has become a major global health concern^{1,2}. Obesity decreases life expectancy and quality of life by increasing the risk and severity of many diseases, among them type-2 diabetes³, cardiovascular disease⁴, asthma⁵, covid-19⁶, sleep apnoea⁷, periodontal disease⁸, hypertension⁹, osteoarthritis, fatty liver disease, stroke, dementia, and several cancers^{1,10}. Although obesity as an epidemiological concern began in the world's most affluent counties^{2,11}, as the global standard of living has risen, obesity has also become a problem in developing countries^{12,13}. As of 2020, over 1.9 billion people in the world were overweight, of which over 650 million were obese¹⁴.

Many approaches have been used to combat obesity^{15,16}. Epidemiologically, programs that increase the activity level¹⁷ or alter people's diet¹⁸ have been implemented. In addition, several evolutionary explanations for obesity have been proposed, each one with its own prognostic solutions¹⁹⁻²¹. At the individual level, obesity can be addressed by several means, among them diet modification^{22,23}, increased activity^{24,25}, cognitive therapy^{26,27}, and appetite suppressive drugs^{28,29}. When other options have failed, bariatric surgery becomes a viable alternative³⁰.

Bariatric surgery is a laparoscopic procedure whereby the volume of the stomach and/or the length of the small intestine are reduced^{31,32}. There are several types of surgeries, affecting the size of the stomach, the path food takes, and the amount and section of intestine removed. Bariatric surgery has two general effects: first, the amount of food that can be consumed at a given feeding bout is reduced, and second, the absorption of this food also decreases^{33,34}. Bariatric surgery is one of the few effective long-term cures for severe obesity, and it also relieves many associated conditions, leading to an increase of both lifespan and quality of life^{30,35}.

Lozano *et al.* **Bariatric surgery: preparations & quality of life**

As with any surgery, bariatric surgery patients receive pre-op information and post-op care instructions. However, unlike some other surgical procedures, bariatric surgery leads not only physical changes, but also social and psychological changes³⁶. Information necessary to help patients deal with these major changes in their lives is relatively recent and still limited³⁷. In Estonia, there has been work on the preparation necessary for surgery and post-surgical complications³⁸ and on post-surgical nursing-care recommendations³⁹. However, no research exists on the social and psychological effects. Hence, the aim of our research is to examine from a psychological, social, and medical perspective: (1) the preparations for bariatric surgery, and (2) the consequences after surgery.

METHODS

Background information

The research was conducted in Estonia, where as of 2016, 35.2% of the population was overweight (BMI > 25) and 19.2% obese (BMI >30)⁴⁰. The costs of bariatric surgery in Estonia are covered by the government for people with a BMI of over 40 who have not been able to lose weight by other means, and for people with a BMI of over 35 who have at least one obesity-related disease⁴¹.

From 2004 to 2017, 5752 bariatric surgeries were performed in Estonia. In the first couple of years, fewer than 10 surgeries per year were conducted but the frequency has been increasing, and since 2013, the number of surgeries has been stable at almost 700 surgeries per year. In 2017, there were 560 surgeries per 1 million people, a rate that is similar to that of Norway (566) and Sweden (565), and higher than the rate in Finland (166) and in Denmark (125)⁴².

Lozano *et al.* **Bariatric surgery: preparations & quality of life**

Study subjects

Participants were recruited using an Estonian Facebook social media group called “Patients of bariatric surgery”, created on July 2014 and as of January 21, 2022, had 974 members. An invitation was sent to the group. Interested people were told about the aims of the research, and those who agreed to participate were asked to read and sign an informed consent form. Participants were adults (≥ 18 years old) who had undergone bariatric surgery at least a year before.

Interviews

This research is qualitative and phenomenological. Qualitative methods of research address everything that is connected with the issue at hand: people's experiences, feelings, behavior, expectations, and beliefs. Phenomenological research methods examine a person's experience, usually by means of an interview⁴³.

Participants were asked three questions related to their experiences before surgery (Tab. 1) and seven related to their experiences after surgery (Tab. 2). The pre-surgery questions addressed (1) the reasons for choosing surgery, (2) information sources, and (3) personal emotions and attitudes. After surgery, they were asked about (1) health, (2) nutrition, (3) physical activity, (4) self- image, (5) people’s attitudes, (6) psychological issues, and (7) relationships. Within each sub-category, participants were allowed to speak freely about their experiences. Participants emphasized different issues, but there were some commonalities. The questions were further explained as required, and the replies were followed up by clarifying inquiries.

Lozano *et al.* **Bariatric surgery: preparations & quality of life**

Interviews were audio-recorded and subsequently transcribed using a speech recognition program⁴⁴. The transcriptions were corrected as required, using the recordings for confirmation.

Ethical Considerations

Participants were informed about the goal of the research, the fact that it is voluntary to participate, and that they have the right to refuse to answer any questions, and to remove themselves from the study at any stage. All participants agreed to continue until the end. Participants read, filled, and signed an informed consent form. The data were collected anonymously. The research was approved by the Tartu University human research ethics committee (protocol number 294/T-26).

RESULTS

Of the 12 interviews, nine were in person and three online via Skype; 10 participants were female and 2 were male. The interviews were conducted from August to October, 2019. The shortest one lasted 11 min. and the longest one lasted 32 min.

Choosing and preparing for surgery

There were 3 general questions: (1) reasons for choosing bariatric surgery, (2) getting information about the procedure, and (3) personal emotions and attitudes of other people before the surgery. Nineteen general topics were identified (Tab. 1).

Lozano *et al.* **Bariatric surgery: preparations & quality of life**

Table 1. Changes and experiences before surgery. Blank = not mentioned, Y = yes, N = no, - = Yes, but not enough. See the text for more detailed explanations of the categories.

	1	2	3	4	5	6	7	8	9	10	11	12
1.- Reasons for choosing surgery												
Excess weight	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Health reasons	Y	Y	Y	Y	Y	Y	Y		Y	N	Y	Y
Difficulties moving	Y	Y	Y		Y		Y		Y	Y		Y
Doctor's advice	Y	Y		Y	Y	YN		Y			Y	Y
Suggestion from friends/family	Y		Y	Y	Y			Y		Y		
Desire to have children											Y	
Desire to play/be with children					Y		Y				Y	Y
Wish to be more attractive	Y		Y		Y		Y					
Concerns with family obesity	Y		Y	Y								
Felt negativity from people	Y-		Y	Y	Y			Y			N	Y
Overt insults from people					Y			Y				Y
2.- Information about the surgery												
From health care workers	N	Y	Y	Y	N	Y-	Y	Y	Y	N	Y	N
From the internet		Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	N
From previous surgery patients	Y	Y		Y				Y		Y		N
3.- Emotions and people's attitudes												
Anxiety	Y			Y	N			Y				
Fear	N			N	Y			Y		N		Y
Uncertainty about the future		Y		Y	Y			Y	Y			Y
Supporting attitudes	Y-	Y							Y	Y		N
Condemning attitudes			Y	Y						Y		Y

Lozano *et al.* **Bariatric surgery: preparations & quality of life**

Reasons for choosing bariatric surgery

In all cases, attempts to lose weight via more conventional means, diet and exercise, had not been successful. The two main reasons why people chose to undergo bariatric surgery were because they were “overweight” (technically “obese”), and because of the associated health problems. The main health problems mentioned were high blood pressure, type-2 diabetes, thyroid malfunction, and joint pain. The latter likely contributed to the third main reason: difficulties moving. The one exception not affected by those reasons was more pragmatic and decided to undergo bariatric surgery because of a doctor’s advice and suggestions from family members. In some cases, the surgery had been recommended by a friend or family member who had already undergone bariatric surgery.

Other reasons included the desire to be more sexually attractive, to play or interact more with children or grandchildren, and in one case, to have children in the future. The category “family obesity” refers to the concern expressed by three participants of gaining even more weight in the future because their family members had excess weight that had increased with age. Some participants mentioned that they **felt** negativity from people, and others indicated that they had **actually experienced** overt negativity and insults. One participant had sad memories from high school when she was teased and humiliated.

Obtaining information about bariatric surgery

Participants obtained information from doctors, the internet, and other patients. Most participants said that they were sufficiently informed by their doctor, but others disagreed, or stated that others, but not themselves, would benefit from additional information. Other than that, there was a lot of information in the internet forums and particularly from the Facebook group we used to recruit the participants. Some participants indicated that they would have

Lozano *et al.* **Bariatric surgery: preparations & quality of life**

liked to get more specific information from doctors instead of having to look it up on the internet.

Personal emotions and other people's attitudes

Before the surgery, participants felt anxiety and fear. Patients were mostly afraid of dying on the surgery table, complications from the surgery, and dealing with life after the surgery. Participants were also worried about how others would react to their decision to go through the surgery. About the decision to go to surgery, there was support from relatives and friends, but there were also people who were doubtful and hesitant.

Some participants initially had a negative attitude towards surgery because they thought that surgery was the easy route, instead of just eating less and exercising more. People also thought that there might be complications during the surgery and their health might even get worse. Finally, they feared that they would not be able to change their nutrition anyway and would end up gaining back the weight.

Life after surgery

There were seven general topics: (1) health, (2) nutrition, (3) physical activity, (4) self-image and body perception, (5) people's attitudes, (6) psychological issues, and (7) relationships. Twenty-nine general themes were identified (Tab. 2).

Lozano *et al.* **Bariatric surgery: preparations & quality of life**

Table 2. Changes and experiences after surgery. Blank = not mentioned, 0 = no change, Y = changes either positive or negative, + = positive change, - = negative change. A = agree. See the text for further explanations about the categories.

	1	2	3	4	5	6	7	8	9	10	11	12
1 Health												
General wellbeing	+	+ -	+	+	+	+ -	+	+	+	+	+	+
Weight loss	+	+	+	+	+	+	+	+	+	+	+	+
Blood pressure			+	+	+	+		+	+		+	+
Blood sugar		-		+	+	+						
Joint pain	+					0	0					
Excess skin	-		-	0		-	-	-	-		-	-
Hair loss				-	-		-	-	-	-		-
Nails worsened												-
Heartburn		-							-			-
The lack of vitamins	-			-	-		-	-				-
The lack of iron								-			-	-
Other health changes		+ -		+	+	+	+	+ -	+ -	+	+	+
2 Nutrition												
Changes in food taste		Y	Y	Y	Y	Y	Y	Y	Y	Y		Y
Food choices		Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Food consumption	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Need for counseling	A	A		A	A	A				A	A	A
Eating problems	A			A	A							A

Lozano *et al.* **Bariatric surgery: preparations & quality of life**

Table 2.. cont.

3 Physical activity

Energy and strength + 0 + + + + + + + + +

Amount of movement + + 0 + + + 0 + + + +

4 Self-image

Self-image + 0 0 + + 0 + + + +

Body perception + 0 - + + + + + + +

Clothing changes + + + + + + +

5 People's attitudes

Family and friends + + + 0 - + + + +- + -

Others + + - + + + + +- + -

6 Psychological issues

Self-confidence + 0 + + + + + + + 0

Fears and doubts Y Y Y Y Y Y Y Y

Other issues Y Y Y Y Y Y Y Y

7 Sex and relations

In couple relations 0 0 + + +- + +- - + 0 -

In sexuality 0 0 + - 0 + + 0 +

Lozano *et al.* **Bariatric surgery: preparations & quality of life**

Health

This category included general well-being, weight loss, blood pressure, blood sugar, joint pain, excess skin, hair loss, nail quality, heartburn, lack of vitamins, lack of iron, and other health changes.

Eight people had the surgery without complications and four people had complications. One person had a longer time of surgery and a larger wound because of liver problems. A second person had a leak of the stomach and peritonitis and needed to stay longer in the hospital. The third person had a fever and infection that was cured after a few days in hospital. Finally, a woman found out after surgery that she was pregnant but fortunately, she later gave birth to a healthy baby.

Everyone lost weight. Nearly everyone reported an increase in wellbeing. Nine participants reported that their previous hypertension was alleviated and about half reported blood sugar levels had improved.

Fast weight loss left excess skin but most of the people accepted it as inevitable and dealt with it with suitable underwear. A couple of people removed the excess skin via plastic surgery and a few had planned to do so in the future.

Several participants experienced hair loss 4-6 months after the surgery but it grew back, sometimes even thicker and healthier. Joint pain was mentioned three times. In one case, it decreased. In two cases, it did not change.

There were also several negative consequences. Some people mentioned heartburn and the need for anti-acid medication. Also, there were people who had swings of blood sugar and dumping syndrome. There were also problems with digestion, dryness of skin, vitamin deficiencies (particularly group B vitamins), iron-deficiency, and anemia. A couple of women

Lozano *et al.* **Bariatric surgery: preparations & quality of life**

mentioned changes in their menstrual cycles; in one case, the cycles became more regular, and in the other, the menstruation became longer and more profuse (menorrhagia). Many people mentioned feeling cold and needing warmer clothing.

However, despite some negative changes in health, most people said that they got rid of numerous health concerns or the health problems eased significantly.

Nutrition

Participants mostly mentioned a decrease in the amount of food they consumed, along with changes in their taste and food preferences. About half of the people searched for information independently online, changed their diets, tested their suitability, and established new balanced diets. The other half had difficulties from the start about how much and what to eat. At least three people kept eating the same amounts and vomiting afterwards. There were also problems with digestion and dumping syndrome. Many formerly preferred foods lost their appeal, and some foods that were previously unpleasant became enjoyable. Some interviewees mentioned that eating was not pleasurable anymore and they only ate because of the need to eat. People indicated there was insufficient information from health care workers and the materials and instructions they had received.

One person developed an eating disorder. The participant thought about food constantly and then decreased the amount of food consumed to the point giving up food completely and losing too much weight. Eventually, this person sought psychiatric help and recovered.

Self- image and body perception

We differentiated between two closely linked concepts: “self-image” (who you are) and “body perception” (how you view your body). There was an increase in self-satisfaction: people were happier, became more social, and felt more attractive. Both because of the weight

Lozano *et al.* **Bariatric surgery: preparations & quality of life**

reduction and the psychological effects, several participants indicated that they had changed their wardrobe. The size of clothing changed, but also the style. Women started wearing more colorful and revealing clothing. If in the past, people had difficulties to find clothing that fit but now they had more options. Women often changed their hair color.

People's attitudes to the surgery

The attitudes of close people were both positive and negative. Most participants' relatives were positive and supportive. In contrast, several people said that their family members were strongly against the surgery and did not support them in their choice. A couple of people actually hid their decision to have the surgery and revealed it only after the procedure had been completed. The attitude of friends, colleagues and acquaintances was generally positive but there were also negative opinions.

Psychological Issues

Most people experienced an increase in self-confidence. They felt more acceptable in society, and more attractive to opposite sex. Women felt more feminine. Both men stated that they were self-confident before the surgery and in that respect did not feel any different. The increase in self-confidence led to greater courage communicating with people, and a desire for new activities and directions in life, for example, to go back to school.

On the other hand, there were also negative effects. The biggest concern was that the weight loss would not be permanent. In some cases, the degree of self-criticism increased and people were still unhappy with their own image. One person developed body dysmorphia, followed by self-injury/cutting, and eventually recovered with psychiatric help. Some people had problems with depression and insomnia. In general, people stated the need to get psychological counseling before and after surgery. They indicated the need for emotional support from professionals when they did not get it from their partner or family.

Lozano *et al.* **Bariatric surgery: preparations & quality of life**

Relationships and sex

For most people who already were in relationships, their sexual relationship improved, but in some cases, earlier problems were exacerbated. Oddly, one reason mentioned for a break-up was that the partner did not like the new slimmer body and he/she preferred the previous lush figure. People found new partners, married and had children. Some participants, including both males, did not report any changes in their relationships or sexuality. Single people, including most single women, found more self-confidence, started communicating and socializing more, and had more sexual relationships.

DISCUSSION

Obesity is a global epidemic and a major health concern and bariatric surgery is an increasingly common treatment option. Here, we examined people's experiences before and after bariatric surgery. Experiences **before** surgery were divided into: (1) reasons for choosing surgery, (2) getting surgery information, (3) personal emotions and attitudes. Experiences **after** surgery were grouped into: (1) health, (2) nutrition, (3) physical activity, (4) self- image, (5) people's attitudes, (6) psychological issues, and (7) relationships.

Experiences before surgery

The reasons for choosing bariatric surgery were unsurprising: people were overweight, had related health problems, had difficulties moving, and hoped for positive changes, such as having a family and looking better. Other researchers report similar findings⁴⁵⁻⁴⁷, which can

Lozano *et al.* **Bariatric surgery: preparations & quality of life**

be summarized as primarily medical reasons first and secondarily, psychological and social reasons.

A Brazilian study of 12 women stands apart in that the motivation for bariatric surgery was mostly social: the desire to look better, to be more accepted in society, and to enter the labour market⁴⁸. Cohn *et al.*⁴⁹ also point out that people choosing bariatric surgery desired to improve their professional lives. Participants in our study did mention negative attitudes towards overweight people, but did not mention any such experiences specifically at their place of work. Overt discrimination and negative attitudes towards obese people at the workplace might not be as prevalent in Estonia.

Information about bariatric surgery came from medical professionals and the internet. Participants used official medical sites on the internet, but they also indicated that they valued social media groups where people share common experiences. Cohn *et al.*⁴⁹ offered the same conclusion. In our study, most people thought that they had received sufficient information from their physicians, but others disagreed. Whereas attending information meetings in person can also contribute to the decision to undergo bariatric surgery⁴⁹, this factor was not mentioned by the participants of our study, who instead preferred to communicate online.

Personal emotions and people's attitudes before surgery were complex. Family members' attitudes were bipolar. Most participants' relatives were supportive but several were against the surgery. Similarly, friends', colleagues', and acquaintances' attitudes were generally positive but there were also negative opinions. There was uncertainty, anxiety, and apprehension about the surgery itself, personal changes afterwards, and reactions from other people. Similar psychological and sociological concerns have been reported recently in the literature^{5,49}. Hence, the decision to undergo bariatric surgery, although primarily motivated by health concerns, is also strongly affected by psychological and social factors.

Experiences after surgery

Health improved as expected. In the immediate aftermath, four out of 12 people had post-operative complications, but not after they had left the hospital. Physicians dealing with emergencies of bariatric surgery patients need to know the type of surgery performed and the common emergencies arising from bariatric surgeries⁵⁰.

Everyone reported weight loss, so the main goal of the surgery was achieved, and several related health problems also decreased. This is not unexpected⁵¹ and it explains the increasing popularity of bariatric surgeries. However, weight loss and the associated health benefits are highly variable⁵². Skin and hair problems are to be expected due to micronutrient and mineral deficiencies⁵³, but in due time, people in our study managed to alter their diets as required. Other studies have reported a decrease in joint pain, particularly in the knee⁵¹. In our study, joint pain decreased in one case, but did not change in two cases. An explanation might be that due to the previous weight, leg joints were already damaged, perhaps permanently.

Several people mentioned feeling cold, which has not been reported elsewhere, but is not particularly surprising, as a layer of fat is a common thermal insulator for mammals living in cold environments⁵⁴. This discomfort must be weighed against the benefit of becoming less susceptible to heat stress, which might be important in warmer climates but perhaps not so much in Estonia, other than in the sauna. Several other minor health concerns were mentioned, but the net effect was that bariatric surgery had a positive effect, not only weight loss, but also improving related health problems.

Nutritional changes dealt mostly with the amount of food consumed, and changes in food preferences and flavors. Similar findings have been reported elsewhere⁵⁵. In our study, about half the people found information on their own and changed their diet as necessary. The other

Lozano *et al.* **Bariatric surgery: preparations & quality of life**

half had difficulties from the start and would have benefited from receiving more information and support from medical personnel⁵⁵.

Another important but not unexpected effect was a deficiency in some micronutrients: iron, zinc, and vitamin B12. Micronutrient and mineral deficiencies can be manifested as readily visible skin problems⁵³. Solving this issue is usually a matter of being aware of the possibility and taking the right nutritional supplements⁵⁶. Nutrition support from professionals might be particularly important for some patients⁵⁷, about half in our sample.

After surgery, one participant in our study developed what might be considered the opposite disorder. From its description, it approached anorexia nervosa, even if was not formally diagnosed as such. Oddly, obesity is officially classified as a nutritional disorder, but anorexia nervosa as an eating disorder⁵⁸. However, both obesity and anorexia nervosa share many biological, psychological, and sociological components and might actually be two extreme expressions of the same predispositions⁵⁹. At the population level, the two disorders might even be causally linked²¹.

Physical activity increased substantially: moving became easier and more pleasant, the distances became longer, and people had more stamina and strength. This has been reported elsewhere⁶⁰, with the caveat that some improvements decreased after one year⁶¹. Other studies have reported a decrease in joint pain, particularly knee pain⁵¹. In our study, one person mentioned a decrease in joint pain decreased in one case, but two people indicated that it had not changed. An explanation might be that due to the previous weight, leg joints were already damaged, perhaps permanently.

Self-Image and body perception did not necessarily change, but people generally felt happier, more social, and more attractive. Others have reported cases whereby patients struggled with their self-identity³⁶, but in our study, it was not an issue. About half of men are satisfied with the results of their surgeries, compared to only 12% of women⁶². Women in our

Lozano *et al.* **Bariatric surgery: preparations & quality of life**

study reported more changes in clothing style than men did, so changes in women, unlike those in men, had more external manifestations and were more readily visible to society.

People's attitudes after the surgery continued being both positive and negative, just like before the surgery, whether from family, friends, colleagues, or acquaintances. Benson Davies *et al.*⁶³ concluded that family support is important to maintain the weight loss up to 6 years after surgery. Perhaps more important than the attitudes towards the surgery itself and the weight loss, attitudes towards the ensuing behavioral and lifestyle changes might be more contentious. Bariatric surgery leads to changes in self-worth, physical activity, psychological state, and life directions, changes that are bound to lead to new and different social interactions.

Psychological issues were generally positive, with improvements in self-esteem leading to a more positive attitude in life. However, just because the changes were generally positive does not mean they were easy to manage. Furthermore, there were some problems with depression, addiction, self-injury, and insomnia. Our study's participants indicated the need for emotional guidance from professionals, particularly when family members were unsupportive. The potential benefits of psychological counseling, both before surgery to prepare patients for the forthcoming changes, and after to help them navigate these changes and identify problems before they develop, have been previously noted⁵⁵. Detailed cost-benefit analyses need to be conducted to determine the degree to which more extensive psychological counseling can become an integral part of the procedure.

Sex and relationships generally improved but there were some cases whereby problems couples already had, actually intensified. Some people became more sexually active and others did not report any differences. Single people became more sexually engaged and sexually active. Despite a better sex life, some marriages deteriorated.

Lozano *et al.* **Bariatric surgery: preparations & quality of life**

We did not have enough male participants to note differences between the sexes. In couples, “sexual function” of bariatric surgery patients and their partners generally improves⁶⁴. However, among single people, “sexual satisfaction” increases in women, but not in men⁶⁵. This is despite that erectile function and testosterone do increase in men after losing weight⁶⁶. This apparent incongruity occurs because “sexual function” is not the same as “sexual satisfaction”. For males, bariatric surgery might improve sexual function, placing patients at par with every other male and still facing the main hurdle: access to women. In contrast, for females, access of males is seldom a problem. Female sexuality is not about the actual act, but rather about receiving attention from men and then choosing to act (or not) as they see fit. It is reasonable to expect that the sexual benefits of losing weight, whether from bariatric surgery or otherwise, will be greater in women than in men.

Future work on the sexuality and relationships of bariatric surgery patients within couples would benefit from comparing them with regular couples, obese couples who underwent no surgery, couples undergoing other comparable life-altering procedures, and of course, cases in which the male, the female, or both partners undergo the procedure.

Finally, two other issues came up: the sex ratio of bariatric patients and the cost of the procedure. First, based on the literature and this study, most patients tend to be female, and hence, most studies are female biased and might not be directly applicable to males. Obesity is female biased in developing countries, but male biased in developed countries⁶⁷. Medical practitioners ought to do more to address this bias and perhaps attract more male patients⁶⁸. Why are more women than men attracted to the procedure? Of all the benefits that we examined and found in the literature, only one, sex and relationships, seems to benefit women more than men. Is it important enough to explain the effect?

Second, in Estonia, the state covers the costs of bariatric surgery for residents. In countries where the government does not pay, the cost weighs heavily on the decision to undergo the

Lozano *et al.* **Bariatric surgery: preparations & quality of life**

procedure⁴⁶ . Whether the procedure is financed individually or by the state, bariatric surgery does reduce the total costs of health care over a person's lifetime⁶⁹ .Public health officials and economists will have to work together and decide the degree to which bariatric surgery will continue to play a role in the battle against obesity⁷⁰ .

In conclusion, bariatric surgery leads to large physical changes, which, in turn, have drastic effects on people's everyday lives, their internal mental states, and their roles in society. Potential patients ought to be better prepared for these changes, have more realistic expectations, and be willing to seek counseling both before and after surgery. Concurrently, medical practitioners ought to be aware of the magnitude of the changes this surgery will bring about in their patients' lives, and help them to prepare for surgery and to adjust to the changes afterwards.

ACKNOWLEDGEMENTS

We thank the participants in the study.

CONFLICTS OF INTEREST

None

AUTHOR CONTRIBUTIONS

Study concept and design: TP and LL; acquisition of data: TP and LL; analysis and interpretation of data: all authors; initial drafting of the manuscript: LL; final writing and critical revision of the manuscript: GAL.

REFERENCES

1. Blüher M. Obesity: global epidemiology and pathogenesis. *Nat Rev Endocrinol*. 2019;15(5):288-298. <https://doi.org/10.1038/s41574-019-0176-8>.
2. Bray GA. Obesity: a time bomb to be defused. *Lancet*. 1998;352(9123):160-161. [https://doi.org/10.1016/S0140-6736\(98\)22029-0](https://doi.org/10.1016/S0140-6736(98)22029-0).
3. Pasquali R, Oriolo C. Obesity and androgens in women. *Frontiers of Hormone Research*. 2019;53:120-134. <https://doi.org/10.1159/000494908>.
4. Ortega FB, Lavie CJ, Blair SN. Obesity and cardiovascular disease. *Circulation Research*. 2016;118(11):1752-1770. <https://doi.org/10.1161/CIRCRESAHA.115.306883>.
5. Peters U, Dixon AE, Forno E. Obesity and asthma. *J Allergy Clin Immunol*. 2018;141(4):1169-1179. <https://doi.org/10.1016/j.jaci.2018.02.004>.
6. Dietz W, Santos-Burgoa C. Obesity and its implications for COVID-19 mortality. *Obesity*. 2020;28(6):1005-1005. <https://doi.org/10.1002/oby.22818>.
7. Meurling IJ, Shea DO, Garvey JF. Obesity and sleep: a growing concern. *Current Opinion in Pulmonary Medicine*. 2019;25(6):602-608. <https://doi.org/10.1097/MCP.0000000000000627>.
8. Suvan JE, Finer N, D'Aiuto F. Periodontal complications with obesity. *Periodontology 2000*. 2018;78(1):98-128. <https://doi.org/10.1111/prd.12239>.
9. Seravalle G, Grassi G. Obesity and hypertension. *Pharmacology Research*. 2017;122:1-7. <https://doi.org/10.1016/j.phrs.2017.05.013>.
10. Kinlen D, Cody D, O'Shea D. Complications of obesity. *QJM: An International Journal of Medicine*. 2018;111(7):437-443. <https://doi.org/10.1093/qjmed/hcx152>.
11. Seidell JC. Obesity in Europe: scaling an epidemic. *Int J Obes*. 1995;19(SUPPL. 3):S1-S4.
12. Shetty P, Schmidhuber J. Introductory lecture: the epidemiology and determinants of obesity in developed and developing countries. *International Journal for Vitamin and Nutrition Research*. 2006;76(4):157-162. <https://doi.org/10.1024/0300-9831.76.4.157>.
13. VanItallie TB. Worldwide epidemiology of obesity. *Pharmacoeconomics*. 1994;5(Suppl 1):1-7. <https://doi.org/10.2165/00019053-199400051-00003>.
14. World Health Organization. Obesity and Overweight. 2020; <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>. Accessed 20 April, 2021.
15. Wirth A, Wabitsch M, Hauner H. The prevention and treatment of obesity. *Deutsches Ärzteblatt International*. 2014;111(42):705-713. <https://doi.org/10.3238/arztebl.2014.0705>.
16. Sharma AM, Iacobellis G. Treatment of obesity: a challenging task. *Contributions to Nephrology*. 2006;151:212-220. <https://doi.org/10.1159/000095331>.
17. Yuksel HS, Şahin FN, Maksimovic N, Drid P, Bianco A. School-Based intervention programs for preventing obesity and promoting physical activity and fitness: a systematic review. *Int J Environ Res Public Health*. 2020;17(1):347. <https://doi.org/10.3390/ijerph17010347>.
18. Gittelsohn J, Trude A. Diabetes and obesity prevention: changing the food environment in low-income settings. *Nutr Rev*. 2017;75(suppl 1):62-69. <https://doi.org/10.1093/nutrit/nuw038>.
19. Speakman JR. Evolutionary perspectives on the obesity epidemic: adaptive, maladaptive, and neutral viewpoints. *Annu Rev Nutr*. 2013;33:289-317. <https://doi.org/10.1146/annurev-nutr-071811-150711>.

Lozano *et al.* **Bariatric surgery: preparations & quality of life**

20. Prentice AM, Hennig BJ, Fulford AJ. Evolutionary origins of the obesity epidemic: Natural selection of thrifty genes or genetic drift following predation release? *Int J Obes.* 2008;32(11):1607-1610. <https://doi.org/10.1038/ijo.2008.147>.
21. Lozano GA. Obesity and sexually selected anorexia nervosa. *Med Hypotheses.* 2008;71:933-940. <https://doi.org/10.1016/j.mehy.2008.07.013>.
22. Ebbeling CB, Feldman HA, Klein GL, *et al.* Effects of a low carbohydrate diet on energy expenditure during weight loss maintenance: randomized trial. *Br Med J.* 2018;363:k4583. <https://doi.org/10.1136/bmj.k4583>.
23. Stewart WK, Fleming LW. Features of a successful therapeutic fast of 382 days' duration. *Postgrad Med J.* 1973;49(569):203-209. <https://doi.org/10.1136/pgmj.49.569.203>.
24. Petridou A, Siopi A, Mougios V. Exercise in the management of obesity. *Metabolism.* 2019;92:163-169. <https://doi.org/10.1016/j.metabol.2018.10.009>.
25. Swift DL, McGee JE, Earnest CP, Carlisle E, Nygard M, Johannsen NM. The effects of exercise and physical activity on weight loss and maintenance. *Prog Cardiovasc Dis.* 2018;61(2):206-213. <https://doi.org/10.1016/j.pcad.2018.07.014>.
26. Castelnuovo G, Pietrabissa G, Manzoni GM, *et al.* Cognitive behavioral therapy to aid weight loss in obese patients: current perspectives. *Psychol Res Behav Manag.* 2017;10:165-173. <https://doi.org/10.2147/PRBM.S113278>.
27. Barrett S, Begg S, O'Halloran P, Kingsley M. Integrated motivational interviewing and cognitive behaviour therapy for lifestyle mediators of overweight and obesity in community-dwelling adults: a systematic review and meta-analyses. *BMC Publ Health.* 2018;18(1):Article number: 1160. <https://doi.org/10.1186/s12889-018-6062-9>.
28. Dragano NRV, Fernø J, Diéguez C, López M, Milbank E. Recent updates on obesity treatments: available drugs and future directions. *Neuroscience.* 2020;437:215-239. <https://doi.org/10.1016/j.neuroscience.2020.04.034>.
29. Kanj A, Levine D. Overcoming obesity: weight-loss drugs are underused. *Cleve Clin J Med.* 2020;87(10):602-604. <https://doi.org/10.3949/ccjm.87a.19102>.
30. Kissler HJ, Settmacher U. Bariatric surgery to treat obesity. *Semin Nephrol.* 2013;33(1):75-89. <https://doi.org/10.1016/j.semnephrol.2012.12.004>.
31. Wolfe BM, Kvach E, Eckel RH. Treatment of obesity: weight loss and bariatric surgery. *Circulation Research.* 2016;118(11):1844-1855. <https://doi.org/10.1161/CIRCRESAHA.116.307591>.
32. Nguyen NT, Varela JE. Bariatric surgery for obesity and metabolic disorders: state of the art. *Nature Reviews Gastroenterology & Hepatology.* 2017;14(3):160-169. <https://doi.org/10.1038/nrgastro.2016.170>.
33. Dumon KR, Murayama KM. Bariatric surgery outcomes. *Surgical Clinics of North America.* 2011;91(6):1313-1338, x. <https://doi.org/10.1016/j.suc.2011.08.014>.
34. Korenkov M. Bariatric surgery. *Contributions to Nephrology.* 2006;151:243-253. <https://doi.org/10.1159/000095334>.
35. Cordero P, Li J, Oben JA. Bariatric surgery as a treatment for metabolic syndrome. *J R Coll Physicians Edinb.* 2017;47(4):364-368. <https://doi.org/10.4997/JRCPE.2017.414>.
36. Griauzde DH, Ibrahim AM, Fisher N, Stricklen A, Ross R, Ghaferi AA. Understanding the psychosocial impact of weight loss following bariatric surgery: a qualitative study. *BMC Obesity.* 2018;5(1):38. <https://doi.org/10.1186/s40608-018-0215-3>.
37. Opozda M, Wittert G, Chur-Hansen A. Patients' expectations and experiences of eating behaviour change after bariatric procedures. *Clin Obes.* 2018;8(5):355-365. <https://doi.org/10.1111/cob.12273>.

Lozano *et al.* **Bariatric surgery: preparations & quality of life**

38. Paas M, Schönberg M. *Bariaatrilise kirurgia patsiendile suunatud õendusabi pre- ja postoperatiivses perioodis [Nursing care for bariatric surgery patients pre- and post surgery]*. Tallinn, Tallinna Tervishoiu Kõrgkool; 2015.
39. Kahr K, Palm R. *Õendusabi patsiendile maovähendusoperatsiooni järgselt [Nursing care after bariatric surgery]*, Tallinna Tervishoiu Kõrgkool; 2015.
40. Tekkel M, Veideman T. Eesti täiskasvanud rahvastiku tervisekäitumise uuring, 2016 [Health Behaviour Among Estonian Adult Population, 2016]. Tallinn: Tervise Arengu Instituut [National Institute for Health Development]; 2017.
41. Estonian Health Insurance Fund. Bariaatrilise patsiendi käsitlemine enne ja pärast kirurgilist sekkumist [Dealing with bariatric surgery patients before and after surgery]. 2022; <https://www.ravijuhend.ee/patsiendivarav/juhendid/209/bariaatrilise-patsiendi-kasitlus-enne-ja-parast-kirurgilist-sekkumist>. Accessed January 31, 2022.
42. Bariatric Services. Facts about bariatric surgery in Estonia. 2020; <https://www.bariatricservices.eu/bariatric-surgery-in-estonia/>. Accessed January 28, 2022.
43. Elo S, Kyngäs H. The qualitative content analysis process. *J Adv Nurs*. 2008;62(1):107-115. <https://doi.org/10.1111/j.1365-2648.2007.04569.x>.
44. Alumäe T, Tilk O, Ullah A. Advanced rich transcription system for Estonian speech. Paper presented at: Human Language Technologies – The Baltic Perspective 2018; Amsterdam.
45. Homer CV, Tod AM, Thompson AR, Allmark P, Goyder E. Expectations and patients' experiences of obesity prior to bariatric surgery: a qualitative study. *BMJ Open*. 2016;6(2):e009389. <https://doi.org/10.1136/bmjopen-2015-009389>.
46. Roberson DW, Neil JA, Pories ML, Rose MA. Tipping point: factors influencing a patient's decision to proceed with bariatric surgery. *Surg Obes Relat Dis*. 2016;12(5):1086-1090. <https://doi.org/10.1016/j.soard.2016.01.009>.
47. Pearl RL, Wadden TA, Walton K, Allison KC, Tronieri JS, Williams NN. Health and appearance: factors motivating the decision to seek bariatric surgery. *Surg Obes Relat Dis*. 2019;15(4):636-642. <https://doi.org/10.1016/j.soard.2019.01.015>.
48. Oliveira DM, Merighi MA, Jesus MC. A decisão da mulher obesa pela cirurgia bariátrica à luz da fenomenologia social [The obese woman's decision for bariatric surgery in light of social phenomenology]. *Rev Esc Enferm USP*. 2014;48(6):970-976. <https://doi.org/10.1590/S0080-623420140000700002>.
49. Cohn I, Raman J, Sui Z. Patient motivations and expectations prior to bariatric surgery: a qualitative systematic review. *Obes Rev*. 2019;20(11):1608-1618. <https://doi.org/10.1111/obr.12919>.
50. Lim R, Beekley A, Johnson DC, Davis KA. Early and late complications of bariatric operation. *Trauma Surg Acute Care Open*. 2018;3(1):e000219. <https://doi.org/10.1136/tsaco-2018-000219>.
51. Stefanik JJ, Felson DT, Apovian CM, *et al.* Changes in pain sensitization after bariatric surgery. *Arthritis Care Res*. 2018;70(10):1525-1528. <https://doi.org/10.1002/acr.23513>.
52. Courcoulas AP, Christian NJ, Belle SH, *et al.* Weight change and health outcomes at 3 years after bariatric surgery among individuals with severe obesity. *JAMA*. 2013;310(22):2416-2425. <https://doi.org/10.1001/jama.2013.280928>.
53. Manzoni APdS, Weber MB. Skin changes after bariatric surgery. *An Bras Dermatol*. 2015;90(2):157-166. <https://doi.org/10.1590/abd1806-4841.20153139>.
54. Blix AS. Adaptations to polar life in mammals and birds. *J Exp Biol*. 2016;219(8):1093-1105. <https://doi.org/10.1242/jeb.120477>.

Lozano *et al.* **Bariatric surgery: preparations & quality of life**

55. Lin HC, Tsao LI. Living with my small stomach: the experiences of post-bariatric surgery patients within 1 year after discharge. *J Clin Nurs*. 2018;27(23-24):4279-4289. <https://doi.org/10.1111/jocn.14616>.
56. Bal BS, Finelli FC, Shope TR, Koch TR. Nutritional deficiencies after bariatric surgery. *Nat Rev Endocrinol*. 2012;8(9):544-556. <https://doi.org/10.1038/nrendo.2012.48>.
57. Sharman M, Hensher M, Wilkinson S, *et al.* What are the support experiences and needs of patients who have received bariatric surgery? *Health Expect*. 2017;20(1):35-46. <https://doi.org/10.1111/hex.12423>.
58. World Health Organization. International Classification of Diseases, 11th Revision. 2019/2021; <https://icd.who.int/browse11/l-m/en>. Accessed 15 September, 2022.
59. Day J, Ternouth A, Collier DA. Eating disorders and obesity: two sides of the same coin? *Epidemiol Psychiatr Sci*. 2009;18(2):96-100. <https://doi.org/10.1017/S1121189X00000956>.
60. Kim D, Seo J, Ha KH, Kim DJ. Maintaining physical activity is associated with reduced major adverse cardiovascular events in people newly diagnosed with diabetes. *J Obes Metab Syndr*. 2022;31(2):187-195. <https://doi.org/10.7570/jomes22007>.
61. King WC, Chen JY, Belle SH, *et al.* Change in pain and physical function following bariatric surgery for severe obesity. *JAMA*. 2016;315(13):1362-1371. <https://doi.org/10.1001/jama.2016.3010>.
62. Lacerda RMR, Castanha CR, Castanha AR, Campos JM, Ferraz Á AB, Vilar L. Perception of body image by patients undergoing bariatric surgery. *Rev Col Bras Cir*. 2018;45(2):e1793. <https://doi.org/10.1590/0100-6991e-20181793>.
63. Benson-Davies S, Davies ML, Kattelman K. Understanding eating and exercise behaviors in post roux-en-y gastric bypass patients: a quantitative and qualitative study. *Bariatr Surg Pract Patient Care*. 2013;8(2):61-68. <https://doi.org/10.1089/bari.2013.9989>.
64. Gokalp F, Koras O, Ugur M, *et al.* Bariatric surgery has positive effects on patients' and their partners' sexual function: a prospective study. *Andrology*. 2021;9(4):1119-1125. <https://doi.org/10.1111/andr.13000>.
65. Goitein D, Zendel A, Segev L, Feigin A, Zippel D. Bariatric surgery improves sexual function in obese patients. *Isr Med Assoc J*. 2015;17(10):616-619.
66. Sarhan MD, Khattab M, Sarhan MD, Maurice KK, Hassan H. Impact of bariatric surgery on male sexual health: a prospective study. *Obes Surg*. 2021;31(9):4064-4069. <https://doi.org/10.1007/s11695-021-05522-7>.
67. Kanter R, Caballero B. Global gender disparities in obesity: a review. *Adv Nutr*. 2012;3(4):491-498. <https://doi.org/10.3945/an.112.002063>.
68. Fuchs HF, Broderick RC, Harnsberger CR, *et al.* Benefits of bariatric surgery do not reach obese men. *J Laparoendosc Adv Surg Tech*. 2015;25(3):196-201. <https://doi.org/10.1089/lap.2014.0639>.
69. Alsumali A, Eguale T, Bairdain S, Samnaliev M. Cost-effectiveness analysis of bariatric surgery for morbid obesity. *Obes Surg*. 2018;28(8):2203-2214. <https://doi.org/10.1007/s11695-017-3100-0>.
70. Lim J, Cho YH, Yamamoto H, Eng A, Markovic T, Kim KK. Governmental or social support of bariatric surgery in the Asia-Pacific region. *J Obes Metab Syndr*. 2017;26(1):10-14. <https://doi.org/10.7570/jomes.2017.26.1.10>.